

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07H 21/04, C07K 14/705, C12N 15/09, 15/63, C12Q 1/68</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/57132</b>																					
			<b>(43) International Publication Date:</b> 11 November 1999 (11.11.99)																					
<b>(21) International Application Number:</b> PCT/US99/09970		<b>(72) Inventors:</b> JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). McCOY, John, M.; 56 Howard Street, Reading, MA 01867 (US). LaVALLIE, Edward, R.; 113 Ann Lee Road, Harvard, MA 01451 (US). COLLINS-RACIE, Lisa, A.; 124 School Street, Acton, MA 01720 (US). EVANS, Cheryl; 18801 Bent Willow Circle, Germantown, MD 20874 (US). MERBERG, David; 2 Orchard Drive, Acton, MA 01720 (US). TREACY, Maurice; 12 Foxrock Court, Dublin 18 (IE). AGOSTINO, Michael, J.; 26 Wolcott Avenue, Andover, MA 01810 (US). STEININGER, Robert, J., II; 100 Reed Street, Cambridge, MA 02140 (US). BOWMAN, Michael, R.; 50 Aldrich Road, Canton, MA 02021 (US). DiBLASIO-SMITH, Elizabeth; 17 Chestnut Road, Tyngsboro, MA 01879 (US). WIDOM, Angela; 19 Cherokee Road, Acton, MA 01720 (US).																						
<b>(22) International Filing Date:</b> 7 May 1999 (07.05.99)		<b>(74) Agent:</b> MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).																						
<b>(30) Priority Data:</b> <table border="0"><tr><td>60/084,564</td><td>7 May 1998 (07.05.98)</td><td>US</td></tr><tr><td>60/087,645</td><td>2 June 1998 (02.06.98)</td><td>US</td></tr><tr><td>60/093,712</td><td>22 July 1998 (22.07.98)</td><td>US</td></tr><tr><td>60/094,935</td><td>31 July 1998 (31.07.98)</td><td>US</td></tr><tr><td>60/095,880</td><td>10 August 1998 (10.08.98)</td><td>US</td></tr><tr><td>60/096,068</td><td>11 August 1998 (11.08.98)</td><td>US</td></tr><tr><td>Not furnished</td><td>6 May 1999 (06.05.99)</td><td>US</td></tr></table>		60/084,564	7 May 1998 (07.05.98)	US	60/087,645	2 June 1998 (02.06.98)	US	60/093,712	22 July 1998 (22.07.98)	US	60/094,935	31 July 1998 (31.07.98)	US	60/095,880	10 August 1998 (10.08.98)	US	60/096,068	11 August 1998 (11.08.98)	US	Not furnished	6 May 1999 (06.05.99)	US	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
60/084,564	7 May 1998 (07.05.98)	US																						
60/087,645	2 June 1998 (02.06.98)	US																						
60/093,712	22 July 1998 (22.07.98)	US																						
60/094,935	31 July 1998 (31.07.98)	US																						
60/095,880	10 August 1998 (10.08.98)	US																						
60/096,068	11 August 1998 (11.08.98)	US																						
Not furnished	6 May 1999 (06.05.99)	US																						
<b>(71) Applicant:</b> GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US).		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>																						
<b>(54) Title:</b> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM																								
<b>(57) Abstract</b>  Novel polynucleotides and the proteins encoded thereby are disclosed.																								

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5           (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366;
- (c)     a polynucleotide comprising the nucleotide sequence of the full-  
10     length protein coding sequence of clone bn365\_53 deposited under accession number ATCC 98752;
- (d)     a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;
- (e)     a polynucleotide comprising the nucleotide sequence of a mature  
15     protein coding sequence of clone bn365\_53 deposited under accession number ATCC 98752;
- (f)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;
- (g)     a polynucleotide encoding a protein comprising the amino acid  
20     sequence of SEQ ID NO:2;
- (h)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
- (i)     a polynucleotide which is an allelic variant of a polynucleotide of  
25     (a)-(f) above;
- (j)     a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- (k)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- 30     (l)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:1.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366; the nucleotide sequence of the full-length

## 5           SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

This application is a continuation-in-part of the following applications:

- 10           (1)     provisional application Ser. No. 60/084,564, filed May 7, 1998;  
            (2)     provisional application Ser. No. 60/087,645, filed June 2, 1998;  
            (3)     provisional application Ser. No. 60/093,712, filed July 22, 1998;  
            (4)     provisional application Ser. No. 60/094,935, filed July 31, 1998;  
            (5)     provisional application Ser. No. 60/095,880, filed August 10, 1998;  
            (6)     provisional application Ser. No. 60/096,068, filed August 11, 1998;  
15   all of which are incorporated by reference herein.

FIELD OF THE INVENTION

20       The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins.

BACKGROUND OF THE INVENTION

25       Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by  
35   virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins and the polynucleotides encoding them that the present invention is directed.



protein coding sequence of clone bn365\_53 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone bn365\_53 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752. In further preferred  
5 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a polynucleotide encoding  
10 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:2.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:1.

15 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
20 consisting of:

(aa) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

(ab) the nucleotide sequence of the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;

25 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

30 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1; and

(bb) the nucleotide sequence of the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:1, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:1 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:1, but excluding the poly(A) tail at the 3' end of SEQ ID NO:1. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the  
15 cDNA sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366.

20 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

25 (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and

(c) the amino acid sequence encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:2. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:2, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:2.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:3 from nucleotide 1358 to nucleotide 1915;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752;
- (e) a polynucleotide encoding the full-length protein encoded by the  
15 cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
20 insert of clone bo342\_2 deposited under accession number ATCC 98752;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:4;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:3.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915; the nucleotide sequence of SEQ ID NO:3 from nucleotide 1358 to nucleotide 1915; the nucleotide sequence of the full-length protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising the amino acid sequence from amino acid 280 to amino acid 289 of SEQ ID NO:4.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:3.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and
    - (ab) the nucleotide sequence of the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3; and

(bb) the nucleotide sequence of the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:3 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:3, but excluding the poly(A) tail at the 3' end of SEQ ID NO:3. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:3 from nucleotide 1358 to nucleotide 1915, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:3 from nucleotide 1358 to nucleotide 1915, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:3 from nucleotide 1358 to nucleotide 1915.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of SEQ  
ID NO:4 having biological activity, the fragment comprising the amino acid sequence from  
10 amino acid 280 to amino acid 289 of SEQ ID NO:4.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5;

15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dn721\_8 deposited under accession number ATCC 98752;

20 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dn721\_8 deposited under accession number ATCC 98752;

25 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;

30 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:5.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689; the nucleotide sequence of the full-length  
10 protein coding sequence of clone dn721\_8 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone dn721\_8 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752. In further preferred  
15 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having  
20 biological activity, the fragment comprising the amino acid sequence from amino acid 318 to amino acid 327 of SEQ ID NO:6.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:5.

Further embodiments of the invention provide isolated polynucleotides produced  
25 according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (aa) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(ab) the nucleotide sequence of the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

5 and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (ba) SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5; and

(bb) the nucleotide sequence of the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

15 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5, and extending  
20 contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:5 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:5, but excluding the poly(A) tail at the 3' end of SEQ ID NO:5. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689, and extending  
25 contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and



(c) the amino acid sequence encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred  
5 -- embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising the amino acid sequence from  
10 amino acid 318 to amino acid 327 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 20 to nucleotide 484;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 18 to nucleotide 892;
- 20 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dn834\_1 deposited under accession number ATCC 98752;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
25 protein coding sequence of clone dn834\_1 deposited under accession number ATCC 98752;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:8;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:7.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 20 to nucleotide 484; the nucleotide sequence of SEQ ID NO:7 from nucleotide 18 to nucleotide 892; the nucleotide sequence of the full-length protein coding sequence of clone dn834\_1 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone dn834\_1 deposited  
15 under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 72 to amino acid 81 of SEQ ID NO:8.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:7.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and

(ab) the nucleotide sequence of the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

5 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

10 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7; and

15 (bb) the nucleotide sequence of the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:7 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:7, but excluding the poly(A) tail at the 3' end of SEQ ID NO:7. Also preferably the polynucleotide isolated  
25 according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 20 to nucleotide 484, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide 20 to nucleotide 484, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 20 to  
30 nucleotide 484. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:7 from nucleotide 18 to nucleotide 892, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:7 from nucleotide

18 to nucleotide 892, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:7 from nucleotide 18 to nucleotide 892.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 72 to amino acid 81 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;

5 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

10 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

15 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:9.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420; the nucleotide sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420; the nucleotide sequence of the full-length  
20 protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752. In further preferred  
25 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having  
30 biological activity, the fragment comprising the amino acid sequence from amino acid 98 to amino acid 107 of SEQ ID NO:10.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:9.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(ab) the nucleotide sequence of the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9; and

(bb) the nucleotide sequence of the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:9 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:9, but excluding the poly(A) tail at the 3' end of SEQ ID NO:9. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420, and extending

contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420. Also preferably the polynucleotide isolated according to the above  
5 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:10;

(b) a fragment of the amino acid sequence of SEQ ID NO:10, the  
15 fragment comprising eight contiguous amino acids of SEQ ID NO:10; and

(c) the amino acid sequence encoded by the cDNA insert of clone  
pd278\_5 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising the amino acid sequence  
25 from amino acid 98 to amino acid 107 of SEQ ID NO:10.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID  
NO:11;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:11 from nucleotide 918 to nucleotide 1295;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pe80\_1 deposited under accession number  
ATCC 98752;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pe80\_1 deposited under accession number ATCC 98752;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:11.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 918 to nucleotide 1295; the nucleotide sequence of the full-length protein coding sequence of clone pe80\_1 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone pe80\_1 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having



biological activity, the fragment comprising the amino acid sequence from amino acid 58 to amino acid 67 of SEQ ID NO:12.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:11.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
10 consisting of:

(aa) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(ab) the nucleotide sequence of the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11; and

(bb) the nucleotide sequence of the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:11 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:11, but excluding the poly(A) tail at the 3' end of SEQ ID NO:11. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:11 from nucleotide 918 to nucleotide 1295, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:11 from nucleotide 918 to nucleotide 1295, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:11 from nucleotide 918 to nucleotide 1295.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
- (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising the amino acid sequence from amino acid 58 to amino acid 67 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 348 to nucleotide 428;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:13.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428; the nucleotide sequence of SEQ ID NO:13 from nucleotide 348 to nucleotide 428; the nucleotide sequence of the full-length protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 35  
5 to amino acid 44 of SEQ ID NO:14.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:13.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 10 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 15 (aa) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and
    - (ab) the nucleotide sequence of the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 20 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
25 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:13, but excluding the poly(A) tail at the 3' end of SEQ ID NO:13; and
    - (bb) the nucleotide sequence of the cDNA insert of clone  
30 pm113\_1 deposited under accession number ATCC 98752;
  - (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
  - (iii) amplifying human DNA sequences; and
  - (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:13 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:13, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:13. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:13 from nucleotide 348 to nucleotide 428, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:13 from  
15 nucleotide 348 to nucleotide 428, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:13 from nucleotide 348 to nucleotide 428.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:14;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;
- 25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:14. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
30 of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:14.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:15 from nucleotide 108 to nucleotide 1496;

5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm749\_8 deposited under accession number ATCC 98752;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;

10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm749\_8 deposited under accession number ATCC 98752;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;

15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:16;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;

20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:15.

30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:15 from nucleotide 108 to nucleotide 1496; the nucleotide sequence of the full-length protein coding sequence of clone pm749\_8 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone pm749\_8 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

of clone pm749\_8 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 226 to amino acid 235 of SEQ ID NO:16.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:15.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(ab) the nucleotide sequence of the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15; and

(bb) the nucleotide sequence of the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

- 5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:15 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:15, but excluding the poly(A) tail at the 3' end of SEQ ID NO:15. Also preferably the
- 10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:15 from nucleotide 108 to nucleotide 1496, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:15 from nucleotide 108 to nucleotide 1496, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:15 from nucleotide
- 15 108 to nucleotide 1496.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
- 20 (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;
- the protein being substantially free from other mammalian proteins. Preferably such
- 25 protein comprises the amino acid sequence of SEQ ID NO:16. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:16, or a protein comprising a fragment of the amino acid sequence of SEQ
- 30 ID NO:16 having biological activity, the fragment comprising the amino acid sequence from amino acid 226 to amino acid 235 of SEQ ID NO:16.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:



(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023;

5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752;

10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752;

15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:18;

20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:17.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023; the nucleotide sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023; the nucleotide sequence of the full-length protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752; or the

nucleotide sequence of a mature protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 325 to amino acid 334 of SEQ ID NO:18.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:17.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and

(ab) the nucleotide sequence of the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17; and

- (bb) the nucleotide sequence of the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:17 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:17, but excluding the poly(A) tail at the 3' end of SEQ ID NO:17. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
- (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:18. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:18, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18 having biological activity, the fragment comprising the amino acid sequence from amino acid 325 to amino acid 334 of SEQ ID NO:18.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19;

10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pv296\_5 deposited under accession number ATCC 98752;

15 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pv296\_5 deposited under accession number ATCC 98752;

20 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:20;

25 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

30 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:19.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299; the nucleotide sequence of the full-length protein coding sequence of clone pv296\_5 deposited under accession number ATCC 98752; or the nucleotide sequence of a mature protein coding sequence of clone pv296\_5 deposited under accession number ATCC 98752. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:20.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:19.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and

(ab) the nucleotide sequence of the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19; and

(bb) the nucleotide sequence of the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:19 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:19, but excluding the poly(A) tail at the 3' end of SEQ ID NO:19. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:20;

(b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and

(c) the amino acid sequence encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:20. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:20, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20 having biological activity, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:20.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;

10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone er311\_20 deposited under accession number ATCC 98781;

15 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone er311\_20 deposited under accession number ATCC 98781;

20 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;

25 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

30 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:21.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008; the nucleotide sequence of the full-length protein coding sequence of clone er311\_20 deposited under accession number ATCC 98781; or the nucleotide sequence of a mature protein coding sequence of clone er311\_20 deposited under accession number ATCC 98781. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 328 to amino acid 337 of SEQ ID NO:22.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:21.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

(ab) the nucleotide sequence of the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:



(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21; and

(bb) the nucleotide sequence of the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:21 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:21, but excluding the poly(A) tail at the 3' end of SEQ ID NO:21. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:22;

(b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and

(c) the amino acid sequence encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising the amino acid sequence from amino acid 328 to amino acid 337 of SEQ ID NO:22.

5 In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23;

10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043;

15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;

20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;

25 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

30 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:23.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043; the nucleotide sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043; the nucleotide sequence of the full-length protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781; or the nucleotide sequence of a mature protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:24, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 255 to amino acid 264 of SEQ ID NO:24.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:23.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and
    - (ab) the nucleotide sequence of the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23; and

(bb) the nucleotide sequence of the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:23 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:23, but excluding the poly(A) tail at the 3' end of SEQ ID NO:23. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:24;

(b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and

(c) the amino acid sequence encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:24. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
10 of SEQ ID NO:24, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 255 to amino acid 264 of SEQ ID NO:24.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 47 to nucleotide 1099;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781;

30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:25.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:25 from nucleotide 47 to nucleotide 1099; the nucleotide sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099; the nucleotide sequence of the full-length protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781; or the nucleotide sequence of a mature protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:26, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 170 to amino acid 179 of SEQ ID NO:26.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:25.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(ab) the nucleotide sequence of the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25; and

(bb) the nucleotide sequence of the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:25 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:25, but excluding the poly(A) tail at the 3' end of SEQ ID NO:25. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide 47 to nucleotide 1099, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from nucleotide 47 to nucleotide 1099, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide

47 to nucleotide 1099. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:26;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:26. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 20 of SEQ ID NO:26, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising the amino acid sequence from amino acid 170 to amino acid 179 of SEQ ID NO:26.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;
  - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259;
  - (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pl87\_1 deposited under accession number ATCC 98781;
- 30 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;



(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pl87\_1 deposited under accession number ATCC 98781;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:27.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259; the nucleotide sequence of the full-length protein coding sequence of clone pl87\_1 deposited under accession number ATCC 98781; or the nucleotide sequence of a mature protein coding sequence of clone pl87\_1 deposited under accession number ATCC 98781. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:28.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:27.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

(ab) the nucleotide sequence of the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

20 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27; and

25 (bb) the nucleotide sequence of the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:27 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:27, but excluding the poly(A) tail at the 3' end of SEQ ID NO:27. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:28;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 20 of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 37 to amino acid 46 of SEQ ID NO:28.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm514\_4 deposited under accession number ATCC 98781;
- 30 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm514\_4 deposited under accession number ATCC 98781;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:30;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:29.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284; the nucleotide sequence of the full-length protein coding sequence of clone pm514\_4 deposited under accession number ATCC 98781; or the nucleotide sequence of a mature protein coding sequence of clone pm514\_4 deposited under accession number ATCC 98781. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 365 to amino acid 374 of SEQ ID NO:30.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:29.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10 (aa) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and

(ab) the nucleotide sequence of the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

20 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29; and

25 (bb) the nucleotide sequence of the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:29 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:29, but excluding the poly(A) tail at the 3' end of SEQ ID NO:29. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:30. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:30, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30 having biological activity, the fragment comprising the amino acid sequence from amino acid 365 to amino acid 374 of SEQ ID NO:30.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:32;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:31.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997; the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997; the nucleotide sequence of the full-length protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:32, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having

biological activity, the fragment comprising the amino acid sequence from amino acid 322 to amino acid 331 of SEQ ID NO:32.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:31.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:31, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:31; and

(ab) the nucleotide sequence of the cDNA insert of clone  
co155\_12 deposited under accession number ATCC 98808;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:31, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:31; and

(bb) the nucleotide sequence of the cDNA insert of clone  
co155\_12 deposited under accession number ATCC 98808;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ



ID NO:31 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:31, but excluding the poly(A) tail at the 3' end of SEQ ID NO:31. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;
- (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
- (c) the amino acid sequence encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:32. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:32, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32 having biological activity, the fragment comprising the amino acid sequence from amino acid 322 to amino acid 331 of SEQ ID NO:32.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 21 to nucleotide 1343;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343;

5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:34;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;

20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:33.

30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:33 from nucleotide 21 to nucleotide 1343; the nucleotide sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343; the nucleotide sequence of the full-length protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808. In other preferred embodiments, the

polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:34, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 215 to amino acid 224 of SEQ ID NO:34.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:33.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and

(ab) the nucleotide sequence of the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33; and

(bb) the nucleotide sequence of the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:33 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:33, but excluding the poly(A) tail at the 3' end of SEQ ID NO:33. Also preferably the  
10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide 21 to nucleotide 1343, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 21 to nucleotide 1343, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:33 from nucleotide  
15 21 to nucleotide 1343. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343, to a nucleotide sequence corresponding to the 3' end of  
20 said sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
- 25 (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such  
30 protein comprises the amino acid sequence of SEQ ID NO:34. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:34, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 215 to amino acid 224 of SEQ ID NO:34.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide 557;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
10 NO:35 from nucleotide 235 to nucleotide 899;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone lv2\_47 deposited under accession number ATCC 98808;
- (e) a polynucleotide encoding the full-length protein encoded by the  
15 cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone lv2\_47 deposited under accession number ATCC 98808;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA  
20 insert of clone lv2\_47 deposited under accession number ATCC 98808;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment  
25 comprising eight contiguous amino acids of SEQ ID NO:36;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:35.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide 557; the nucleotide sequence of SEQ ID NO:35 from nucleotide 235 to nucleotide 899; the nucleotide sequence of the full-length protein coding sequence of clone lv2\_47 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone lv2\_47 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:36 from amino acid 58 to amino acid 164. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 77 to amino acid 86 of SEQ ID NO:36.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:35.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and
    - (ab) the nucleotide sequence of the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

5 (ba) SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35; and

(bb) the nucleotide sequence of the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;

10 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:35 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:35, but excluding the poly(A) tail at the 3' end of SEQ ID NO:35. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide  
20 557, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide 557, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide 557. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
25 NO:35 from nucleotide 235 to nucleotide 899, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:35 from nucleotide 235 to nucleotide 899, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:35 from nucleotide 235 to nucleotide 899.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:36;

(b) the amino acid sequence of SEQ ID NO:36 from amino acid 58 to amino acid 164;

(c) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and

(d) the amino acid sequence encoded by the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:36 or the amino acid sequence of SEQ ID NO:36 from amino acid 58 to amino acid 164. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment preferably  
10 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:36, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36 having biological activity, the fragment comprising the amino acid sequence from amino acid 77 to amino acid 86 of SEQ ID NO:36.

In one embodiment, the present invention provides a composition comprising an  
15 isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 104 to nucleotide 499;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:37 from nucleotide 215 to nucleotide 499;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808;

30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:38;



(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:37.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:37 from nucleotide 104 to nucleotide 499; the nucleotide sequence of SEQ ID NO:37 from nucleotide 215 to nucleotide 499; the nucleotide sequence of the full-length protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:38, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:38.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:37.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and

(ab) the nucleotide sequence of the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37; and

(bb) the nucleotide sequence of the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:37 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:37, but excluding the poly(A) tail at the 3' end of SEQ ID NO:37. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide 104 to nucleotide 499, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from nucleotide 104 to nucleotide 499, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide

104 to nucleotide 499. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:37 from nucleotide 215 to nucleotide 499, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:37 from  
5 nucleotide 215 to nucleotide 499, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:37 from nucleotide 215 to nucleotide 499.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:38;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:38. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:38, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:38.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pm96\_9 deposited under accession  
30 number ATCC 98808;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pm96\_9 deposited under accession number ATCC 98808;

5 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:40;

10 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

15 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:39.

20 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861; the nucleotide sequence of the full-length protein coding sequence of clone pm96\_9 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone pm96\_9 deposited under accession number ATCC 98808. In other preferred embodiments, the  
25 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
30 preferably thirty) contiguous amino acids of SEQ ID NO:40, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 110 to amino acid 119 of SEQ ID NO:40.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:39.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 5 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 10 (aa) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
- (ab) the nucleotide sequence of the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 15 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 20 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 25 pm96\_9 deposited under accession number ATCC 98808;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

- 30 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:39 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:39, but excluding the poly(A) tail at the 3' end of SEQ ID NO:39. Also preferably the

polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment comprising the amino acid sequence from amino acid 110 to amino acid 119 of SEQ ID NO:40.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:42;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:41.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762; the nucleotide sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762; the nucleotide sequence of the full-length protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having

biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:42.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:41.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:41, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:41; and

(ab) the nucleotide sequence of the cDNA insert of clone  
pu261\_1 deposited under accession number ATCC 98808;

15 (ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the  
probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that  
hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

(ba) SEQ ID NO:41, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:41; and

(bb) the nucleotide sequence of the cDNA insert of clone  
pu261\_1 deposited under accession number ATCC 98808;

(ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41, and  
extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ



ID NO:41 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:41, but excluding the poly(A) tail at the 3' end of SEQ ID NO:41. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
- (b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:42. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:42, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42 having biological activity, the fragment comprising the amino acid sequence from amino acid 115 to amino acid 124 of SEQ ID NO:42.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pw214\_15 deposited under accession number ATCC 98808;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pw214\_15 deposited under accession number ATCC 98808;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:44;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:43.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824; the nucleotide sequence of the full-length protein coding sequence of clone pw214\_15 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone pw214\_15 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having  
5 biological activity, the fragment comprising the amino acid sequence from amino acid 36 to amino acid 45 of SEQ ID NO:44.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:43.

Further embodiments of the invention provide isolated polynucleotides produced  
10 according to a process selected from the group consisting of:

(a) a process comprising the steps of:  
(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

15 (aa) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

(ab) the nucleotide sequence of the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43; and

30 (bb) the nucleotide sequence of the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:43 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:43, but excluding the poly(A) tail at the 3' end of SEQ ID NO:43. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:44;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:44. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:44, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44 having biological activity, the fragment comprising the amino acid sequence from amino acid 36 to amino acid 45 of SEQ ID NO:44.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qb56\_19 deposited under accession number ATCC 98808;

5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qb56\_19 deposited under accession number ATCC 98808;

10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:46;

15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:45.

25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383; the nucleotide sequence of the full-length protein coding sequence of clone qb56\_19 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone qb56\_19 deposited under accession number ATCC 98808. In other preferred embodiments, the  
30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:46, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 58 to amino acid 67 of SEQ ID NO:46.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:45.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

15 (ab) the nucleotide sequence of the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

20 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45; and

(bb) the nucleotide sequence of the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;

30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:45 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:45, but excluding the poly(A) tail at the 3' end of SEQ ID NO:45. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:46. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:46, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising the amino acid sequence from amino acid 58 to amino acid 67 of SEQ ID NO:46.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide 1273;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 242 to nucleotide 1273;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:48;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:47.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide 1273; the nucleotide sequence of SEQ ID NO:47 from nucleotide 242 to nucleotide 1273; the nucleotide sequence of the full-length protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808. In further preferred



embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a polynucleotide encoding  
5 a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 179 to amino acid 188 of SEQ ID NO:48.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:47.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
15 consisting of:

- (aa) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and

- (ab) the nucleotide sequence of the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47; and

- (bb) the nucleotide sequence of the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:47 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:47, but excluding the poly(A) tail at the 3' end of SEQ ID NO:47. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide  
10 1273, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide 1273, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide 1273. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:47 from nucleotide 242 to nucleotide 1273, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:47 from nucleotide 242 to nucleotide 1273, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:47 from nucleotide 242 to nucleotide 1273.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
- (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48. In further preferred embodiments, the present invention provides a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising the amino acid sequence from amino acid 179 to amino acid 188 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qf116\_2 deposited under accession number ATCC 98808;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qf116\_2 deposited under accession number ATCC 98808;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least
- 30 25% of the length of SEQ ID NO:49.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097; the nucleotide sequence of the full-length protein coding sequence of clone qf116\_2 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone qf116\_2

deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808. In further preferred  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 147  
10 to amino acid 156 of SEQ ID NO:50.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:49.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - 20 (aa) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and
    - (ab) the nucleotide sequence of the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - 25 (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
  - (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (ba) SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49; and

- (bb) the nucleotide sequence of the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

10 ID NO:49 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:49, but excluding the poly(A) tail at the 3' end of SEQ ID NO:49. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097, and extending contiguously from a nucleotide sequence corresponding to the 5' end

15 of said sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group

20 consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
- (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
- (c) the amino acid sequence encoded by the cDNA insert of clone
- 25 qf116\_2 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:50. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment preferably

30 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:50, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50 having biological activity, the fragment comprising the amino acid sequence from amino acid 147 to amino acid 156 of SEQ ID NO:50.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:51 from nucleotide 595 to nucleotide 741;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:51.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741; the nucleotide sequence of SEQ ID NO:51

from nucleotide 595 to nucleotide 741; the nucleotide sequence of the full-length protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808; or the nucleotide sequence of a mature protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 92 to amino acid 101 of SEQ ID NO:52.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:51.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and

(ab) the nucleotide sequence of the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51; and

(bb) the nucleotide sequence of the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

10 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:51 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:51, but excluding the poly(A) tail at the 3' end of SEQ ID NO:51. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741. Also preferably the polynucleotide isolated according to the above  
20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:51 from nucleotide 595 to nucleotide 741, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:51 from nucleotide 595 to nucleotide 741, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:51 from nucleotide 595 to nucleotide 741.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:52;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and

(c) the amino acid sequence encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:52. In further preferred



embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:52, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52 having biological activity, the fragment comprising the amino acid sequence from amino acid 92 to amino acid 101 of SEQ ID NO:52.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:54;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:53.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196; the nucleotide sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196; the nucleotide sequence of the full-length protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 40 to amino acid 49 of SEQ ID NO:54.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:53.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and

(ab) the nucleotide sequence of the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53; and

(bb) the nucleotide sequence of the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:53 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:53, but excluding the poly(A) tail at the 3' end of SEQ ID NO:53. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:54;
- (b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
5 am748\_5 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:54. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment preferably  
10 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:54, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54 having biological activity, the fragment comprising the amino acid sequence from amino acid 40 to amino acid 49 of SEQ ID NO:54.

In one embodiment, the present invention provides a composition comprising an  
15 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 51 to nucleotide 1310;
- 20 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cj507\_1 deposited under accession number ATCC 98817;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817;
- 25 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cj507\_1 deposited under accession number ATCC 98817;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817;
- 30 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:55.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 51 to nucleotide 1310; the nucleotide sequence of the full-length protein coding sequence of clone cj507\_1 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone cj507\_1 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:56, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 205 to amino acid 214 of SEQ ID NO:56.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:55.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:55, but excluding the poly(A) tail at the 3' end of SEQ ID NO:55; and

- (ab) the nucleotide sequence of the cDNA insert of clone  
cj507\_1 deposited under accession number ATCC 98817;  
(ii) hybridizing said probe(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C; and  
5 (iii) isolating the DNA polynucleotides detected with the  
probe(s);

and

- (b) a process comprising the steps of:  
(i) preparing one or more polynucleotide primers that  
10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from  
the group consisting of:

- (ba) SEQ ID NO:55, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:55; and  
(bb) the nucleotide sequence of the cDNA insert of clone  
15 cj507\_1 deposited under accession number ATCC 98817;  
(ii) hybridizing said primer(s) to human genomic DNA in  
conditions at least as stringent as 4X SSC at 50 degrees C;  
(iii) amplifying human DNA sequences; and  
(iv) isolating the polynucleotide products of step (b)(iii).

- 20 Preferably the polynucleotide isolated according to the above process comprises a  
nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:55, and  
extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ  
ID NO:55 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:55, but  
excluding the poly(A) tail at the 3' end of SEQ ID NO:55. Also preferably the  
25 polynucleotide isolated according to the above process comprises a nucleotide sequence  
corresponding to the cDNA sequence of SEQ ID NO:55 from nucleotide 51 to nucleotide  
1310, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
of said sequence of SEQ ID NO:55 from nucleotide 51 to nucleotide 1310, to a nucleotide  
sequence corresponding to the 3' end of said sequence of SEQ ID NO:55 from nucleotide  
30 51 to nucleotide 1310.

In other embodiments, the present invention provides a composition comprising  
a protein, wherein said protein comprises an amino acid sequence selected from the group  
consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;

(b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and

(c) the amino acid sequence encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
10 of SEQ ID NO:56, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 205 to amino acid 214 of SEQ ID NO:56.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:57 from nucleotide 195 to nucleotide 1328;

20 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cn922\_5 deposited under accession number ATCC 98817;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;

25 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cn922\_5 deposited under accession number ATCC 98817;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;

30 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:58;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

5 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:57.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:57 from nucleotide 195 to nucleotide 1328; the nucleotide sequence of the full-length protein coding sequence of clone cn922\_5 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone cn922\_5 deposited under accession number ATCC 98817. In other preferred embodiments, the  
15 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
20 preferably thirty) contiguous amino acids of SEQ ID NO:58, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 184 to amino acid 193 of SEQ ID NO:58.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:57.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and



- (ab) the nucleotide sequence of the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 cn922\_5 deposited under accession number ATCC 98817;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:57 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:57, but excluding the poly(A) tail at the 3' end of SEQ ID NO:57. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:57 from nucleotide 195 to nucleotide 1328, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:57 from nucleotide 195 to nucleotide 1328, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:57 from nucleotide
- 30 195 to nucleotide 1328.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:58;

- (b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:58. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:58, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58 having biological activity, the fragment comprising the amino acid sequence from amino acid 184 to amino acid 193 of SEQ ID NO:58.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:59 from nucleotide 76 to nucleotide 942;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cw691\_11 deposited under accession
- 20 number ATCC 98817;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;
- (e) a polynucleotide comprising the nucleotide sequence of a mature
- 25 protein coding sequence of clone cw691\_11 deposited under accession number ATCC 98817;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;
- (g) a polynucleotide encoding a protein comprising the amino acid
- 30 sequence of SEQ ID NO:60;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:59.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:59 from nucleotide 76 to nucleotide 942; the nucleotide sequence of the full-length protein coding sequence of clone cw691\_11 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone cw691\_11 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:60, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 139 to amino acid 148 of SEQ ID NO:60.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:59.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and

- (ab) the nucleotide sequence of the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 cw691\_11 deposited under accession number ATCC 98817;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:59 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:59, but excluding the poly(A) tail at the 3' end of SEQ ID NO:59. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:59 from nucleotide 76 to nucleotide 942, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:59 from nucleotide 76 to nucleotide 942, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:59 from nucleotide
- 30 76 to nucleotide 942.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;

(b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and

(c) the amino acid sequence encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;

- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:60. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:60, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60 having biological activity, the fragment comprising the amino acid sequence from amino acid 139 to amino acid 148 of SEQ ID NO:60.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 11 to nucleotide 1252;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 1252;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817;

30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:62;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:61.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:61 from nucleotide 11 to nucleotide 1252; the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 1252; the nucleotide sequence of the full-length protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:62, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 202 to amino acid 211 of SEQ ID NO:62.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:61.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

(ab) the nucleotide sequence of the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61; and

(bb) the nucleotide sequence of the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:61 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:61, but excluding the poly(A) tail at the 3' end of SEQ ID NO:61. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 11 to nucleotide 1252, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from nucleotide 11 to nucleotide 1252, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide

11 to nucleotide 1252. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 1252, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:61 from  
5 nucleotide 119 to nucleotide 1252, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 1252.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:62;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:62. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:62, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62 having biological activity, the fragment comprising the amino acid sequence from amino acid 202 to amino acid 211 of SEQ ID NO:62.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:63 from nucleotide 451 to nucleotide 1296;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817;



(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:64;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:63.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296; the nucleotide sequence of SEQ ID NO:63 from nucleotide 451 to nucleotide 1296; the nucleotide sequence of the full-length protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:64, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having

biological activity, the fragment comprising the amino acid sequence from amino acid 203 to amino acid 212 of SEQ ID NO:64.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:63.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:63, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:63; and

(ab) the nucleotide sequence of the cDNA insert of clone  
cw1640\_1 deposited under accession number ATCC 98817;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:63, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:63; and

(bb) the nucleotide sequence of the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:63 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:63, but excluding the poly(A) tail at the 3' end of SEQ ID NO:63. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:63 from nucleotide 451 to nucleotide 1296, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:63 from nucleotide 451 to nucleotide 1296, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:63 from nucleotide 451 to nucleotide 1296.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:64;
- (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:64. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:64, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising the amino acid sequence from amino acid 203 to amino acid 212 of SEQ ID NO:64.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827;

5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;

10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;

15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:66;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;

20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:65.

30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827; the nucleotide sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827; the nucleotide sequence of the full-length protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide

encodes the full-length or a mature protein encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:66.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:65.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65; and

(ab) the nucleotide sequence of the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65; and

(bb) the nucleotide sequence of the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:65 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:65, but excluding the poly(A) tail at the 3' end of SEQ ID NO:65. Also preferably the  
10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:65 from nucleotide  
15 66 to nucleotide 827. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827, to a nucleotide sequence corresponding to the 3' end of  
20 said sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:66;

25 (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and

(c) the amino acid sequence encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins. Preferably such  
30 protein comprises the amino acid sequence of SEQ ID NO:66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:66, or a protein comprising a fragment of the amino acid sequence of SEQ

ID NO:66 having biological activity, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:66.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

5           (a)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67;

          (b)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529;

10          (c)     a polynucleotide comprising the nucleotide sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529;

          (d)     a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817;

15          (e)     a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;

          (f)     a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817;

20          (g)     a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;

          (h)     a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:68;

25          (i)     a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;

          (j)     a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

          (k)     a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

30          (l)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

          (m)     a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:67.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529; the nucleotide sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529; the nucleotide sequence of the full-length protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence from amino acid 58 to amino acid 67 of SEQ ID NO:68.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:67.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and
    - (ab) the nucleotide sequence of the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:



(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67; and

(bb) the nucleotide sequence of the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:67 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:67, but excluding the poly(A) tail at the 3' end of SEQ ID NO:67. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:68;

(b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and

(c) the amino acid sequence encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:68. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:68, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68 having biological activity, the fragment comprising the amino acid sequence  
10 from amino acid 58 to amino acid 67 of SEQ ID NO:68.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 31 to nucleotide 543;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:69 from nucleotide 88 to nucleotide 543;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone di393\_2 deposited under accession  
20 number ATCC 98817;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817;
- (f) a polynucleotide comprising the nucleotide sequence of a mature  
25 protein coding sequence of clone di393\_2 deposited under accession number ATCC 98817;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817;
- (h) a polynucleotide encoding a protein comprising the amino acid  
30 sequence of SEQ ID NO:70;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

5 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:69.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:69 from nucleotide 31 to nucleotide 543; the nucleotide sequence of SEQ ID NO:69 from nucleotide 88 to nucleotide 543; the nucleotide sequence of the full-length protein coding sequence of clone di393\_2 deposited under accession number ATCC 98817; or the nucleotide sequence of a mature protein coding sequence of clone di393\_2 deposited  
15 under accession number ATCC 98817. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological  
20 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 80 to amino acid 89 of SEQ ID NO:70.

25 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:69.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and

- (ab) the nucleotide sequence of the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 di393\_2 deposited under accession number ATCC 98817;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).
- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:69 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:69, but excluding the poly(A) tail at the 3' end of SEQ ID NO:69. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide 31 to nucleotide 543, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from nucleotide 31 to nucleotide 543, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide
- 30 31 to nucleotide 543. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:69 from nucleotide 88 to nucleotide 543, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:69 from

nucleotide 88 to nucleotide 543, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:69 from nucleotide 88 to nucleotide 543.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
5 consisting of:

- (a) the amino acid sequence of SEQ ID NO:70;
- (b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
10 di393\_2 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:70. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment preferably  
15 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:70, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70 having biological activity, the fragment comprising the amino acid sequence from amino acid 80 to amino acid 89 of SEQ ID NO:70.

In one embodiment, the present invention provides a composition comprising an  
20 isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dj167\_2 deposited under accession  
25 number ATCC 98818;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dj167\_2 deposited under accession number  
30 ATCC 98818;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:72;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:71.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356; the nucleotide sequence of the full-length protein coding sequence of clone dj167\_2 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone dj167\_2 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 195 to amino acid 204 of SEQ ID NO:72.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:71.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

(ab) the nucleotide sequence of the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71; and

(bb) the nucleotide sequence of the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:71 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:71, but excluding the poly(A) tail at the 3' end of SEQ ID NO:71. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356, to a nucleotide

sequence corresponding to the 3' end of said sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:72;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:72. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:72, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72 having biological activity, the fragment comprising the amino acid sequence from amino acid 195 to amino acid 204 of SEQ ID NO:72.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490;
- (d) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343;
- (e) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090;
- (f) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;



(g) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090;

(h) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;

(i) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:74;

(j) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;

(k) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(h) above;

(l) a polynucleotide which encodes a species homologue of the protein of (i) or (j) above ;

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j); and

(n) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(j) and that has a length that is at least 25% of the length of SEQ ID NO:73.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490; the nucleotide sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490; the nucleotide sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343; the nucleotide sequence of the full-length protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090; or the nucleotide sequence of a mature protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090. In yet other preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:74 from amino acid 637 to amino acid 1036. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or

a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence from amino acid 513 to amino acid 522 of SEQ ID NO:74.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:73.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

(ab) the nucleotide sequence of the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73; and

(bb) the nucleotide sequence of the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73, and

extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:73 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:73, but excluding the poly(A) tail at the 3' end of SEQ ID NO:73. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
5 corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490. Also preferably the polynucleotide isolated according to the  
10 above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490. Also  
15 preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343, to a nucleotide sequence corresponding to the 3' end of said  
20 sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- 25 (b) the amino acid sequence of SEQ ID NO:74 from amino acid 637 to amino acid 1036;
- (c) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
- (d) the amino acid sequence encoded by the cDNA insert of clone  
30 dj167\_19 deposited under accession number ATCC 207090;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:74 or the amino acid sequence of SEQ ID NO:74 from amino acid 637 to amino acid 1036. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:74 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:74, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74 having biological activity, the fragment comprising the amino acid sequence  
5 from amino acid 513 to amino acid 522 of SEQ ID NO:74.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- 10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:75.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441; the nucleotide sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441; the nucleotide sequence of the full-length protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:76, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 223 to amino acid 232 of SEQ ID NO:76.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:75.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and
    - (ab) the nucleotide sequence of the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
  - (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75; and

(bb) the nucleotide sequence of the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:75 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:75, but excluding the poly(A) tail at the 3' end of SEQ ID NO:75. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:76;

(b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and

(c) the amino acid sequence encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:76. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
10 of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising the amino acid sequence from amino acid 223 to amino acid 232 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592;

20 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dx146\_12 deposited under accession number ATCC 98818;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;

25 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dx146\_12 deposited under accession number ATCC 98818;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;

30 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

5 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:77.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592; the nucleotide sequence of the full-length protein coding sequence of clone dx146\_12 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone dx146\_12 deposited under accession number ATCC 98818. In other preferred embodiments, the  
15 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
20 preferably thirty) contiguous amino acids of SEQ ID NO:78, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 247 to amino acid 256 of SEQ ID NO:78.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
25 ID NO:77.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and



- (ab) the nucleotide sequence of the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77; and

- (bb) the nucleotide sequence of the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and

- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:77 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:77, but excluding the poly(A) tail at the 3' end of SEQ ID NO:77. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;

- (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:78. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising the amino acid sequence from amino acid 247 to amino acid 256 of SEQ ID NO:78.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:79 from nucleotide 19 to nucleotide 948;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID
- 20 NO:79 from nucleotide 337 to nucleotide 948;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818;
- (e) a polynucleotide encoding the full-length protein encoded by the
- 25 cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA
- 30 insert of clone dx219\_13 deposited under accession number ATCC 98818;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:80;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:79.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:79 from nucleotide 19 to nucleotide 948; the nucleotide sequence of SEQ ID NO:79 from nucleotide 337 to nucleotide 948; the nucleotide sequence of the full-length protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:80, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 150 to amino acid 159 of SEQ ID NO:80.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:79.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and

(ab) the nucleotide sequence of the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79; and

(bb) the nucleotide sequence of the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:79 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:79, but excluding the poly(A) tail at the 3' end of SEQ ID NO:79. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 19 to nucleotide 948, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from nucleotide 19 to nucleotide 948, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide

19 to nucleotide 948. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:79 from nucleotide 337 to nucleotide 948, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:79 from  
5 nucleotide 337 to nucleotide 948, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:79 from nucleotide 337 to nucleotide 948.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:80;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:80. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:80, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80 having biological activity, the fragment comprising the amino acid sequence from amino acid 150 to amino acid 159 of SEQ ID NO:80.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:81 from nucleotide 62 to nucleotide 286;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:82;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:81.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286; the nucleotide sequence of SEQ ID NO:81 from nucleotide 62 to nucleotide 286; the nucleotide sequence of the full-length protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:82, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the

fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:82.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:81.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group  
10 consisting of:

(aa) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and

(ab) the nucleotide sequence of the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25 (ba) SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81; and

(bb) the nucleotide sequence of the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:81 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:81, but excluding the poly(A) tail at the 3' end of SEQ ID NO:81. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:81 from nucleotide 62 to nucleotide 286, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:81 from nucleotide 62 to nucleotide 286, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:81 from nucleotide 62 to nucleotide 286.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:82;
- (b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:82. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:82, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82 having biological activity, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:82.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83;



(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 141 to nucleotide 572;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572;

5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:84;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;

20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

25 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:83.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID  
30 NO:83 from nucleotide 141 to nucleotide 572; the nucleotide sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572; the nucleotide sequence of the full-length protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide

encodes the full-length or a mature protein encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:84, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76 of SEQ ID NO:84.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:83.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:83; and

(ab) the nucleotide sequence of the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:83; and

(bb) the nucleotide sequence of the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:83 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:83. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 141 to nucleotide 572, and extending contiguously from a nucleotide sequence  
10 corresponding to the 5' end of said sequence of SEQ ID NO:83 from nucleotide 141 to nucleotide 572, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide 141 to nucleotide 572. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572, and  
15 extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:84;
- (b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:84. In further preferred embodiments, the present invention provides a protein comprising a fragment of the  
30 amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:84, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76 of SEQ ID NO:84.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:85 from nucleotide 505 to nucleotide 3210;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:85.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210; the nucleotide sequence of SEQ ID NO:85

from nucleotide 505 to nucleotide 3210; the nucleotide sequence of the full-length protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:86, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 465 to amino acid 474 of SEQ ID NO:86.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:85.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and

(ab) the nucleotide sequence of the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85; and

(bb) the nucleotide sequence of the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
10 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:85 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:85, but excluding the poly(A) tail at the 3' end of SEQ ID NO:85. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210. Also preferably the polynucleotide isolated according to the above  
20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:85 from nucleotide 505 to nucleotide 3210, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:85 from nucleotide 505 to nucleotide 3210, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:85 from nucleotide 505 to nucleotide 3210.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:86;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and

(c) the amino acid sequence encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:86. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:86, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising the amino acid sequence from amino acid 465 to amino acid 474 of SEQ ID NO:86.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 522 to nucleotide 899;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:87.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899; the nucleotide sequence of SEQ ID NO:87 from nucleotide 522 to nucleotide 899; the nucleotide sequence of the full-length protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:88.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:87.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
    - (aa) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and
    - (ab) the nucleotide sequence of the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
  - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and



(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

5 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:87, but excluding the poly(A) tail at the 3' end of SEQ ID NO:87; and

10 (bb) the nucleotide sequence of the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

15 (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:87 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:87, but  
20 excluding the poly(A) tail at the 3' end of SEQ ID NO:87. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899, to a nucleotide  
25 sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:87 from nucleotide 522 to nucleotide 899, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:87 from  
30 nucleotide 522 to nucleotide 899, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:87 from nucleotide 522 to nucleotide 899.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
- (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:88. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:88, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 138 to amino acid 147 of SEQ ID NO:88.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 6 to nucleotide 452;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:89 from nucleotide 399 to nucleotide 452;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:90;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:89.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:89 from nucleotide 6 to nucleotide 452; the nucleotide sequence of SEQ ID NO:89 from nucleotide 399 to nucleotide 452; the nucleotide sequence of the full-length protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818; or the nucleotide sequence of a mature protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:90, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 69 to amino acid 78 of SEQ ID NO:90.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:89.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and

(ab) the nucleotide sequence of the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89; and

(bb) the nucleotide sequence of the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:89 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:89, but excluding the poly(A) tail at the 3' end of SEQ ID NO:89. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89 from nucleotide 6 to nucleotide 452, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from nucleotide 6 to nucleotide 452, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:89 from nucleotide

6 to nucleotide 452. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:89 from nucleotide 399 to nucleotide 452, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:89 from  
5 nucleotide 399 to nucleotide 452, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:89 from nucleotide 399 to nucleotide 452.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:90;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:90. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:90, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising the amino acid sequence from amino acid 69 to amino acid 78 of SEQ ID NO:90.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:91 from nucleotide 682 to nucleotide 1179;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:92;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:91.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179; the nucleotide sequence of SEQ ID NO:91 from nucleotide 682 to nucleotide 1179; the nucleotide sequence of the full-length protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the

fragment comprising the amino acid sequence from amino acid 191 to amino acid 200 of SEQ ID NO:92.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:91.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

10

(aa) SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91; and

(ab) the nucleotide sequence of the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;

15

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

25

(ba) SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91; and

(bb) the nucleotide sequence of the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:91 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:91, but excluding the poly(A) tail at the 3' end of SEQ ID NO:91. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:91 from nucleotide 682 to nucleotide 1179, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:91 from nucleotide 682 to nucleotide 1179, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:91 from nucleotide 682 to nucleotide 1179.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;
- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and
- (c) the amino acid sequence encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:92. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:92, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment comprising the amino acid sequence from amino acid 191 to amino acid 200 of SEQ ID NO:92.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93;



(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bf169\_13 deposited under accession number ATCC 98822;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone bf169\_13 deposited under accession number ATCC 98822;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:94;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:93.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077; the nucleotide sequence of the full-length protein coding sequence of clone bf169\_13 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone bf169\_13 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein

comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 332 to amino acid 341 of SEQ ID NO:94.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:93.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

- (ab) the nucleotide sequence of the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (ba) SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93; and

- (bb) the nucleotide sequence of the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:93 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:93, but excluding the poly(A) tail at the 3' end of SEQ ID NO:93. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
- (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:94. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:94, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 332 to amino acid 341 of SEQ ID NO:94.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bl152\_12 deposited under accession number ATCC 98822;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone bl152\_12 deposited under accession number ATCC 98822;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:96;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:95.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735; the nucleotide sequence of the full-length protein coding sequence of clone bl152\_12 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone bl152\_12 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:96, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 97 to amino acid 106 of SEQ ID NO:96.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:95.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

15 (ab) the nucleotide sequence of the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

20 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:95, but excluding the poly(A) tail at the 3' end of SEQ ID NO:95; and

(bb) the nucleotide sequence of the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;

30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:95 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:95, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:95. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:96;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone  
bl152\_12 deposited under accession number ATCC 98822;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:96. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:96, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96 having biological activity, the fragment comprising the amino acid sequence from amino acid 97 to amino acid 106 of SEQ ID NO:96.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone bz578\_1 deposited under accession number ATCC 98822;

5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone bz578\_1 deposited under accession number ATCC 98822;

10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:98;

15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

20 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:97.

25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816; the nucleotide sequence of the full-length protein coding sequence of clone bz578\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone bz578\_1 deposited under accession number ATCC 98822. In other preferred embodiments, the  
30 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most

preferably thirty) contiguous amino acids of SEQ ID NO:98, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:98.

5 Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:97.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

10 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and

15 (ab) the nucleotide sequence of the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

20 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

25 (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:97, but excluding the poly(A) tail at the 3' end of SEQ ID NO:97; and

(bb) the nucleotide sequence of the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;

30 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).



Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:97 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:97, but  
5 excluding the poly(A) tail at the 3' end of SEQ ID NO:97. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816, to a nucleotide  
10 sequence corresponding to the 3' end of said sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:98;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:98. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
25 of SEQ ID NO:98, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98 having biological activity, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:98.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide 992;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99 from nucleotide 765 to nucleotide 992;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:100;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:99.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide 992; the nucleotide sequence of SEQ ID NO:99 from nucleotide 765 to nucleotide 992; the nucleotide sequence of the full-length protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822. In further preferred

embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a polynucleotide  
5 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:100.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:99.

10 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
  - (i) preparing one or more polynucleotide probes that hybridize  
15 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

- (aa) SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99; and

- (ab) the nucleotide sequence of the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;

20 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

- (iii) isolating the DNA polynucleotides detected with the probe(s);

and

25 (b) a process comprising the steps of:

- (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

30 (ba) SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99; and

- (bb) the nucleotide sequence of the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;

- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99, and  
5 extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:99 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:99, but excluding the poly(A) tail at the 3' end of SEQ ID NO:99. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide  
10 992, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide 992, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide 992. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID  
15 NO:99 from nucleotide 765 to nucleotide 992, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:99 from nucleotide 765 to nucleotide 992, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:99 from nucleotide 765 to nucleotide 992.

In other embodiments, the present invention provides a composition comprising  
20 a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
- (b) a fragment of the amino acid sequence of SEQ ID NO:100, the  
fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
- 25 (c) the amino acid sequence encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:100. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:100, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:100.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ch245\_1 deposited under accession number ATCC 98822;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ch245\_1 deposited under accession number ATCC 98822;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;
- 25 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:101.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480; the nucleotide sequence of the full-length protein coding sequence of clone ch245\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone ch245\_1

deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822. In further preferred  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:102, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino  
10 acid 45 to amino acid 54 of SEQ ID NO:102.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:101.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- 15 (a) a process comprising the steps of:
- (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- 20 (aa) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and
- (ab) the nucleotide sequence of the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 25 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that  
30 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101; and

- (bb) the nucleotide sequence of the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:101 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:101, but excluding the poly(A) tail at the 3' end of SEQ ID NO:101. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
- (b) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:102. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:102, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 45 to amino acid 54 of SEQ ID NO:102.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103;

5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cj378\_3 deposited under accession number ATCC 98822;

10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cj378\_3 deposited under accession number ATCC 98822;

15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:104;

20 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

25 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

(k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:103.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541; the nucleotide sequence of the full-length protein coding sequence of clone cj378\_3 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone cj378\_3 deposited



under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:104.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:103.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

- (a) a process comprising the steps of:
    - (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (aa) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and
      - (ab) the nucleotide sequence of the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;
    - (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
    - (iii) isolating the DNA polynucleotides detected with the probe(s);
- and
- (b) a process comprising the steps of:
    - (i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
      - (ba) SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103; and

- (bb) the nucleotide sequence of the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- 5 (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ  
10 ID NO:103 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:103, but excluding the poly(A) tail at the 3' end of SEQ ID NO:103. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541, and extending contiguously from a nucleotide sequence corresponding to the 5' end  
15 of said sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group  
20 consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
- (c) the amino acid sequence encoded by the cDNA insert of clone  
25 cj378\_3 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:104. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment preferably  
30 comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:104, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:104.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:105 from nucleotide 401 to nucleotide 2349;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone cw1481\_1 deposited under accession number ATCC 98822;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone cw1481\_1 deposited under accession number ATCC 98822;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- 20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- 25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 30 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:105.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202; the nucleotide sequence of SEQ ID

NO:105 from nucleotide 401 to nucleotide 2349; the nucleotide sequence of the full-length protein coding sequence of clone cw1481\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone cw1481\_1 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 264 to amino acid 273 of SEQ ID NO:106.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:105.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and

(ab) the nucleotide sequence of the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105; and

(bb) the nucleotide sequence of the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;

5 (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a  
10 nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:105 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:105, but excluding the poly(A) tail at the 3' end of SEQ ID NO:105. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence  
15 corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202. Also preferably the polynucleotide isolated according to the above  
20 process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:105 from nucleotide 401 to nucleotide 2349, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:105 from nucleotide 401 to nucleotide 2349, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:105 from nucleotide 401 to nucleotide 2349.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:106;

30 (b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and

(c) the amino acid sequence encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:106. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:106, or a protein comprising a fragment of the amino acid sequence of SEQ  
5 ID NO:106 having biological activity, the fragment comprising the amino acid sequence from amino acid 264 to amino acid 273 of SEQ ID NO:106.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905;
- 15 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:108;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:107.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905; the nucleotide sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905; the nucleotide sequence of the full-length protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:108, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 474 to amino acid 483 of SEQ ID NO:108.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:107.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and

(ab) the nucleotide sequence of the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107; and

(bb) the nucleotide sequence of the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:107 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:107, but excluding the poly(A) tail at the 3' end of SEQ ID NO:107. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:



- (a) the amino acid sequence of SEQ ID NO:108;
- (b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:108. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:108, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 474 to amino acid 483 of SEQ ID NO:108.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 16 to nucleotide 369;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 103 to nucleotide 369;

- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822;

- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;

- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822;

- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;

- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:110;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:109.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:109 from nucleotide 16 to nucleotide 369; the nucleotide sequence of SEQ ID NO:109 from nucleotide 103 to nucleotide 369; the nucleotide sequence of the full-length protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:110, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:110.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:109.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and

(ab) the nucleotide sequence of the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109; and

(bb) the nucleotide sequence of the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:109 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:109, but excluding the poly(A) tail at the 3' end of SEQ ID NO:109. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109 from nucleotide 16 to nucleotide 369, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:109 from nucleotide 16 to nucleotide 369, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:109 from nucleotide

16 to nucleotide 369. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:109 from nucleotide 103 to nucleotide 369, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:109 from  
5 nucleotide 103 to nucleotide 369, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:109 from nucleotide 103 to nucleotide 369.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:110;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:110. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
20 of SEQ ID NO:110, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising the amino acid sequence from amino acid 54 to amino acid 63 of SEQ ID NO:110.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID  
30 NO:111 from nucleotide 2255 to nucleotide 2539;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:112;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:111.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539; the nucleotide sequence of SEQ ID NO:111 from nucleotide 2255 to nucleotide 2539; the nucleotide sequence of the full-length protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112

having biological activity, the fragment comprising the amino acid sequence from amino acid 53 to amino acid 62 of SEQ ID NO:112.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:111.

5 Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize  
10 in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:111, but excluding the poly(A) tail at the  
3' end of SEQ ID NO:111; and

(ab) the nucleotide sequence of the cDNA insert of clone  
km225\_1 deposited under accession number ATCC 98822;

15 (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

20 (b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:111, but excluding the poly(A) tail at the  
25 3' end of SEQ ID NO:111; and

(bb) the nucleotide sequence of the cDNA insert of clone  
km225\_1 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

30 (iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ

ID NO:111 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:111, but excluding the poly(A) tail at the 3' end of SEQ ID NO:111. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:111 from nucleotide 2255 to nucleotide 2539, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:111 from nucleotide 2255 to nucleotide 2539, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:111 from nucleotide 2255 to nucleotide 2539.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
- (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
- (c) the amino acid sequence encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:112. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:112, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112 having biological activity, the fragment comprising the amino acid sequence from amino acid 53 to amino acid 62 of SEQ ID NO:112.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 1734 to nucleotide 2030;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030;
- 5 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;
- 10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;
- 15 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:114;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;
- 20 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and
- 25 (m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:113.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:113 from nucleotide 1734 to nucleotide 2030; the nucleotide sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030; the nucleotide sequence of the full-length protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822. In other preferred

30



embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:114, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 44 to amino acid 53 of SEQ ID NO:114.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:113.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and

(ab) the nucleotide sequence of the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113; and

(bb) the nucleotide sequence of the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

5 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:113 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:113, but excluding the poly(A) tail at the 3' end of SEQ ID NO:113. Also preferably the  
10 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113 from nucleotide 1734 to nucleotide 2030, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 1734 to nucleotide 2030, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:113  
15 from nucleotide 1734 to nucleotide 2030. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030, to a nucleotide  
20 sequence corresponding to the 3' end of said sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 25 (a) the amino acid sequence of SEQ ID NO:114;  
(b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and  
(c) the amino acid sequence encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;  
30 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:114. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids

of SEQ ID NO:114, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114 having biological activity, the fragment comprising the amino acid sequence from amino acid 44 to amino acid 53 of SEQ ID NO:114.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350;
- 10 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822;
- 15 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above;
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above ;
- 30 (l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i); and

(m) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(i) and that has a length that is at least 25% of the length of SEQ ID NO:115.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350; the nucleotide sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350; the nucleotide sequence of the full-length protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids of SEQ ID NO:116, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino acid 87 to amino acid 96 of SEQ ID NO:116.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ ID NO:115.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

(i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115; and

(ab) the nucleotide sequence of the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;

(ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and

(iii) isolating the DNA polynucleotides detected with the probe(s);

and

(b) a process comprising the steps of:

(i) preparing one or more polynucleotide primers that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(ba) SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115; and

(bb) the nucleotide sequence of the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;

(ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;

(iii) amplifying human DNA sequences; and

(iv) isolating the polynucleotide products of step (b)(iii).

Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:115 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:115, but excluding the poly(A) tail at the 3' end of SEQ ID NO:115. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350. Also preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:116;

- (b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and
- (c) the amino acid sequence encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:116. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids
- 10 of SEQ ID NO:116, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising the amino acid sequence from amino acid 87 to amino acid 96 of SEQ ID NO:116.

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:117 from nucleotide 837 to nucleotide 1094;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone my340\_1 deposited under accession
- 20 number ATCC 98822;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;
- (e) a polynucleotide comprising the nucleotide sequence of a mature
- 25 protein coding sequence of clone my340\_1 deposited under accession number ATCC 98822;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;
- (g) a polynucleotide encoding a protein comprising the amino acid
- 30 sequence of SEQ ID NO:118;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above;

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above ;

5 (k) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h); and

(l) a polynucleotide that hybridizes under stringent conditions to any one of the polynucleotides specified in (a)-(h) and that has a length that is at least 25% of the length of SEQ ID NO:117.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:117 from nucleotide 837 to nucleotide 1094; the nucleotide sequence of the full-length protein coding sequence of clone my340\_1 deposited under accession number ATCC 98822; or the nucleotide sequence of a mature protein coding sequence of clone my340\_1 deposited under accession number ATCC 98822. In other preferred embodiments, the  
15 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
20 preferably thirty) contiguous amino acids of SEQ ID NO:118, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:118.

Other embodiments provide the gene corresponding to the cDNA sequence of SEQ  
25 ID NO:117.

Further embodiments of the invention provide isolated polynucleotides produced according to a process selected from the group consisting of:

(a) a process comprising the steps of:

30 (i) preparing one or more polynucleotide probes that hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:

(aa) SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117; and

- (ab) the nucleotide sequence of the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;
- (ii) hybridizing said probe(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C; and
- 5 (iii) isolating the DNA polynucleotides detected with the probe(s);

and

- (b) a process comprising the steps of:
- (i) preparing one or more polynucleotide primers that
- 10 hybridize in 6X SSC at 65 degrees C to a nucleotide sequence selected from the group consisting of:
- (ba) SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117; and
- (bb) the nucleotide sequence of the cDNA insert of clone
- 15 my340\_1 deposited under accession number ATCC 98822;
- (ii) hybridizing said primer(s) to human genomic DNA in conditions at least as stringent as 4X SSC at 50 degrees C;
- (iii) amplifying human DNA sequences; and
- (iv) isolating the polynucleotide products of step (b)(iii).

- 20 Preferably the polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117, and extending contiguously from a nucleotide sequence corresponding to the 5' end of SEQ ID NO:117 to a nucleotide sequence corresponding to the 3' end of SEQ ID NO:117, but excluding the poly(A) tail at the 3' end of SEQ ID NO:117. Also preferably the
- 25 polynucleotide isolated according to the above process comprises a nucleotide sequence corresponding to the cDNA sequence of SEQ ID NO:117 from nucleotide 837 to nucleotide 1094, and extending contiguously from a nucleotide sequence corresponding to the 5' end of said sequence of SEQ ID NO:117 from nucleotide 837 to nucleotide 1094, to a nucleotide sequence corresponding to the 3' end of said sequence of SEQ ID NO:117 from nucleotide
- 30 837 to nucleotide 1094.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:118;



(b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and

(c) the amino acid sequence encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;

5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:118. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) contiguous amino acids  
10 of SEQ ID NO:118, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118 having biological activity, the fragment comprising the amino acid sequence from amino acid 38 to amino acid 47 of SEQ ID NO:118.

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial,  
15 yeast, insect and mammalian cells, transformed with such polynucleotide compositions. Also provided by the present invention are organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein.

Processes are also provided for producing a protein, which comprise:

20 (a) growing a culture of the host cell transformed with such polynucleotide compositions in a suitable culture medium; and

(b) purifying the protein from the culture.

The protein produced according to such methods is also provided by the present invention.

25 Protein compositions of the present invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically  
30 effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are schematic representations of the pED6 and pNOTs vectors, respectively, used for deposit of clones disclosed herein.

### DETAILED DESCRIPTION

#### 5 ISOLATED PROTEINS AND POLYNUCLEOTIDES

Nucleotide and amino acid sequences, as presently determined, are reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone in accordance with known methods. The predicted amino acid sequence (both full-length  
10 and mature forms) can then be determined from such nucleotide sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have determined to be the reading frame best identifiable with sequence information available  
15 at the time of filing.

As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell  
20 in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

#### Clone "bn365\_53"

A polynucleotide of the present invention has been identified as clone "bn365\_53".  
25 bn365\_53 was isolated from a human adult placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bn365\_53 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein  
30 as "bn365\_53 protein").

The nucleotide sequence of bn365\_53 as presently determined is reported in SEQ ID NO:1, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bn365\_53 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bn365\_53 should be approximately 650 bp.

The nucleotide sequence disclosed herein for bn365\_53 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bn365\_53 demonstrated at least some similarity with sequences identified as AA242967 (zr65g11.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 668324 5') and N40141 (yw73c12.r1 Homo sapiens cDNA clone 257878 5'). The predicted amino acid sequence disclosed herein for bn365\_53 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bn365\_53 protein demonstrated at least some similarity to sequences identified as D63484 (KIAA0150 protein [Homo sapiens]) and to the GAGE-1 to GAGE-6 family of human proteins expressed in tumors (GenBank Accession Numbers U19142-U19147). The amino acid sequence of SEQ ID NO:2 contains two RGD (Arg-Gly-Asp) motifs (around residues 12 and 75): the sequence Arg-Gly-Asp, found in fibronectin, is crucial for its interaction with its cell surface receptor, an integrin. What has been called the 'RGD' tripeptide is also found in the sequences of a number of other proteins, where it has been shown to play a role in cell adhesion. These proteins are: some forms of collagens, fibrinogen, vitronectin, von Willebrand factor (VWF), snake disintegrins, and slime mold discoidins. Based upon sequence similarity, bn365\_53 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of bn365\_53 indicates that it may contain one or more repetitive elements.

#### Clone "bo342\_2"

A polynucleotide of the present invention has been identified as clone "bo342\_2". bo342\_2 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bo342\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "bo342\_2 protein").

The nucleotide sequence of bo342\_2 as presently determined is reported in SEQ ID NO:3, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bo342\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:4. Amino

acids 372 to 384 of SEQ ID NO:4 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 385. Amino acids 1 to 13 are also a possible leader/signal sequence, with the predicted mature amino acid sequence beginning in that case at amino acid 14. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the bo342\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bo342\_2 should be approximately 2600 bp.

The nucleotide sequence disclosed herein for bo342\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bo342\_2 demonstrated at least some similarity with sequences identified as AA306000 (EST177027 Jurkat T-cells VI Homo sapiens cDNA 5' end) and W94256 (ze12b02.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 358731 3' similar to contains Alu repetitive element). Based upon sequence similarity, bo342\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts six potential transmembrane domains within the bo342\_2 protein sequence, centered around amino acids 300, 320, 380, 410, 430, and 490 of SEQ ID NO:4, respectively. The nucleotide sequence of bo342\_2 indicates that it may contain Alu or other repetitive elements.

#### Clone "dn721\_8"

A polynucleotide of the present invention has been identified as clone "dn721\_8". dn721\_8 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dn721\_8 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dn721\_8 protein").

The nucleotide sequence of dn721\_8 as presently determined is reported in SEQ ID NO:5, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dn721\_8 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:6.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dn721\_8 should be approximately 2900 bp.

The nucleotide sequence disclosed herein for dn721\_8 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dn721\_8 demonstrated at least some similarity with sequences identified as H63637 (yr34b12.r1 Homo sapiens cDNA clone 207167 5'), N31598 (yy20b12.s1 Homo sapiens cDNA clone 271775 3'), and R61419 (yh15e05.r1 Homo sapiens cDNA clone 37671 5'). Based upon sequence similarity, dn721\_8 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two possible transmembrane domains within the dn721\_8 protein sequence, one centered around amino acid 269 and another around amino acid 457 of SEQ ID NO:6.

#### Clone "dn834\_1"

A polynucleotide of the present invention has been identified as clone "dn834\_1". dn834\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dn834\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dn834\_1 protein").

The nucleotide sequence of dn834\_1 as presently determined is reported in SEQ ID NO:7, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dn834\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:8.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dn834\_1 should be approximately 900 bp.

The nucleotide sequence disclosed herein for dn834\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dn834\_1 demonstrated at least some similarity with sequences identified as AA544005 (vj83h07.r1 Soares mouse mammary gland NbMMG Mu's musculus cDNA clone 935677 5'), AL022163 (Human DNA sequence \*\*\* SEQUENCING IN PROGRESS \*\*\* from clone 551E13; HTGS phase 1), L44560 (Homo sapiens thymus mRNA (randomly primed, normalized), single-pass sequence), and T72271 (Human B cell surface antigen cDNA). The predicted amino acid sequence disclosed herein for dn834\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the

BLASTX search protocol. The predicted dn834\_1 protein demonstrated at least some similarity to sequences identified as R47496 (Translated sequence of domains I and II of celD cDNA in clone pCNP4). Based upon sequence similarity, dn834\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer  
5 program predicts three potential transmembrane domains within the dn834\_1 protein sequence, centered around amino acids 59, 84, and 145 of SEQ ID NO:8, respectively.

dn834\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 18 kDa was detected in membrane fractions using SDS  
polyacrylamide gel electrophoresis.

10

#### Clone "pd278\_5"

A polynucleotide of the present invention has been identified as clone "pd278\_5". A cDNA clone was first isolated from a human fetal kidney cDNA library using methods  
15 which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate pd278\_5 from a human adult kidney cDNA library. pd278\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to  
20 herein as "pd278\_5 protein").

The nucleotide sequence of pd278\_5 as presently determined is reported in SEQ ID NO:9, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pd278\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:10. Amino  
25 acids 61 to 73 of SEQ ID NO:10 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 74. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pd278\_5 protein.

30 There are two additional and mutually overlapping possible open reading frames close to the 5' end of SEQ ID NO:9 (bases 82 - 420 and bases 119 - 414). The translated open reading frame of bases 119 - 414 has a predicted leader/signal sequence from amino acid 49 to amino acid 61, with the predicted mature amino acid sequence beginning at

amino acid 62. Each of the additional possible open reading frames has a predicted transmembrane domain.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pd278\_5 should be approximately 2000 bp.

5 The nucleotide sequence disclosed herein for pd278\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pd278\_5 demonstrated at least some similarity with sequences identified as AA292241 (zt50d11.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 725781 5'), AA428245 zw51d10.s1 Soares total fetus Nb2HF8 9w Homo sapiens  
10 cDNA clone 773587 3'), AA599487 (ag23f05.s1 Jia bone marrow stroma Homo sapiens cDNA clone 1071201 3'), AA827135 (ob53b03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE 1335053 3'), H54322 (yq90d03.s1 Homo sapiens cDNA clone 203045 3'), and T22170 (Human gene signature HUMGS03741). The predicted amino acid sequence disclosed herein for pd278\_5 was searched against the GenPept and GeneSeq amino acid  
15 sequence databases using the BLASTX search protocol. The predicted pd278\_5 protein demonstrated at least some similarity to sequences identified as R13144 (Deleted in Colorectal Carcinomas) and X13885 (extensin (AA 1-620) [Nicotiana tabacum]). Based upon sequence similarity, pd278\_5 proteins and each similar protein or peptide may share at least some activity.

20

#### Clone "pe80\_1"

A polynucleotide of the present invention has been identified as clone "pe80\_1". pe80\_1 was isolated from a human adult blood (chronic myelogenous leukemia K562) cDNA library using methods which are selective for cDNAs encoding secreted proteins  
25 (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pe80\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pe80\_1 protein").

The nucleotide sequence of pe80\_1 as presently determined is reported in SEQ ID  
30 NO:11, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pe80\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:12.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pe80\_1 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for pe80\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pe80\_1 demonstrated at least some similarity with sequences identified as AA291078 (zs47b04.r1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone  
5 IMAGE:700591 5'), AA429912 (zw66e06.s1 Soares testis NHT Homo sapiens cDNA clone 781186 3'), H82367 (yv79d06.r1 Homo sapiens cDNA clone 248939 5' similar to contains Alu repetitive element;contains OFR repetitive element), Q60627 (Human brain Expressed Sequence Tag EST02640), and R20261 (yg20a02.r1 Homo sapiens cDNA clone 32587 5').  
10 Based upon sequence similarity, pe80\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two possible transmembrane domains within the pe80\_1 protein sequence, one centered around amino acid 58 and another around amino acid 109 of SEQ ID NO:12. The nucleotide sequence of pe80\_1 indicates that it may contain an Alu repetitive element.

15 Clone "pm113\_1"

A polynucleotide of the present invention has been identified as clone "pm113\_1". pm113\_1 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis  
20 of computer analysis of the amino acid sequence of the encoded protein. pm113\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm113\_1 protein").

The nucleotide sequence of pm113\_1 as presently determined is reported in SEQ ID NO:13, and includes a poly(A) tail. What applicants presently believe to be the proper  
25 reading frame and the predicted amino acid sequence of the pm113\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:14. Amino acids 41 to 53 of SEQ ID NO:14 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 54. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
30 should the predicted leader/signal sequence not be separated from the remainder of the pm113\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm113\_1 should be approximately 1700 bp.



The nucleotide sequence disclosed herein for pm113\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pm113\_1 demonstrated at least some similarity with sequences identified as AA009482 (zi04c03.r1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 429796 5'), AA350890 (EST58401 Infant brain Homo sapiens cDNA 3' end), AC003030 (Human DNA from chromosome 19-specific cosmid R29828, genomic sequence, complete sequence), H98961 (yx11b02.s1 Homo sapiens cDNA clone 261387 3'), R07796 (yf15e05.r1 Homo sapiens cDNA clone), T22151 (Human gene signature HUMGS03721), and W68491 (zd34h02.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 342579 5'). Based upon sequence similarity, pm113\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "pm749\_8"

A polynucleotide of the present invention has been identified as clone "pm749\_8". pm749\_8 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pm749\_8 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm749\_8 protein").

The nucleotide sequence of pm749\_8 as presently determined is reported in SEQ ID NO:15, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pm749\_8 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:16.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm749\_8 should be approximately 2300 bp.

The nucleotide sequence disclosed herein for pm749\_8 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pm749\_8 demonstrated at least some similarity with sequences identified as AA314025 (EST185879 Colon carcinoma (HCC) cell line II Homo sapiens cDNA 5' end) and AA374458 (EST86612 HSC172 cells I Homo sapiens cDNA 5' end). The predicted amino acid sequence disclosed herein for pm749\_8 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol.

The predicted pm749\_8 protein demonstrated at least some similarity to sequences identified as D89169 (similar to *Saccharomyces cerevisiae* SCD6 protein, SWISS-PROT Accession Number P45978 [*Schizosaccharomyces pombe*]) and U30384 (Scd6p [*Saccharomyces cerevisiae*]). Based upon sequence similarity, pm749\_8 proteins and each  
5 similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pm749\_8 protein sequence centered around amino acid 138 of SEQ ID NO:16.

Clone "pt31\_4"

10 A polynucleotide of the present invention has been identified as clone "pt31\_4". pt31\_4 was isolated from a human adult blood (lymphoblastic leukemia MOLT-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pt31\_4  
15 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pt31\_4 protein").

The nucleotide sequence of pt31\_4 as presently determined is reported in SEQ ID NO:17, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pt31\_4 protein corresponding  
20 to the foregoing nucleotide sequence is reported in SEQ ID NO:18. Amino acids 19 to 31 of SEQ ID NO:18 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pt31\_4  
25 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pt31\_4 should be approximately 3200 bp.

The nucleotide sequence disclosed herein for pt31\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
30 FASTA search protocols. pt31\_4 demonstrated at least some similarity with sequences identified as AA348130 (EST54532 Fetal heart II *Homo sapiens* cDNA 5' end), AA350691 (EST58082 Infant brain *Homo sapiens* cDNA 5' end), AC001226 (Genomic sequence from Human 13, complete sequence), H22773 (ym54c06.r1 *Homo sapiens* cDNA clone 52351 5'), and R21869 (yh22b10.s1 *Homo sapiens* cDNA clone 130459 3'). The predicted amino

acid sequence disclosed herein for pt31\_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pt31\_4 protein demonstrated at least some similarity to sequences identified as U53147 (C01B7.6 [Caenorhabditis elegans]). Based upon sequence similarity, pt31\_4 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five potential transmembrane domains within the pt31\_4 protein sequence, centered around amino acids 90, 110, 210, 410, and 590 of SEQ ID NO:18, respectively.

10        Clone "pv296\_5"

A polynucleotide of the present invention has been identified as clone "pv296\_5". pv296\_5 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pv296\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pv296\_5 protein").

The nucleotide sequence of pv296\_5 as presently determined is reported in SEQ ID NO:19, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pv296\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:20.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pv296\_5 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for pv296\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pv296\_5 demonstrated at least some similarity with sequences identified as AA022471 (ze70c01.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 364320 3'), AA335246 (EST39647 Epididymus Homo sapiens cDNA 5' end), and AA481308 (zv06a05.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 752816 5'). Based upon sequence similarity, pv296\_5 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pv296\_5 protein sequence centered around amino acid 32 of SEQ ID NO:20.

Clone "er311\_20"

A polynucleotide of the present invention has been identified as clone "er311\_20". er311\_20 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
5 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. er311\_20 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "er311\_20 protein").

The nucleotide sequence of er311\_20 as presently determined is reported in SEQ  
10 ID NO:21, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the er311\_20 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:22. Amino acids 654 to 666 of SEQ ID NO:22 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 667. Due to the  
15 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the er311\_20 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone er311\_20 should be approximately 2800 bp.

20 The nucleotide sequence disclosed herein for er311\_20 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. er311\_20 demonstrated at least some similarity with sequences identified as AF035526 (Mus musculus kanadapin mRNA, complete cds), R18277 (yg01c06.r1 Homo sapiens cDNA clone 31018 5' similar to SP:ZK632.2 CE00419  
25 COILED COIL PROTEIN), R47371 (Hf060-r Homo sapiens cDNA clone f060-r), and Z40133 (H. sapiens partial cDNA sequence; clone c-1sh08). The predicted amino acid sequence disclosed herein for er311\_20 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted er311\_20 protein demonstrated at least some similarity to sequences identified as  
30 AF035526 (kanadapin [Mus musculus]) and Z22181 (ZK632.2 [Caenorhabditis elegans]). The mouse kanadapin protein and the predicted er311\_20 protein both contain poly-glutamic acid stretches within their C-terminal portions. Based upon sequence similarity, er311\_20 proteins and each similar protein or peptide may share at least some activity.

The TopPredII computer program predicts two potential transmembrane domains within the er311\_20 protein sequence, one centered around amino acid 667 and another at the extreme C-terminus of SEQ ID NO:22.

er311\_20 protein was expressed in a COS cell expression system, and an expressed  
5 protein band of approximately 91 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "fh149\_12"

A polynucleotide of the present invention has been identified as clone "fh149\_12".  
10 fh149\_12 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. fh149\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein  
15 as "fh149\_12 protein").

The nucleotide sequence of fh149\_12 as presently determined is reported in SEQ ID NO:23, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the fh149\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:24. Amino  
20 acids 133 to 145 of SEQ ID NO:24 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 146. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the fh149\_12 protein.

25 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone fh149\_12 should be approximately 2500 bp.

The nucleotide sequence disclosed herein for fh149\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fh149\_12 demonstrated at least some similarity with sequences  
30 identified as AA653557 (ag67b07.s1 Gessler Wilms tumor Homo sapiens cDNA clone 1127989 3'), AA191185 (zq45b09.r1 Stratagene hNT neuron (#937233) Homo sapiens cDNA clone 632633 5'), H20588 (yn63d06.r1 Homo sapiens cDNA clone 173099 5'), R16294 (yf93b09.r1 Homo sapiens cDNA clone 30087 5'), T08702 (Rat OCT-1 gene),

T25120 (Human gene signature HUMGS07278), U38652 (Mus musculus transmembrane transporter (Lx1) mRNA, complete cds), U77086 (Human organic cation transporter 1 (hOCT1) mRNA, complete cds), and Z66539 (H.sapiens creatine transporter gene). The predicted amino acid sequence disclosed herein for fh149\_12 was searched against the  
5 GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted fh149\_12 protein demonstrated at least some similarity to sequences identified as D17546 (Collagen [Mus musculus]), R77676 (Rat OCT-1 protein), and U77086 (organic cation transporter 1 [Homo sapiens]). The fh149\_12 protein also shows  
10 some homology to organic cation transporters from rat (GenBank L27651) and pig (GenBank Y09400) cells. Based upon sequence similarity, fh149\_12 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts eleven potential transmembrane domains within the fh149\_12 protein sequence, centered around amino acids 40, 112, 139, 162, 200, 229, 349, 376, 405, 436, and 467 of SEQ ID NO:24, respectively.

15

#### Clone "pc201\_6"

A polynucleotide of the present invention has been identified as clone "pc201\_6". pc201\_6 was isolated from a human adult retina (retinoblastoma WERI-Rb1) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat.  
20 No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pc201\_6 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pc201\_6 protein").

The nucleotide sequence of pc201\_6 as presently determined is reported in SEQ  
25 ID NO:25, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pc201\_6 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:26. Amino acids 20 to 32 of SEQ ID NO:26 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 33. Due to the hydrophobic nature  
30 of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pc201\_6 protein.

A partial cDNA clone related to pc201\_6, pc201\_SP, was also isolated from a human adult retina (retinoblastoma WERI-Rb1) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. The pc201\_SP clone appears to encode a splice variant of the pc201\_6 protein. The amino acid sequence of the predicted pc201\_SP splice variant protein comprises the amino acid sequence reported in SEQ ID NO:177.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pc201\_6 should be approximately 2500 bp.

The nucleotide sequence disclosed herein for pc201\_6 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pc201\_6 demonstrated at least some similarity with sequences identified as AA256414 (zr80d11.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 682005 5' similar to WP EEED8.9 CE01893), AA342139 (EST47690 Fetal spleen Homo sapiens cDNA 3' end), AC004085 (Homo sapiens; HTGS phase 1, 72 unordered pieces), AF035950 (Homo sapiens putative DDB p127-associated protein mRNA, partial cds), and H10436 (ym08d09.s1 Homo sapiens cDNA clone 47394 3'). The predicted amino acid sequence disclosed herein for pc201\_6 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pc201\_6 protein demonstrated at least some similarity to sequences identified as AF035950 (putative DDB p127-associated protein [Homo sapiens]) and U23484 (EEED8.5 [Caenorhabditis elegans]). Based upon sequence similarity, pc201\_6 proteins and each similar protein or peptide may share at least some activity.

#### Clone "pl87\_1"

A polynucleotide of the present invention has been identified as clone "pl87\_1". pl87\_1 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pl87\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pl87\_1 protein").

The nucleotide sequence of pl87\_1 as presently determined is reported in SEQ ID NO:27, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pl87\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28.

5 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pl87\_1 should be approximately 700 bp.

The nucleotide sequence disclosed herein for pl87\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pl87\_1 demonstrated at least some similarity with sequences  
10 identified as AA371861 (EST83927 Parathyroid gland tumor I Homo sapiens cDNA 5' end) and AA861863 (ak39e11.s1 Soares testis NHT Homo sapiens cDNA clone IMAGE:1408364 3'). Based upon sequence similarity, pl87\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domains within the pl87\_1 protein sequence centered around amino acid  
15 50 of SEQ ID NO:28.

pl87\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 22 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### 20 Clone "pm514\_4"

A polynucleotide of the present invention has been identified as clone "pm514\_4". pm514\_4 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis  
25 of computer analysis of the amino acid sequence of the encoded protein. pm514\_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm514\_4 protein").

The nucleotide sequence of pm514\_4 as presently determined is reported in SEQ ID NO:29, and includes a poly(A) tail. What applicants presently believe to be the proper  
30 reading frame and the predicted amino acid sequence of the pm514\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:30.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm514\_4 should be approximately 3000 bp.



The nucleotide sequence disclosed herein for pm514\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pm514\_4 demonstrated at least some similarity with sequences identified as AA393855 (zv64g11.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 758468 5' similar to WP ZK1248.14 CE02898), AA427943 (zw53d10.s1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773779 3'), AA434561 (zw53d10.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 773779 5'), W49736 (zc41a03.r1 Soares senescent fibroblasts NbHSF Homo sapiens cDNA clone 324844 5'), and U95822 (Human putative transmembrane GTPase mRNA, partial cds). The predicted amino acid sequence disclosed herein for pm514\_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted pm514\_4 protein demonstrated at least some similarity to sequences identified as U95822 (putative transmembrane GTPase [Homo sapiens]). Based upon sequence similarity, pm514\_4 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pm514\_4 protein sequence, centered around amino acid 600 of SEQ ID NO:30.

#### Clone "co155\_12"

A polynucleotide of the present invention has been identified as clone "co155\_12". co155\_12 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. co155\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "co155\_12 protein").

The nucleotide sequence of co155\_12 as presently determined is reported in SEQ ID NO:31, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the co155\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:32. Amino acids 21 to 33 of SEQ ID NO:32 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 34. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain

should the predicted leader/signal sequence not be separated from the remainder of the co155\_12 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone co155\_12 should be approximately 2700 bp.

5       The nucleotide sequence disclosed herein for co155\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. co155\_12 demonstrated at least some similarity with sequences identified as AA578373 (nl23d11.s1 NCI\_CGAP\_HSC1 Homo sapiens cDNA clone IMAGE:1041525, mRNA sequence), N43800 (yy42h09.r1 Homo sapiens cDNA clone 10 273953 5'), and W40418 (zc82c10.r1 Pancreatic Islet Homo sapiens cDNA clone 328818 5', mRNA sequence). The predicted amino acid sequence disclosed herein for co155\_12 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted co155\_12 protein demonstrated at least some similarity to the sequences identified as L12721 (transmembrane domain encoded by 15 1099-1167) and AF004849 (human serine/threonine protein kinase). Based upon sequence similarity, co155\_12 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five additional potential transmembrane domains within the co155\_12 protein sequence, centered around amino acids 90, 180, 470, 580, and 610 of SEQ ID NO:32, respectively.

20

#### Clone "fn189\_13"

A polynucleotide of the present invention has been identified as clone "fn189\_13". fn189\_13 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was 25 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. fn189\_13 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "fn189\_13 protein").

30       The nucleotide sequence of fn189\_13 as presently determined is reported in SEQ ID NO:33, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the fn189\_13 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:34. Amino acids 9 to 21 of SEQ ID NO:34 are a predicted leader/signal sequence, with the predicted

mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the fn189\_13 protein.

- 5           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone fn189\_13 should be approximately 3800 bp.

The nucleotide sequence disclosed herein for fn189\_13 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fn189\_13 demonstrated at least some similarity with sequences  
10 identified as AA144270 (mr14d12.r1 Soares mouse 3NbMS Mus musculus cDNA clone 597431 5') and N27605 (yx44e10.r1 Homo sapiens cDNA clone 264618 5'). The predicted amino acid sequence disclosed herein for fn189\_13 was searched against the GenPept, GeneSeq, and SWISS\_PROT amino acid sequence databases using the BLASTX search protocol. The predicted fn189\_13 protein demonstrated at least some similarity to  
15 sequences identified as P32857 (PROTEIN PTM1 PRECURSOR [Saccharomyces cerevisiae]) and U64598 (weakly similar to S. cerevisiae PTM1 precursor (SP:P32857) [Caenorhabditis elegans]). Based upon sequence similarity, fn189\_13 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts five additional potential transmembrane domains within the fn189\_13  
20 protein sequence, centered around amino acids 225, 260, 340, 360, and 420 of SEQ ID NO:34, respectively.

#### Clone "lv2\_47"

A polynucleotide of the present invention has been identified as clone "lv2\_47".  
25 lv2\_47 was isolated from a human adult thyroid cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. lv2\_47 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as  
30 "lv2\_47 protein").

The nucleotide sequence of lv2\_47 as presently determined is reported in SEQ ID NO:35, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the lv2\_47 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:36. The TopPredII

computer program predicts a potential transmembrane domain within the lv2\_47 protein sequence of SEQ ID NO:36, centered around amino acid 60.

Another potential lv2\_47 reading frame and predicted amino acid sequence is encoded by basepairs 365 to 880 of SEQ ID NO:35 and is reported in SEQ ID NO:178.

5 Amino acids 49 to 61 of SEQ ID NO:178 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 62. Due to the hydrophobic nature of this predicted leader/signal sequence, it is likely to act as a transmembrane domain should it not be separated from the remainder of the protein of SEQ ID NO:178. The TopPredII computer program predicts two additional potential  
10 transmembrane domains within the SEQ ID NO:178 amino acid sequence.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone lv2\_47 should be approximately 1950 bp.

The nucleotide sequence disclosed herein for lv2\_47 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and  
15 FASTA search protocols. lv2\_47 demonstrated at least some similarity with sequences identified as AA007293 (zh97f07.r1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 429253 5'), AA447347 (zw93g06.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 784570 5' similar to WP:F43E2.7 CE07243), AA522451 (ng30h09.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA clone IMAGE:936353), AA526614 (ni52g12.s1  
20 NCI\_CGAP\_Ov2 Homo sapiens cDNA clone 980518), F18178 (H.sapiens EST sequence (002-T4-28) from skeletal muscle, mRNA sequence), H46569 (yo20f10.s1 Homo sapiens cDNA clone 178507 3'), and T22574 (Human gene signature HUMGS04190). Based upon sequence similarity, lv2\_47 proteins and each similar protein or peptide may share at least some activity.

25

#### Clone "ml243\_1"

A polynucleotide of the present invention has been identified as clone "ml243\_1". ml243\_1 was isolated from a human adult brain (caudate nucleus) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No.  
30 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ml243\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ml243\_1 protein").

The nucleotide sequence of ml243\_1 as presently determined is reported in SEQ ID NO:37, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ml243\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:38. Amino acids 25 to 37 of SEQ ID NO:38 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 38. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ml243\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ml243\_1 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for ml243\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ml243\_1 demonstrated at least some similarity with sequences identified as N66656 (yy71a06.s1 Homo sapiens cDNA clone 278962 3'), R17513 (yg02g12.r1 Homo sapiens cDNA clone 31064 5'), Z83837 (Human DNA sequence from Fosmid 113D11 on chromosome 22q11.2-qter contains ESTs, CpG island), and Z84468 (Human DNA sequence from clone 299D3; HTGS phase 1). Based upon sequence similarity, ml243\_1 proteins and each similar protein or peptide may share at least some activity.

ml243\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 16 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

#### Clone "pm96\_9"

A polynucleotide of the present invention has been identified as clone "pm96\_9". pm96\_9 was isolated from a human fetal kidney (293 cell line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pm96\_9 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pm96\_9 protein").

The nucleotide sequence of pm96\_9 as presently determined is reported in SEQ ID NO:39, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pm96\_9 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:40.

5       The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pm96\_9 should be approximately 3600 bp.

The nucleotide sequence disclosed herein for pm96\_9 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pm96\_9 demonstrated at least some similarity with sequences  
10 identified as AA444024 (zv44d12.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 756503 5'), AA488901 (aa55h09.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:824897 3'), R16408 (yf40b02.r1 Homo sapiens cDNA clone 129291 5'), T19732 (Human gene signature HUMGS00806), U52112 (Homo sapiens Xq28 genomic DNA in the region of the L1CAM locus containing the genes for neural cell adhesion molecule L1  
15 (L1CAM), arginine-vasopressin receptor (AVPR2), C1 p115 (C1), ARD1 N-acetyltransferase related protein (TE2), renin-binding protein (RbP), host cell factor 1 (HCF1), and interleukin-1 receptor-associated kinase (IRAK) genes, complete cds, and Xq28lu2 gene), and Z82250 (Human DNA sequence from cosmid N86D4 on chromosome 22q12-qter contains STS). Based upon sequence similarity, pm96\_9 proteins and each similar protein  
20 or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain at the extreme C-terminus of the pm96\_9 protein sequence (SEQ ID NO:40).

#### Clone "pu261\_1"

25       A polynucleotide of the present invention has been identified as clone "pu261\_1". pu261\_1 was isolated from a human adult blood (promyelocytic leukemia HL-60) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein.  
30       pu261\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pu261\_1 protein").

The nucleotide sequence of pu261\_1 as presently determined is reported in SEQ ID NO:41, and includes a poly(A) tail. What applicants presently believe to be the proper

reading frame and the predicted amino acid sequence of the pu261\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:42. Amino acids 116 to 128 of SEQ ID NO:42 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 129. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pu261\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pu261\_1 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for pu261\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pu261\_1 demonstrated at least some similarity with sequences identified as H16093 (ym20g10.r1 Homo sapiens cDNA clone 48582 5'). Based upon sequence similarity, pu261\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the pu261\_1 protein sequence centered around amino acid 70 of SEQ ID NO:42.

#### Clone "pw214\_15"

A polynucleotide of the present invention has been identified as clone "pw214\_15". pw214\_15 was isolated from a human adult brain (cerebellum) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. pw214\_15 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pw214\_15 protein").

The nucleotide sequence of pw214\_15 as presently determined is reported in SEQ ID NO:43, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pw214\_15 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:44.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pw214\_15 should be approximately 1800 bp.

The nucleotide sequence disclosed herein for pw214\_15 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and

FASTA search protocols. pw214\_15 demonstrated at least some similarity with sequences identified as AA173391 (zp03a07.r1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone 595284 5'), AA253067 (zr52a10.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 667002 5'), AA523652 ni64d09.s1 NCI\_CGAP\_Pr12 Homo sapiens cDNA clone 981617), and H41832 (yo07b08.r1 Homo sapiens cDNA clone 177207 5'). Based upon sequence similarity, pw214\_15 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the pw214\_15 protein sequence centered around amino acid 15 of SEQ ID NO:44.

10

#### Clone "qb56\_19"

A polynucleotide of the present invention has been identified as clone "qb56\_19". qb56\_19 was isolated from a human adult bladder (carcinoma 5637) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qb56\_19 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qb56\_19 protein").

The nucleotide sequence of qb56\_19 as presently determined is reported in SEQ ID NO:45, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qb56\_19 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:46. Amino acids 18 to 40 of SEQ ID NO:46 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 41. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qb56\_19 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qb56\_19 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for qb56\_19 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qb56\_19 demonstrated at least some similarity with sequences identified as AA632658 (np87c12.s1 NCI\_CGAP\_Thy1 Homo sapiens cDNA clone



IMAGE:1133302), N56430 (JJ8973F Homo sapiens cDNA clone JJ8973 5'), and W05470 (za87f11.r1 Soares fetal lung NbHL19W Homo sapiens cDNA clone 299565 5'). Based upon sequence similarity, qb56\_19 proteins and each similar protein or peptide may share at least some activity.

- 5 qb56\_19 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 14 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "qc646\_1"

- 10 A polynucleotide of the present invention has been identified as clone "qc646\_1". qc646\_1 was isolated from a human adult neural tissue (neuroepithelioma HTB-10 line) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded
- 15 protein. qc646\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qc646\_1 protein").

- The nucleotide sequence of qc646\_1 as presently determined is reported in SEQ ID NO:47, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qc646\_1 protein
- 20 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Amino acids 12 to 24 of SEQ ID NO:48 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Amino acids 32 to 44 are also a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 45, or are a transmembrane domain. Due to the hydrophobic
- 25 nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the qc646\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qc646\_1 should be approximately 1800 bp.

- The nucleotide sequence disclosed herein for qc646\_1 was searched against the
- 30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qc646\_1 demonstrated at least some similarity with sequences identified as AA470035 (zt94a07.r1 Soares testis NHT Homo sapiens cDNA clone 729972 5'), and AA483957 (ne76e11.s1 NCI\_CGAP\_Ew1 Homo sapiens cDNA clone

IMAGE:910220). The predicted amino acid sequence disclosed herein for qc646\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted qc646\_1 protein demonstrated at least some similarity to sequences identified as D88666 (PS-PLA1 (serine phospholipid-specific phospholipase A) [Rattus norvegicus]), M93284 (lipase related protein 2 [Homo sapiens]), and R30739 (C-terminally truncated GPL(1-319)), as well as lipases from various other species. Rat PS-PLA1, serine phospholipid-specific phospholipase A, is a member of the lipase family and is secreted from activated platelets. Based upon sequence similarity, qc646\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the qc646\_1 protein sequence, one centered around amino acid 190 and another around amino acid 325 of SEQ ID NO:48. The nucleotide sequence of qc646\_1 indicates that it may contain Alu repetitive elements.

Clone "qf116\_2"

A polynucleotide of the present invention has been identified as clone "qf116\_2". qf116\_2 was isolated from a human adult bladder (carcinoma 5637) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qf116\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qf116\_2 protein").

The nucleotide sequence of qf116\_2 as presently determined is reported in SEQ ID NO:49, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qf116\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:50.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qf116\_2 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for qf116\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qf116\_2 demonstrated at least some similarity with sequences identified as D50810 (placental leucine aminopeptidase [Homo sapiens]), R94512 (GTVap (short version), insulin-cleaving aminopeptidase from GLUT-4 vesicles), and U32990

(vp165 [Rattus norvegicus]). Human placental leucine aminopeptidase/oxytocinase (P-LAP), a member of the type II membrane-spanning zinc metallopeptidase family, degrades several peptide hormones such as oxytocin and vasopresin, suggesting a role in maintaining homeostasis during pregnancy. The predicted P-LAP amino acid sequence  
5 contains the HEXXH consensus sequence of zinc metallopeptidases, indicating that the enzyme belongs to this family, which includes aminopeptidase N and aminopeptidase A. The deduced P-LAP amino acid sequence also contains a hydrophobic region near the N-terminus, suggesting that the enzyme is a type II integral membrane protein. Results suggest that the enzyme is synthesized as an integral membrane protein and is released  
10 into blood under some physiological conditions. (See Rogi *et al.*, 1996, *J. Biol. Chem.* 271(1): 56-61, which is incorporated by reference herein.) Based upon sequence similarity, qf116\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two potential transmembrane domains within the qf116\_2 protein sequence, one centered around amino acid 25 and another around amino  
15 acid 290 of SEQ ID NO:50.

#### Clone "qf662\_3"

A polynucleotide of the present invention has been identified as clone "qf662\_3". qf662\_3 was isolated from a human adult bladder (carcinoma 5637) cDNA library using  
20 methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. qf662\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "qf662\_3 protein").

25 The nucleotide sequence of qf662\_3 as presently determined is reported in SEQ ID NO:51, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the qf662\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:52. Amino acids 133 to 145 of SEQ ID NO:52 are a predicted leader/signal sequence, with the  
30 predicted mature amino acid sequence beginning at amino acid 146. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the qf662\_3 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone qf662\_3 should be approximately 1000 bp.

The nucleotide sequence disclosed herein for qf662\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. qf662\_3 demonstrated no significant similarity with sequences in these databases. The nucleotide sequence of qf662\_3 indicates that it may contain repetitive elements.

Clone "am748\_5"

A polynucleotide of the present invention has been identified as clone "am748\_5". am748\_5 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. am748\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "am748\_5 protein").

The nucleotide sequence of am748\_5 as presently determined is reported in SEQ ID NO:53, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the am748\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:54. Amino acids 14 to 26 of SEQ ID NO:54 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the am748\_5 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone am748\_5 should be approximately 1550 bp.

The nucleotide sequence disclosed herein for am748\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. am748\_5 demonstrated at least some similarity with sequences identified as AA418860 (zv98g04.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 767862 5' similar to gb:X14008\_ma1 LYSOZYME C PRECURSOR (HUMAN); contains Alu repetitive element; contains element PTR5 repetitive element), AC003007 (Human

Chromosome 16 BAC clone CIT987SK-A-61E3, complete sequence), H73304 (yu27c10.r1 Homo sapiens cDNA clone 235026 5' similar to contains Alu repetitive element), N35175 (yx83d10.r1 Homo sapiens cDNA clone 268339 5' similar to gb X14008\_ma1 LYSOZYME C PRECURSOR (HUMAN); contains Alu repetitive element),  
5 N41479 (yy05a11.r1 Homo sapiens cDNA clone 270332 5' similar to gb:X14008\_ma1 LYSOZYME C PRECURSOR (HUMAN)), Q81139 (HPLA2-8 gene), T04964 (EST02852 Homo sapiens cDNA clone HFBCI77), and U18391 (Human Alu sequence clone A8). The predicted amino acid sequence disclosed herein for am748\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol.  
10 The predicted am748\_5 protein demonstrated at least some similarity to sequences identified as X55777 (put. ORF [Homo sapiens]) and R13556 (Protein encoded downstream of hhc\_M oncoprotein). Based upon sequence similarity, am748\_5 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of am748\_5 indicates that it may contain one or more of the following repetitive  
15 elements: Alu, L1.

#### Clone "cj507\_1"

A polynucleotide of the present invention has been identified as clone "cj507\_1". cj507\_1 was isolated from a human fetal brain cDNA library using methods which are  
20 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cj507\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cj507\_1 protein").

25 The nucleotide sequence of cj507\_1 as presently determined is reported in SEQ ID NO:55, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cj507\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
30 cj507\_1 should be approximately 2100 bp.

The nucleotide sequence disclosed herein for cj507\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cj507\_1 demonstrated at least some similarity with sequences

identified as AA100356 (zn46a02.r1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 550442 5' similar to contains element PTR5 repetitive element), AA228100 (zr56g04.s1 Soares NhHMPu S1 Homo sapiens cDNA clone 667446 3'), AA479997 (zv18b07.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 753973 5' similar to contains  
5 element PTR5 repetitive element, mRNA sequence), and X85324 (H.sapiens mRNA for non polymorphic CAG repeat (CAG12)). Based upon sequence similarity, cj507\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the cj507\_1 protein sequence centered around amino acid 265 of SEQ ID NO:56. The  
10 nucleotide sequence of cj507\_1 indicates that it may contain a GCA simple repeat region.

cj507\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 47 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

15      Clone "cn922\_5"

A polynucleotide of the present invention has been identified as clone "cn922\_5". cn922\_5 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
20 analysis of the amino acid sequence of the encoded protein. cn922\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cn922\_5 protein").

The nucleotide sequence of cn922\_5 as presently determined is reported in SEQ ID NO:57, and includes a poly(A) tail. What applicants presently believe to be the proper  
25 reading frame and the predicted amino acid sequence of the cn922\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:58.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cn922\_5 should be approximately 2200 bp.

The nucleotide sequence disclosed herein for cn922\_5 was searched against the  
30 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cn922\_5 demonstrated at least some similarity with sequences identified as H34191 (EST110864 Rattus sp. cDNA 5' end), R18707 (yf98f02.r1 Homo sapiens cDNA clone 30546 5'), T26556 (Human gene signature HUMGS08801), and

Z83230 (*Caenorhabditis elegans* cosmid F56A8). The predicted amino acid sequence disclosed herein for cn922\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cn922\_5 protein demonstrated at least some similarity to sequences identified as AB004535  
5 (HYPOTHETICAL 105.9 KD PROTEIN IN AAC3-RFC5 INTERGENIC REGION [Schizosaccharomyces pombe]) and Z83230 (F56A8.a and F56A8.1 [*Caenorhabditis elegans*]). Based upon sequence similarity, cn922\_5 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts six potential transmembrane domains within the cn922\_5 protein sequence, centered around  
10 amino acids 25, 100, 135, 190, 290, and 370 of SEQ ID NO:58, respectively. The nucleotide sequence of cn922\_5 indicates that it may contain one or more of the following repetitive elements: MER, L1.

Clone "cw691\_11"

15 A polynucleotide of the present invention has been identified as clone "cw691\_11". cw691\_11 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cw691\_11 is a full-length  
20 clone, including the entire coding sequence of a secreted protein (also referred to herein as "cw691\_11 protein").

The nucleotide sequence of cw691\_11 as presently determined is reported in SEQ ID NO:59, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cw691\_11 protein  
25 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:60.

Another potential cw691\_11 reading frame and predicted amino acid sequence is encoded by basepairs 542 to 970 of SEQ ID NO:59 and is reported in SEQ ID NO:179. Amino acids 34 to 46 of SEQ ID NO:179 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 47. Due to the  
30 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the protein of SEQ ID NO:179.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw691\_11 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for cw691\_11 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw691\_11 demonstrated at least some similarity with sequences identified as AA363712 (EST74158 Pancreas I Homo sapiens cDNA 5' end similar to similar to C. elegans hypothetical protein R10E12.1), AA521201 (aa74c10.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone 826674 3'), AA527142 (ni07a10.s1 NCI\_CGAP\_Br2 Homo sapiens cDNA clone IMAGE 967290, mRNA sequence), AA745501 (ny64d03.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1283045, mRNA sequence), N73108 (yv69a09.r1 Homo sapiens cDNA clone 247960 5'), T19938 (Human gene signature HUMGS01070), and W77963 (zd70d09.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 346001 5' similar to WP:R10E12.1 CE00310). The predicted amino acid sequence disclosed herein for cw691\_11 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cw691\_11 protein demonstrated at least some similarity to sequences identified as P82971 (Bioadhesive precursor protein from cDNA 52), U73679 (YNK1-a [Caenorhabditis elegans]), and Z29561 (R10E12.1 [Caenorhabditis elegans]). Based upon sequence similarity, cw691\_11 proteins and each similar protein or peptide may share at least some activity.

#### Clone "cw1000\_2"

A polynucleotide of the present invention has been identified as clone "cw1000\_2". cw1000\_2 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cw1000\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cw1000\_2 protein").

The nucleotide sequence of cw1000\_2 as presently determined is reported in SEQ ID NO:61, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cw1000\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:62. Amino



acids 24 to 36 of SEQ ID NO:62 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 37. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the cw1000\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw1000\_2 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for cw1000\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw1000\_2 demonstrated at least some similarity with sequences identified as AA446779 (zw89d11.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 784149 5', mRNA sequence), AA493561 (nh04f07.s1 NCI\_CGAP\_Thy1 Homo sapiens cDNA clone 943333 similar to WP:F15G9.4 CE01552 IG SUPERFAMILY REPEATS ;contains element MSR1 repetitive element), H35690 (EST111696 Rattus sp. cDNA similar to Opioid binding protein/cell adhesion-like molecule), R18502 (yf96a05.r1 Homo sapiens cDNA clone 30376 5'), T21582 (Human gene signature HUMGS02965), T39504 (ya06g11.r1 Homo sapiens cDNA clone 60740 5'), T46848 (yb94b01.r1 Homo sapiens cDNA clone 78793 5'), T51129 (yb94b01.s1 Homo sapiens cDNA clone 78793 3'), and W67535 (zd40g11.s1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 343172 3' similar to PIR S05539 S05539 glycophorin C - human ;contains element MSR1 repetitive element). The predicted amino acid sequence disclosed herein for cw1000\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cw1000\_2 protein demonstrated at least some similarity to sequences identified as M24406 (poliovirus receptor [Homo sapiens]), R07130 (H20B receptor), W04404 (Human CRTAM; Cytotoxic or Regulatory T-cell associated Mol.; CRTAM), X13890 (glycophorin C [Homo sapiens]), and X90569 (elastic titin [Homo sapiens]). Based upon sequence similarity, cw1000\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the cw1000\_2 protein sequence centered around amino acid 358 of SEQ ID NO:62. The nucleotide sequence of cw1000\_2 indicates that it may contain a GCC1 repeat element.

cw1000\_2 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 57 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5           Clone "cw1640\_1"

A polynucleotide of the present invention has been identified as clone "cw1640\_1". cw1640\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
10 analysis of the amino acid sequence of the encoded protein. cw1640\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cw1640\_1 protein").

The nucleotide sequence of cw1640\_1 as presently determined is reported in SEQ ID NO:63, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the cw1640\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:64. Amino acids 123 to 135 of SEQ ID NO:64 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 136. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a  
20 transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the cw1640\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw1640\_1 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for cw1640\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw1640\_1 demonstrated at least some similarity with sequences identified as AA075643 (zm88a12.r1 Stratagene ovarian cancer (#937219) Homo sapiens cDNA clone 544990 5' similar to SW:ACT\_EUPCR P20360 ACTIN), AA411334 (zv29e11.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 755084 5' similar to  
30 WP:C49H3.8 CE04234 ACTIN-LIKE PROTEIN ), AA913364 (ol37b07.s1 Soares NFL\_T\_GBC\_S1 Homo sapiens cDNA clone IMAGE:1525621 3' similar to WP:C49H3.8 CE04234 ACTIN-LIKE PROTEIN, mRNA sequence), N25416 (yx40g10.r1 Homo sapiens cDNA clone 264258 5' similar to SP ACT2\_PLAFA P14883 ACTIN), R96887

(yq61g10.r1 Homo sapiens cDNA clone 200322 5'), W37097 (zb98h03.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 320885 5'), W44778 (zb98h03.s1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 320885 3'), W61038 (zc54g09.r1 Soares senescent fibroblasts NbHSF Homo), W76570 (zd66f12.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 345647 5' similar to SW:ACT\_PROCL P45521 ACTIN), and W82519 (mf05b01.r1 Soares mouse p3NMF19.5 Mus musculus cDNA clone). The predicted amino acid sequence disclosed herein for cw1640\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cw1640\_1 protein demonstrated at least some similarity to sequences identified as J00068 (alpha-actin [Homo sapiens]), J01163 (actin [Oxytricha fallax]), R22026 (A. chrysogenum actin), R50328 (Drug resistant structural protein), U42436 (Similar to actin-like protein [Caenorhabditis elegans]), and U90439 (actin isolog [Arabidopsis thaliana]). Based upon sequence similarity, cw1640\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "d24\_1"

A polynucleotide of the present invention has been identified as clone "d24\_1". A cDNA clone was first isolated from a human adult blood (peripheral blood mononuclear cells treated with concanavalin A and phorbol myristate acetate) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate d24\_1 from a human adult blood (peripheral blood mononuclear cells treated with phytohemagglutinin, phorbol myristate acetate, and mixed lymphocyte reaction) cDNA library. d24\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "d24\_1 protein").

The nucleotide sequence of d24\_1 as presently determined is reported in SEQ ID NO:65, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the d24\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:66. Amino acids 124 to 136 of SEQ ID NO:66 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 137. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should

the predicted leader/signal sequence not be separated from the remainder of the d24\_1 protein. The mRNA sequence encoding amino acids 172 to 175 of SEQ ID NO:66 may not be present in alternatively-spliced forms of d24\_1 mRNA molecules.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
5 d24\_1 should be approximately 2000 bp.

The nucleotide sequence disclosed herein for d24\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. d24\_1 demonstrated at least some similarity with sequences identified as AA478740 (zv14g12.s1 Soares NhHMPu S1 Homo sapiens cDNA clone  
10 753670 3'), AA479444 (zv14g12.r1 Soares NhHMPu S1 Homo sapiens cDNA clone 753670 5', mRNA sequence), AA278581 (zs76f09.r1 Soares NbHTGBC Homo sapiens cDNA clone 703433 5' similar to WP T04A8.12 CE01067 YEAST 107.9KD PGK1-MAK32 INTERGENIC HYPOTHETICAL PROTEIN), H05202 (yl85h02.r1 Homo sapiens cDNA clone 45213 5' similar to SP T04A8.12m CE01067 YEAST 107.9KD  
15 PGK1-MAK32 INTERGENIC HYPOTHETICAL PROTEIN), R74287 (yi57e07.r1 Homo sapiens cDNA clone 143364 5'), U57715 (*Rattus norvegicus* FGF receptor activating protein FRAG1 (FRAG1) mRNA, complete CDs), and Z35663 (*C. elegans* protein of unknown function). The predicted amino acid sequence disclosed herein for d24\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
20 BLASTX search protocol. The predicted d24\_1 protein demonstrated at least some similarity to the sequence identified as U57715 (FGF receptor activating protein FRAG1 [*Rattus norvegicus*]). Lorenzi *et al.* (1996, *Proc. Natl. Acad. Sci. USA* 93:8956, incorporated by reference herein) studied the FRAG1 gene in rat osteosarcoma cells. They concluded that the FRAG1 gene product gets fused to FGF receptor 2 (FGFR2). This  
25 fusion "drastically stimulates the transforming activity and autophosphorylation of the receptor" and causes oncogenicity. Based upon sequence similarity, d24\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the d24\_1 protein sequence, centered around amino acids 34, 154, and 194 of SEQ ID NO:66,  
30 respectively.

d24\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 24 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5           Clone "dd426\_1"

A polynucleotide of the present invention has been identified as clone "dd426\_1". A cDNA clone was first isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
10 analysis of the amino acid sequence of the encoded protein. This cDNA clone was then used to isolate dd426\_1 from a human adult testes (teratocarcinoma NCCIT) cDNA library. dd426\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dd426\_1 protein").

The nucleotide sequence of dd426\_1 as presently determined is reported in SEQ  
15 ID NO:67, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dd426\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:68. Amino acids 76 to 88 of SEQ ID NO:68 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 89. Due to the hydrophobic nature  
20 of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the dd426\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dd426\_1 should be approximately 800 bp.

25           The nucleotide sequence disclosed herein for dd426\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dd426\_1 demonstrated at least some similarity with sequences identified as AA760716 (nz13d06.s1 NCI\_CGAP\_GCB1 Homo sapiens cDNA clone IMAGE:1287659 similar to WP:F13H10.3 CE05624 YEAST YEH4 LIKE PROTEIN;  
30 mRNA sequence), H11919 (ym10e10.r1 Homo sapiens cDNA clone 47462 5'), and Z68748 (Caenorhabditis elegans cosmid F13H10). The predicted amino acid sequence disclosed herein for dd426\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dd426\_1 protein

demonstrated at least some similarity to sequences identified as U39782 (lysine and histidine specific transporter [*Arabidopsis thaliana*]) and Z68748 (F13H10.3 [*Caenorhabditis elegans*]). Based upon sequence similarity, dd426\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program  
5 predicts an additional potential transmembrane domain within the dd426\_1 protein sequence centered around amino acid 30 of SEQ ID NO:68, which may also function as a leader/signal sequence.

dd426\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 12 kDa was detected in membrane fractions using SDS  
10 polyacrylamide gel electrophoresis.

#### Clone "di393\_2"

A polynucleotide of the present invention has been identified as clone "di393\_2". di393\_2 was isolated from a human adult testes cDNA library using methods which are  
15 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. di393\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "di393\_2 protein").

20 The nucleotide sequence of di393\_2 as presently determined is reported in SEQ ID NO:69, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the di393\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:70. Amino acids 7 to 19 of SEQ ID NO:70 are a predicted leader/signal sequence, with the predicted  
25 mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the di393\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
30 di393\_2 should be approximately 600 bp.

The nucleotide sequence disclosed herein for di393\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. di393\_2 demonstrated at least some similarity with sequences

identified as AA669506 (zu85g08.s1 Soares testis NHT Homo sapiens cDNA clone 744830 3', mRNA sequence). Based upon sequence similarity, di393\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the di393\_2 protein sequence centered around amino acid 66 of SEQ ID NO:70.

di393\_2 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 20 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "dj167\_2"

A polynucleotide of the present invention has been identified as clone "dj167\_2". dj167\_2 was isolated from a human adult placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dj167\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dj167\_2 protein").

The nucleotide sequence of dj167\_2 as presently determined is reported in SEQ ID NO:71, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dj167\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:72.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dj167\_2 should be approximately 1550 bp.

The nucleotide sequence disclosed herein for dj167\_2 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dj167\_2 demonstrated at least some similarity with sequences identified as H49161 (yq18d05.r1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone 274208 5'), L12350 (Human thrombospondin 2 (THBS2) mRNA, complete cds), T98917 (ye66b03.s1 Homo sapiens cDNA clone 122669 3' similar to SP:TSP1\_CHICK P35440 THROMBOSPONDIN 1), and X87620 (B.taurus mRNA for complete thrombospondin). The predicted amino acid sequence disclosed herein for dj167\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dj167\_2 protein demonstrated at least some

similarity to sequences identified as L12350 (thrombospondin 2 [Homo sapiens]), M60853 (thrombospondin [Gallus gallus]), R40823 (Human thrombospondin 1), U48245 (protein kinase C-binding protein Nel [Rattus norvegicus]), X87620 (thrombospondin [Bos taurus]), and Z71178 (B0024.14 [Caenorhabditis elegans]). Based upon sequence  
5 similarity, dj167\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three potential transmembrane domains within the dj167\_2 protein sequence, centered around amino acids 140, 215, and 315 of SEQ ID NO:72, respectively.

10       Clone "dj167\_19"

A polynucleotide of the present invention has been identified as clone "dj167\_19". dj167\_19 was isolated from a human adult placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
15 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dj167\_19 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dj167\_19 protein").

The nucleotide sequence of dj167\_19 as presently determined is reported in SEQ ID NO:73, and includes a poly(A) tail. What applicants presently believe to be the proper  
20 reading frame and the predicted amino acid sequence of the dj167\_19 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:74. Amino acids 22 to 34 of SEQ ID NO:74 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 35. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
25 should the predicted leader/signal sequence not be separated from the remainder of the dj167\_19 protein. The dj167\_19 clone is related to that of dj167\_2, and extends further 5'. The dj167\_19 clone appears to contain coding sequences for chorionic somatomammotropin in the opposite orientation at its 5' end between Sfi restriction sites (at nucleotides 16 and 839 of SEQ ID NO:73). The dj167\_2 and dj167\_19 clones may represent  
30 alternatively spliced messenger RNA molecules encoding two different forms of a secreted protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dj167\_19 should be approximately 4500 bp.



Analysis of the dj167\_19 amino acid sequence (SEQ ID NO:74) reveals the following domains: IGFBP cysteine-rich domain at amino acids 60-75; VWF-B cysteine-rich domains at amino acids 174-210, 212-247, 255-291, and 293-328; Chordin cysteine-rich domains at amino acids 336-390, 403-456, 608-662, 679-734, 753-808, and 819-873; Antistatin (protease inhibitor) cysteine-rich domains at amino acids 469-498, 505-532, 539-564, and 567-592; RGD cell attachment sequence at amino acids 314-316, and Asn glycosylation sites at amino acids 71, 113, 330, 474, and 746. The cysteine-rich domains listed above are similar to domains found in the C domain of Von Willebrand Factor (VWF), and in procollagen and thrombospondin. In addition, the amino acid sequence of SEQ ID NO:74 from amino acid 938 to amino acid 960 appears to be a transmembrane domain.

The dj167\_19 transcript is expressed in several cell types, including kidney, pancreas, spleen, and ovary, and is most abundantly expressed in placental tissue.

#### Clone "dw665\_4"

A polynucleotide of the present invention has been identified as clone "dw665\_4". dw665\_4 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dw665\_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dw665\_4 protein").

The nucleotide sequence of dw665\_4 as presently determined is reported in SEQ ID NO:75, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dw665\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76. Amino acids 15 to 27 of SEQ ID NO:76 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Amino acids 16 to 28 of SEQ ID NO:76 are also a predicted leader/signal sequence, with the predicted mature amino acid sequence in that case beginning at amino acid 29. Due to the hydrophobic nature of these predicted leader/signal sequences, each is likely to act as a transmembrane domain should it not be separated from the remainder of the dw665\_4 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dw665\_4 should be approximately 3750 bp.

The nucleotide sequence disclosed herein for dw665\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dw665\_4 demonstrated at least some similarity with sequences identified as AA029053 (zk09f06.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 470051 3'), H77289 (EST27017 WATM1 Homo sapiens cDNA clone 27017, mRNA sequence), and T21722 (Human gene signature HUMGS03170). The predicted amino acid sequence disclosed herein for dw665\_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dw665\_4 protein demonstrated at least some similarity to sequences identified as L35764 (chordin [Xenopus laevis]) and W31559 (Xenopus frog protein "chordin"). Analysis of motifs within the predicted dw665\_4 protein revealed the presence of Chordin cysteine-rich domains at amino acids 37-99, 115-178, and 260-322 of SEQ ID NO:76; an 'RGD' cell-attachment sequence (at amino acids 179-181 of SEQ ID NO:76), which in some proteins has been shown to play a role in cell adhesion; and Asp glycosylation sites at amino acids 118 and 291. Based upon sequence similarity, dw665\_4 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of dw665\_4 indicates that it may contain an AC repetitive element.

dw665\_4 transcripts are expressed in many tissues including kidney, adrenal gland, and prostate tissues, and are most abundantly expressed in pancreas; however, little or no dw665\_4 transcript expression is observed in liver or peripheral blood cells. dw665\_4 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 75 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis; two additional bands at approximately 26 and 30 kDa were also observed. BIACORE binding experiments indicate that dw665\_4 protein has a Chordin-like protein-binding profile, and binds to BMP-2, BMP-4, BMP-7, BMP-12, and GDF-5.

#### Clone "dx146\_12"

A polynucleotide of the present invention has been identified as clone "dx146\_12". dx146\_12 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dx146\_12 is a full-length

clone, including the entire coding sequence of a secreted protein (also referred to herein as "dx146\_12 protein").

The nucleotide sequence of dx146\_12 as presently determined is reported in SEQ ID NO:77, and includes a poly(A) tail. What applicants presently believe to be the proper  
5 reading frame and the predicted amino acid sequence of the dx146\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dx146\_12 should be approximately 2250 bp.

The nucleotide sequence disclosed herein for dx146\_12 was searched against the  
10 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dx146\_12 demonstrated at least some similarity with sequences identified as AA090429 (y0527.seq.F Fetal heart, Lambda ZAP Express Homo sapiens cDNA 5'), AA232068 (zr24a01.r1 Stratagene NT2 neuronal precursor 937230 Homo sapiens cDNA clone 664296 5'), AA886679 (oj47h07.s1 NCI\_CGAP\_Kid3 Homo sapiens  
15 cDNA clone IMAGE:1501501 3' similar to WP:F16A11.2 CE09424 METHANOCOCCUS HYPOTHETICAL PROTEIN 0682 LIKE; mRNA sequence), R61436 (yh15g06.r1 Homo sapiens cDNA clone 37884 5'), and Z81505 (Caenorhabditis elegans cosmid F16A11, complete sequence). The predicted amino acid sequence disclosed herein for dx146\_12 was searched against the GenPept and GeneSeq amino acid sequence  
20 databases using the BLASTX search protocol. The predicted dx146\_12 protein demonstrated at least some similarity to sequences identified as U67515 (hypothetical protein (SP P46850) [Methanococcus jannaschii]) and Z81505 (F16A11.2 [Caenorhabditis elegans]). Based upon sequence similarity, dx146\_12 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a  
25 potential transmembrane domain within the dx146\_12 protein sequence centered around amino acid 405 of SEQ ID NO:78.

dx146\_12 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 50 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

30

Clone "dx219\_13"

A polynucleotide of the present invention has been identified as clone "dx219\_13". dx219\_13 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
5 analysis of the amino acid sequence of the encoded protein. dx219\_13 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dx219\_13 protein").

The nucleotide sequence of dx219\_13 as presently determined is reported in SEQ ID NO:79, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the dx219\_13 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:80. Amino acids 94 to 106 of SEQ ID NO:80 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 107. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a  
15 transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the dx219\_13 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dx219\_13 should be approximately 1200 bp.

The nucleotide sequence disclosed herein for dx219\_13 was searched against the  
20 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. dx219\_13 demonstrated at least some similarity with sequences identified as AA429731 (zw66g05.s1 Soares testis NHT Homo sapiens cDNA clone 781208 3'), AA446067 (zw66e06.r1 Soares testis NHT Homo sapiens cDNA clone 781186 5', mRNA sequence), T23212 (standard; cDNA to mRNA; 161 BP, Human gene signature  
25 HUMGS05005), W29299 (mb99f03.r1 Soares mouse p3NMF19.5 Mus musculus cDNA clone 337565 5'), W87852 (zh68b05.r1 Soares fetal liver spleen 1NFLS S1 Homo sapiens cDNA clone 417201 5'), and Y13897 (Homo sapiens partial mRNA for hypothetical protein). Based upon sequence similarity, dx219\_13 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two  
30 additional potential transmembrane domains within the dx219\_13 protein sequence, one centered around amino acid 160 and another around amino acid 275 of SEQ ID NO:80.

dx219\_13 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 37 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

5           Clone "fm3\_1"

A polynucleotide of the present invention has been identified as clone "fm3\_1". fm3\_1 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
10 analysis of the amino acid sequence of the encoded protein. fm3\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "fm3\_1 protein").

The nucleotide sequence of fm3\_1 as presently determined is reported in SEQ ID NO:81, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the fm3\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:82. Amino acids 7 to 19 of SEQ ID NO:82 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
20 the predicted leader/signal sequence not be separated from the remainder of the fm3\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone fm3\_1 should be approximately 600 bp.

The nucleotide sequence disclosed herein for fm3\_1 was searched against the  
25 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. fm3\_1 demonstrated at least some similarity with sequences identified as T15669 (IB1718 Infant brain, Bento Soares Homo sapiens cDNA 3'end). Based upon sequence similarity, fm3\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional  
30 potential transmembrane domains within the fm3\_1 protein sequence centered around amino acid 85 of SEQ ID NO:82.

fm3\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 9 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "h225\_1"

A polynucleotide of the present invention has been identified as clone "h225\_1". h225\_1 was isolated from a human adult blood (peripheral blood mononuclear cells  
5 treated with phytohemagglutinin and phorbol myristate acetate and mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. h225\_1 is a full-length clone, including the entire coding sequence  
10 of a secreted protein (also referred to herein as "h225\_1 protein").

The nucleotide sequence of h225\_1 as presently determined is reported in SEQ ID NO:83. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the h225\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:84. Amino acids 52 to 64 of SEQ ID NO:84  
15 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 65. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the h225\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
20 h225\_1 should be approximately 832 bp.

The nucleotide sequence disclosed herein for h225\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. h225\_1 demonstrated at least some similarity with sequences identified as AA604374 (no87e01.s1 NCI\_CGAP\_AA1 Homo sapiens cDNA clone  
25 IMAGE:1113816 similar to WP:ZK757.1 CE00467; mRNA sequence), H18393 (yn49a12.r1 Homo sapiens cDNA clone 171742 5' similar to SP:ZK757.1 CE00467), and R23642 (yh35e03.r1 Homo sapiens cDNA clone 131740 5'). The predicted amino acid sequence disclosed herein for h225\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted h225\_1  
30 protein demonstrated at least some similarity to sequences identified as AL022600 (hypothetical protein [Schizosaccharomyces pombe]) and Z48758 (SC9727\_21 unknown [Saccharomyces cerevisiae]). Based upon sequence similarity, h225\_1 proteins and each similar protein or peptide may share at least some activity.

Clone "kj320\_1"

A polynucleotide of the present invention has been identified as clone "kj320\_1". kj320\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
5 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. kj320\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "kj320\_1 protein").

The nucleotide sequence of kj320\_1 as presently determined is reported in SEQ ID  
10 NO:85, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the kj320\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:86. Amino acids 26 to 38 of SEQ ID NO:86 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 39. Due to the hydrophobic nature  
15 of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the kj320\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone kj320\_1 should be approximately 4900 bp.

20 The nucleotide sequence disclosed herein for kj320\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. kj320\_1 demonstrated at least some similarity with sequences identified as A45343 (Sequence 13 from Patent WO9517522), AA284111 (zc36f08.T7 Soares senescent fibroblasts NbHSF Homo sapiens cDNA clone 324423 3' similar to WP  
25 ZK688.8 CE00544 UDP-GALNAC; mRNA sequence), AA375707 (EST88026 HSC172 cells II Homo sapiens cDNA 5' end), AA534406 (nf76b08.s1 NCI\_CGAP\_Co3 Homo sapiens cDNA clone IMAGE 925815), D39885 (Rice cDNA, partial sequence (S1531\_1A)), G10010 (human STS CHLC.GCT16E06.P18287 clone GCT16E06), Q75104 (Cattle GalNAc-transferase), Q95187 (Simple tandem repeat (STR) corresponding  
30 to wg1d10), and U35890 (Rattus norvegicus polypeptide GalNAc transferase T1 mRNA, complete cds). The predicted amino acid sequence disclosed herein for kj320\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted kj320\_1 protein demonstrated at least some

similarity to sequences identified as R66397 (Cattle GalNAc-transferase), U41514 (UDP-GalNAc polypeptide N-acetylgalactosaminyltransferase [Homo sapiens]), and X85018 (UDP-GalNAc polypeptide N-acetylgalactosaminyl transferase [Homo sapiens]). Analysis of motifs within kj320\_1 reveals the presence of the alpha-2-macroglobulin family thiolester region signature. The proteinase-binding alpha-macroglobulins (A2M) are large glycoproteins found in the plasma of vertebrates, in the hemolymph of some invertebrates, and in reptilian and avian egg white. They inhibit all four classes of proteinases by trapping a proteinase with a peptide stretch containing the specific cleavage site (the 'bait' region) which upon proteinase binding induces a conformational change in the protein, trapping the proteinase. Upon cleavage of the 'bait' region, a covalent bond (a thiol-ester bond between the side chains of a cysteine and a glutamine) is formed between the A2M and the proteinase. Based upon sequence similarity, kj320\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of kj320\_1 indicates that it may contain one or more repetitive elements.

kj320\_1 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 136 kDa was detected in conditioned medium using SDS polyacrylamide gel electrophoresis.

#### Clone "ml236\_5"

A polynucleotide of the present invention has been identified as clone "ml236\_5". ml236\_5 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. ml236\_5 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ml236\_5 protein").

The nucleotide sequence of ml236\_5 as presently determined is reported in SEQ ID NO:87, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ml236\_5 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:88. Amino acids 148 to 160 of SEQ ID NO:88 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 161. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a



transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ml236\_5 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ml236\_5 should be approximately 1300 bp.

5        The nucleotide sequence disclosed herein for ml236\_5 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ml236\_5 demonstrated at least some similarity with sequences identified as AA137204 (zl23h11.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 502821 3'), AA307966 (EST17887 Aorta endothelial cells, TNF alpha-treated Homo sapiens cDNA 5' end, mRNA sequence), AA434504 (zw31c03.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 770884 5' similar to WP C45G9.7 CE01858), AA525971 (ni93g09.s1 NCI\_CGAP\_Pr21 Homo sapiens cDNA clone 984448), AA526490 (ni96c11.s1 NCI\_CGAP\_Pr21 Homo sapiens cDNA clone IMAGE 984692, mRNA sequence), AF028823 (Homo sapiens Tax interaction protein 1 mRNA, partial cds), U90913 (Human clone 23665 mRNA sequence), and W73114 (zd55c12.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 344566 5'). The predicted amino acid sequence disclosed herein for ml236\_5 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ml236\_5 protein demonstrated at least some similarity to sequences identified as AF028823 (Tax interaction protein 1 [Homo sapiens]) and U21323 (similar to tight junction protein (ZO-1) (SP Z01\_HUMAN, Q07157) [Caenorhabditis elegans]). Based upon sequence similarity, ml236\_5 proteins and each similar protein or peptide may share at least some activity.

25        ml236\_5 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 14 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "pu282\_10"

30        A polynucleotide of the present invention has been identified as clone "pu282\_10". pu282\_10 was isolated from a human adult blood (promyelocytic leukemia HL-60) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on

the basis of computer analysis of the amino acid sequence of the encoded protein. pu282\_10 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "pu282\_10 protein").

5 The nucleotide sequence of pu282\_10 as presently determined is reported in SEQ ID NO:89, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the pu282\_10 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:90. Amino acids 119 to 131 of SEQ ID NO:90 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 132. Due to the  
10 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the pu282\_10 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone pu282\_10 should be approximately 1050 bp.

15 The nucleotide sequence disclosed herein for pu282\_10 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. pu282\_10 demonstrated at least some similarity with sequences identified as AA311503 (EST182442 Jurkat T-cells VI Homo sapiens cDNA 5' end), AA336709 (EST41341 Endometrial tumor Homo sapiens cDNA 5' end), AA336890  
20 (EST41572 Endometrial tumor), AA385588 (EST99290 Thyroid Homo sapiens cDNA 5' end), AA526889 (ni09e05.s1 NCI\_CGAP\_Br2 Homo sapiens cDNA clone IMAGE:967520), AC003058 (Arabidopsis thaliana "unknown" protein), and T19726 (Human gene signature HUMGS00800). Based upon sequence similarity, pu282\_10 proteins and each similar protein or peptide may share at least some activity. The  
25 TopPredII computer program predicts two additional potential transmembrane domains within the pu282\_10 protein sequence, one centered around amino acid 39 and another around amino acid 95 of SEQ ID NO:90.

pu282\_10 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 16 kDa was detected in conditioned medium  
30 and membrane fractions using SDS polyacrylamide gel electrophoresis.

Clone "at94\_2"

A polynucleotide of the present invention has been identified as clone "at94\_2". at94\_2 was isolated from a human adult blood (lymphocytes and dendritic cells treated with mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as  
5 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. at94\_2 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "at94\_2 protein").

The nucleotide sequence of at94\_2 as presently determined is reported in SEQ ID NO:91, and includes a poly(A) tail. What applicants presently believe to be the proper  
10 reading frame and the predicted amino acid sequence of the at94\_2 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:92. Amino acids 214 to 226 of SEQ ID NO:92 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 227. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should  
15 the predicted leader/signal sequence not be separated from the remainder of the at94\_2 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone at94\_2 should be approximately 4300 bp.

The nucleotide sequence disclosed herein for at94\_2 was searched against the  
20 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. at94\_2 demonstrated at least some similarity with sequences identified as N24317 (yx23d12.r1 Homo sapiens cDNA clone 262583 5'), T30988 (EST25695 Homo sapiens cDNA 5' end similar to None), and U37026 (Rattus norvegicus brain sodium channel beta 2 subunit (SCNB2) mRNA, complete cds). The predicted amino  
25 acid sequence disclosed herein for at94\_2 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted at94\_2 protein demonstrated at least some similarity to the sequence identified as Z49912 (T24F1.2 [Caenorhabditis elegans]). Based upon sequence similarity, at94\_2 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer  
30 program predicts four additional potential transmembrane domains within the at94\_2 protein sequence, centered around amino acids 23, 306, 332, and 364 of SEQ ID NO:92, respectively.

Clone "bf169\_13"

A polynucleotide of the present invention has been identified as clone "bf169\_13". bf169\_13 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
5 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bf169\_13 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "bf169\_13 protein").

The nucleotide sequence of bf169\_13 as presently determined is reported in SEQ  
10 ID NO:93, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bf169\_13 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:94. Amino acids 342 to 354 are a possible leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 355. Due to the hydrophobic nature of this  
15 possible leader/signal sequence, it is likely to act as a transmembrane domain should it not be separated from the remainder of the bf169\_13 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bf169\_13 should be approximately 3000 bp.

The nucleotide sequence disclosed herein for bf169\_13 was searched against the  
20 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bf169\_13 demonstrated at least some similarity with sequences identified as AA227952 (zr56b06.s1 Soares NhHMPu S1 Homo sapiens cDNA clone 667379 3'), AA453914 (zx32e11.r1 Soares total fetus Nb2HF8 9w Homo sapiens cDNA clone 788204 5' similar to contains element TAR1 repetitive element; mRNA sequence),  
25 H46157 (yo13f11.r1 Homo sapiens cDNA clone 177837 5'), H18792 (yn52e02.r1 Homo sapiens cDNA clone 172058 5'), and N24601 (yx72e01.s1 Homo sapiens cDNA clone 267288 3'). The predicted amino acid sequence disclosed herein for bf169\_13 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bf169\_13 protein demonstrated at least some  
30 similarity to sequences identified as L41834 (plant nuclear protein [Ensis minor]) and Z75539 (F28C1.1 [Caenorhabditis elegans]). Analysis of motifs in the predicted bf169\_13 protein revealed a "mitochondrial energy transfer proteins" signature at amino acid 574 of SEQ ID NO:94. Based upon sequence similarity, bf169\_13 proteins and each similar

protein or peptide may share at least some activity. The nucleotide sequence of bf169\_13 indicates that it may contain one or more GCCCCA, GCCC, GGA and/or GC repeat sequences.

bf169\_13 protein was expressed in a COS cell expression system, and an  
5 expressed protein band of approximately 109 kDa was detected in membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "bl152\_12"

A polynucleotide of the present invention has been identified as clone "bl152\_12".  
10 bl152\_12 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bl152\_12 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein  
15 as "bl152\_12 protein").

The nucleotide sequence of bl152\_12 as presently determined is reported in SEQ ID NO:95, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bl152\_12 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:96.

20 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone bl152\_12 should be approximately 1100 bp.

The nucleotide sequence disclosed herein for bl152\_12 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bl152\_12 demonstrated at least some similarity with sequences  
25 identified as AA280876 (zs97d04.s1 NCI\_CGAP\_GCB1 Soares NbHTGBC Homo sapiens cDNA clone 711559 3' similar to contains element MER22 repetitive element), AA280956 (zs97d04.r1 NCI\_CGAP\_GCB1 Soares NbHTGBC Homo sapiens cDNA clone 711559 5'), R21512 (yh19b03.s1 Homo sapiens cDNA clone 130157 3'), R67018 (yi26e05.s1 Homo sapiens cDNA clone 140384 3' similar to contains MER22 repetitive element),  
30 R71877 (yj87d11.s1 Homo sapiens cDNA clone 155733 3' similar to contains MER22 repetitive element), T22941 (Human gene signature HUMGS04666), W46539 (zc30g03.s1 Soares senescent fibroblasts NbHSF Homo sapiens cDNA clone 323860 3', mRNA sequence), and W70065 (zd49c04.s1 Soares fetal heart NbHH19W Homo sapiens cDNA

clone). The predicted amino acid sequence disclosed herein for bl152\_12 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bl152\_12 protein demonstrated at least some similarity to the sequence identified as Z82256 (B0513.2 [Caenorhabditis elegans]). Based upon  
5 sequence similarity, bl152\_12 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of bl152\_12 indicates that it may contain one or more GCC repeat sequences.

bl152\_12 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 25 kDa was detected in conditioned medium using SDS  
10 polyacrylamide gel electrophoresis.

#### Clone "bz578\_1"

15 A polynucleotide of the present invention has been identified as clone "bz578\_1". bz578\_1 was isolated from a human fetal kidney cDNA library using methods and was identified as encoding a novel protein on the basis of computer analysis of the amino acid sequence of the encoded protein. bz578\_1 is a full-length clone, including the entire coding sequence of a novel protein (also referred to herein as "bz578\_1 protein").

20 The nucleotide sequence of bz578\_1 as presently determined is reported in SEQ ID NO:97, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the bz578\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:98.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
25 bz578\_1 should be approximately 1000 bp.

The nucleotide sequence disclosed herein for bz578\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. bz578\_1 demonstrated at least some similarity with sequences identified as T47038 (yb12e08.r1 Homo sapiens cDNA clone 70982 5' contains L1  
30 repetitive element) and Z82975 (Human DNA sequence from PAC 36J3, between markers DXS1192 and DXS102 on chromosome X). The predicted amino acid sequence disclosed herein for bz578\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted bz578\_1 protein

demonstrated at least some similarity to sequences identified as AF051782 (diaphanous 1 [Homo sapiens]), U96963 (diaphanous 1 [mouse]), and U93572 (putative p150 [Homo sapiens]). Based upon sequence similarity, bz578\_1 proteins and each similar protein or peptide may share at least some activity. The nucleotide sequence of bz578\_1 indicates  
5 that it may contain one or more L1 repeat sequences.

Clone "cb123\_1"

A polynucleotide of the present invention has been identified as clone "cb123\_1". cb123\_1 was isolated from a human fetal brain cDNA library using methods which are  
10 selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cb123\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cb123\_1 protein").

15 The nucleotide sequence of cb123\_1 as presently determined is reported in SEQ ID NO:99, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cb123\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:100. Amino acids 44 to 56 of SEQ ID NO:100 are a predicted leader/signal sequence, with the  
20 predicted mature amino acid sequence beginning at amino acid 57. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the cb123\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
25 cb123\_1 should be approximately 1500 bp.

The nucleotide sequence disclosed herein for cb123\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cb123\_1 demonstrated at least some similarity with sequences identified as AA309020 (EST179803 Colon carcinoma (Caco-2) cell line I Homo sapiens  
30 cDNA 5' end, mRNA sequence), R89617 (ym98b08.s1 Homo sapiens cDNA clone 166935 3'), T16814 (NIB1893 Normalized infant brain, Bento Soares Homo sapiens cDNA 3' end similar to EST02882 H. sapiens cDNA clone HFBCL71), T24092 (Human gene signature HUMGS06080), and T55187 (yb43e06.s1 Homo sapiens cDNA clone 73954 3'). The

predicted amino acid sequence disclosed herein for cb123\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cb123\_1 protein demonstrated at least some similarity to the sequence identified as U33331 (orf UL133 [Human cytomegalovirus]). Based upon sequence  
5 similarity, cb123\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts two additional potential transmembrane domains within the cb123\_1 protein sequence, one centered around amino acid 15 and another around amino acid 80 of SEQ ID NO:100.

10 Clone "ch245\_1"

A polynucleotide of the present invention has been identified as clone "ch245\_1". ch245\_1 was isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
15 analysis of the amino acid sequence of the encoded protein. ch245\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ch245\_1 protein").

The nucleotide sequence of ch245\_1 as presently determined is reported in SEQ ID NO:101, and includes a poly(A) tail. What applicants presently believe to be the proper  
20 reading frame and the predicted amino acid sequence of the ch245\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:102. The TopPredII computer program predicts a potential transmembrane domain within the ch245\_1 protein sequence centered around amino acid 87 of SEQ ID NO:102.

Another potential ch245\_1 reading frame and predicted amino acid sequence is  
25 encoded by basepairs 533 to 778 of SEQ ID NO:101 and is reported in SEQ ID NO:180. The TopPredII computer program predicts a potential transmembrane domain within the SEQ ID NO:180 amino acid sequence centered around amino acid 34 of SEQ ID NO:180.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ch245\_1 should be approximately 1350 bp.

30 The nucleotide sequence disclosed herein for ch245\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ch245\_1 demonstrated at least some similarity with sequences identified as AA402307 (zu48f03.r1 Soares ovary tumor NbHOT Homo sapiens cDNA



clone 741245 5', mRNA sequence), H19032 (ym44e04.r1 Homo sapiens cDNA clone 50921 5'), H19323 (ym44e04.s1 Homo sapiens cDNA clone 50921 3'), and N36070 (yy02g11.r1 Homo sapiens cDNA clone 270116 5'). The predicted amino acid sequence disclosed herein for ch245\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted ch245\_1 protein demonstrated at least some similarity to the sequence identified as M58597 (ELAM-1 ligand fucosyltransferase [Homo sapiens]) and U36763 (fatty acid synthase [Mycobacterium bovis]). Based upon sequence similarity, ch245\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "cj378\_3"

A polynucleotide of the present invention has been identified as clone "cj378\_3". cj378\_3 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. cj378\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cj378\_3 protein").

The nucleotide sequence of cj378\_3 as presently determined is reported in SEQ ID NO:103, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the cj378\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:104.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cj378\_3 should be approximately 1400 bp.

The nucleotide sequence disclosed herein for cj378\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cj378\_3 demonstrated at least some similarity with sequences identified as D60138 (Human fetal brain cDNA 5'-end GEN-088A04, mRNA sequence), H19318 (ym44d06.s1 Homo sapiens cDNA clone 51231 3'), H41859 (yo07g06.r1 Homo sapiens cDNA clone 177274 5'), T25386 (Human gene signature HUMGS07551), and T75383 (yc89g05.r1 Homo sapiens cDNA clone 23351 5'). Based upon sequence

similarity, cj378\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain at the N-terminus of the the cj378\_3 protein sequence (SEQ ID NO:104).

5           Clone "cw1481\_1"

A polynucleotide of the present invention has been identified as clone "cw1481\_1". cw1481\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer  
10 analysis of the amino acid sequence of the encoded protein. cw1481\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "cw1481\_1 protein").

The nucleotide sequence of cw1481\_1 as presently determined is reported in SEQ ID NO:105, and includes a poly(A) tail. What applicants presently believe to be the proper  
15 reading frame and the predicted amino acid sequence of the cw1481\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:106.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone cw1481\_1 should be approximately 2380 bp.

The nucleotide sequence disclosed herein for cw1481\_1 was searched against the  
20 GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. cw1481\_1 demonstrated at least some similarity with sequences identified as AA027927 (zk05a10.r1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 469626 5'), AA027928 (zk05a10.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 469626 3' similar to contains MER28.b2 MER28 repetitive element),  
25 AA113357 (zn69g06.s1 Stratagene HeLa cell s3 937216 Homo sapiens cDNA clone 563482 3'), AA252304 (zs12b08.s1 Soares NbHTGBC Homo sapiens cDNA clone 684951 3' similar to contains element MER22 repetitive element), AA976744 (oq09a09.s1 NCI\_CGAP\_GC4 Homo sapiens cDNA clone IMAGE 1585816 3' similar to TR O15025 O15025 KIAA0308 ;contains element MER22 repetitive element; mRNA sequence),  
30 R55084 (yg87a06.r1 Homo sapiens cDNA clone 40244 5'), U00930 (Human clone C4E 1.63 (CAC)<sub>n</sub>/(GTG)<sub>n</sub> repeat-containing mRNA), U00955 (Human clone CE29 8.1 (CAC)<sub>n</sub>/(GTG)<sub>n</sub> repeat-containing mRNA), and W16808 (zb93a09.s1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 320344 3'). The predicted amino acid sequence

disclosed herein for cw1481\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted cw1481\_1 protein demonstrated at least some similarity to sequences identified as AB002306 (KIAA0308 [Homo sapiens]), X15906 (precursor polypeptide), and Z68751 (F01G4.1 [Caenorhabditis elegans]). Based upon sequence similarity, cw1481\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the cw1481\_1 protein sequence centered around amino acid 431 of SEQ ID NO:106, and a putative transmembrane domain within the cw1481\_1 protein sequence centered around amino acid 395 of SEQ ID NO:106. The amino acid sequence of cw1481\_1 indicates that it has a histidine-rich region and a serine-rich region, and it is strongly internally repeated.

#### Clone "dd119\_4"

A polynucleotide of the present invention has been identified as clone "dd119\_4". dd119\_4 was isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. dd119\_4 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "dd119\_4 protein").

The nucleotide sequence of dd119\_4 as presently determined is reported in SEQ ID NO:107, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the dd119\_4 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:108. Amino acids 27 to 39 of SEQ ID NO:108 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 40. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the dd119\_4 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone dd119\_4 should be approximately 3350 bp.

The nucleotide sequence disclosed herein for dd119\_4 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and

FASTA search protocols. dd119\_4 demonstrated at least some similarity with sequences identified as AA151924 (zo30e05.r1 Stratagene colon (#937204) Homo sapiens cDNA clone 588416 5' similar to SW SLIT\_DROME P24014 SLIT PROTEIN PRECURSOR; mRNA sequence), AA193464 (zr41c06.s1 Soares NhHMPu S1 Homo sapiens cDNA clone 665962 3'), AB011135 (Homo sapiens mRNA for KIAA0563 protein, complete cds), G23888 (human STS WI-12393), H04996 (yl74c12.s1 Homo sapiens cDNA clone 43851 3'), M86526 (Rat proline-rich protein (PRP) gene, 5' end, and containing several Alu-like repetitive elements), M86514 (Rat proline-rich protein mRNA, 3' end), W68823 (zd37f04.r1 Soares fetal heart NbHH19W Homo sapiens cDNA clone 342847 5'), and Z54386 (H.sapiens CpG island DNA genomic MseI fragment, clone 10g3, forward read cpg10g3.ft1a). The predicted amino acid sequence disclosed herein for dd119\_4 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted dd119\_4 protein demonstrated at least some similarity to sequences identified as AB011135 (KIAA0563 protein [Homo sapiens]) and M86526 (proline-rich protein [Rattus norvegicus]). The rat proline-rich protein (PRP) is encoded by a single-copy gene and is expressed in the ventral prostate of the rat, with the precursor protein product being cleaved into multiple proline-rich polypeptides. Based upon sequence similarity, dd119\_4 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the dd119\_4 protein sequence centered around amino acid 928 of SEQ ID NO:108.

dd119\_4 protein was expressed in a COS cell expression system, and an expressed protein band of approximately 166 kDa was detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Clone "df202\_3"

A polynucleotide of the present invention has been identified as clone "df202\_3". df202\_3 was isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. df202\_3 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "df202\_3 protein").

The nucleotide sequence of df202\_3 as presently determined is reported in SEQ ID NO:109, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the df202\_3 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:110.

5 Amino acids 17 to 29 of SEQ ID NO:110 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 30. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the df202\_3 protein.

10 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone df202\_3 should be approximately 1600 bp.

The nucleotide sequence disclosed herein for df202\_3 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. df202\_3 demonstrated at least some similarity with sequences  
15 identified as AA138679 (mq76g03.r1 Stratagene mouse melanoma (#937312) Mus musculus cDNA clone 584692 5'), AA283121 (zt17b05.s1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 713361 3'), AA286996 (zs58c10.r1 NCI\_CGAP\_GCB1 Soares NbHTGBC Homo sapiens cDNA clone IMAGE 701682 5'), N54968 (yv38g01.s1 Homo sapiens cDNA clone 245040 3'), T20071 (Human gene signature HUMGS01213), and  
20 W28275 (44g12 Human retina cDNA randomly primed sublibrary Homo sapiens cDNA).

The predicted amino acid sequence disclosed herein for df202\_3 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted df202\_3 protein demonstrated at least some similarity to the sequence identified as Z81137 (W02D9.h [Caenorhabditis elegans]). Based upon sequence  
25 similarity, df202\_3 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts three additional potential transmembrane domains within the df202\_3 protein sequence, centered around amino acids 55, 80, and 108 of SEQ ID NO:110, respectively.

30 Clone "km225\_1"

A polynucleotide of the present invention has been identified as clone "km225\_1". km225\_1 was isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was

identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. km225\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "km225\_1 protein").

5       The nucleotide sequence of km225\_1 as presently determined is reported in SEQ ID NO:111, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the km225\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:112. Amino acids 9 to 21 of SEQ ID NO:112 are a predicted leader/signal sequence, with the  
10       predicted mature amino acid sequence beginning at amino acid 22. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the km225\_1 protein.

      The EcoRI/NotI restriction fragment obtainable from the deposit containing clone  
15       km225\_1 should be approximately 2300 bp.

      The nucleotide sequence disclosed herein for km225\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. km225\_1 demonstrated at least some similarity with sequences identified as AA101603 (zk94h09.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA  
20       clone 490529 3' similar to contains Alu repetitive element; mRNA sequence). Based upon sequence similarity, km225\_1 proteins and each similar protein or peptide may share at least some activity.

#### Clone "mj301\_1"

25       A polynucleotide of the present invention has been identified as clone "mj301\_1". mj301\_1 was isolated from a human adult lymph node cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. mj301\_1 is a full-length clone,  
30       including the entire coding sequence of a secreted protein (also referred to herein as "mj301\_1 protein").

      The nucleotide sequence of mj301\_1 as presently determined is reported in SEQ ID NO:113, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the mj301\_1 protein

corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:114. Amino acids 65 to 77 of SEQ ID NO:114 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 78. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the mj301\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone mj301\_1 should be approximately 2760 bp; however, a band of 550 bp has been detected in restriction digests, possibly due to an internal EcoRI or NotI restriction site in the clone.

The nucleotide sequence disclosed herein for mj301\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. mj301\_1 demonstrated at least some similarity with sequences identified as AA053085 (zl73d01.s1 Stratagene colon (#937204) Homo sapiens cDNA clone 510241 3'), AA347293 (EST53566 Fetal heart II Homo sapiens cDNA 5' end), AA813287 (ai76a07.s1 Soares testis NHT Homo sapiens cDNA clone 1376724 3', mRNA sequence), R45713 (Ha117-f Homo sapiens cDNA clone a117-f), T20114 (Human gene signature HUMGS01258), U46278 (Human clone xs252 mRNA sequence), Z36823 (H.sapiens (xs170) mRNA), and Z36832 (H.sapiens (xs170) mRNA). The human xs170 sequence is differentially expressed in pancreatic cancer cells. The predicted amino acid sequence disclosed herein for mj301\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted mj301\_1 protein demonstrated at least some similarity to the sequence identified as U07818 (putative phospho-beta-glucosidase [Bacillus stearothermophilus]). Based upon sequence similarity, mj301\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts an additional potential transmembrane domain within the mj301\_1 protein sequence centered around amino acid 60 of SEQ ID NO:114.

#### Clone "ml10\_7"

A polynucleotide of the present invention has been identified as clone "ml10\_7". ml10\_7 was isolated from a human adult brain (caudate nucleus) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis

of computer analysis of the amino acid sequence of the encoded protein. ml10\_7 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "ml10\_7 protein").

5 The nucleotide sequence of ml10\_7 as presently determined is reported in SEQ ID NO:115, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the ml10\_7 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:116. Amino acids 30 to 42 of SEQ ID NO:116 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 43. Due to the  
10 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the ml10\_7 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone ml10\_7 should be approximately 1600 bp.

15 The nucleotide sequence disclosed herein for ml10\_7 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. ml10\_7 demonstrated at least some similarity with sequences identified as AA411457 (zv30f06.s1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 755171 3'), AA411585 (zv30f06.r1 Soares ovary tumor NbHOT Homo sapiens  
20 cDNA clone 755171 5', mRNA sequence), AA485512 (zx90b02.r1 Soares ovary tumor NbHOT Homo sapiens cDNA clone 810987 5'), R97588 (yq59b05.r1 Homo sapiens cDNA clone 200049 5' similar to contains MSR1 repetitive element), and T23020 (Human gene signature HUMGS04748). The predicted amino acid sequence disclosed herein for ml10\_7 was searched against the GenPept and GeneSeq amino acid sequence databases using the  
25 BLASTX search protocol. The predicted ml10\_7 protein demonstrated at least some similarity to the sequence identified as R56978 (Human myotonic dystrophy gene protein). Based upon sequence similarity, ml10\_7 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four additional potential transmembrane domains within the ml10\_7 protein sequence, centered  
30 approximately around amino acids 20, 55 (between residues 50 and 60), 85 (between residues 80 and 89), and 175 (between residues 169 and 180) of SEQ ID NO:116, respectively. ml10\_7 appears to represent one member of a group of multiple alternatively spliced transcripts.



Clone "my340\_1"

A polynucleotide of the present invention has been identified as clone "my340\_1". my340\_1 was isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was  
5 identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. my340\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "my340\_1 protein").

The nucleotide sequence of my340\_1 as presently determined is reported in SEQ  
10 ID NO:117, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the my340\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:118.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone my340\_1 should be approximately 1800 bp.

15 The nucleotide sequence disclosed herein for my340\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. my340\_1 demonstrated at least some similarity with sequences identified as AA469015 (nc79g10.r1 NCI\_CGAP\_Pr2 Homo sapiens cDNA clone IMAGE:783618), H85290 (yv86f01.r1 Homo sapiens cDNA clone 249625 5'), L29074  
20 (Homo sapiens fragile X mental retardation protein (FMR-1) gene (6 alternative splices), complete cds), M86699 (Human kinase (TTK) mRNA, complete cds), W19755 (zb38f08.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 305895 5'), W63667 (zc57h10.r1 Soares parathyroid tumor NbHPA Homo sapiens cDNA clone 326467 5', mRNA sequence), and Z84478 (Human DNA sequence). The predicted amino  
25 acid sequence disclosed herein for my340\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted my340\_1 protein demonstrated at least some similarity to the sequence identified as M86699 (kinase [Homo sapiens]). The human TTK kinase can phosphorylate serine, threonine, and tyrosine hydroxyamino acids. Based upon sequence similarity,  
30 my340\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts a potential transmembrane domain within the my340\_1 protein sequence centered around amino acid 50 of SEQ ID NO:28.

Deposit of Clones

Clones bn365\_53, bo342\_2, dn721\_8, dn834\_1, pd278\_5, pe80\_1, pm113\_1, pm749\_8, pt31\_4, and pv296\_5 were deposited on May 7, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.)  
5 as an original deposit under the Budapest Treaty and were given the accession number ATCC 98752, from which each clone comprising a particular polynucleotide is obtainable.

Clones er311\_20, fh149\_12, pc201\_6, pl87\_1, and pm514\_4 were deposited on June 2, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty  
10 and were given the accession number ATCC 98781, from which each clone comprising a particular polynucleotide is obtainable.

Clones co155\_12, fn189\_13, lv2\_47, ml243\_1, pm96\_9, pu261\_1, pw214\_15, qb56\_19, qc646\_1, qf116\_2, and qf662\_3 were deposited on July 2, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession  
15 number ATCC 98808, from which each clone comprising a particular polynucleotide is obtainable.

Clones am748\_5, cj507\_1, cn922\_5, cw691\_11, cw1000\_2, cw1640\_1, d24\_1, dd426\_1, and di393\_2 were deposited on July 16, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 98817, from which each clone comprising a particular polynucleotide is obtainable.  
20

Clones dj167\_2, dw665\_4, dx146\_12, dx219\_13, fm3\_1, h225\_1, kj320\_1, ml236\_5, and pu282\_10, were deposited on July 16, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 98818, from which each clone comprising a particular polynucleotide is obtainable.  
25

Clones at94\_2, bf169\_13, bl152\_12, bz578\_1, cb123\_1, ch245\_1, cj378\_3, cw1481\_1, dd119\_4, df202\_3, km225\_1, mj301\_1, ml10\_7, and my340\_1 were deposited on July 22, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 98822, from which each clone comprising a particular polynucleotide is obtainable.  
30

Clone dj167\_19 was deposited on February 5, 1999 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and was given the accession number ATCC 207090, from which the dj167\_19 clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R. § 1.806.

Each clone has been transfected into separate bacterial cells (*E. coli*) in the composite deposits above. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or pNOTs vector depicted in Figures 1A and 1B, respectively. The pED6dpc2 vector ("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and insertion of the M13 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences provided herein, or from a combination of those sequences. The sequence of an oligonucleotide probe that was used to isolate or to sequence each full-length clone is identified below, and should be most reliable in isolating the clone of interest.

<u>Clone</u>	<u>Probe Sequence</u>
bn365_53	SEQ ID NO:119

	bo342_2	SEQ ID NO:120
	dn721_8	SEQ ID NO:121
	dn834_1	SEQ ID NO:122
	pd278_5	SEQ ID NO:123
5	pe80_1	SEQ ID NO:124
	pm113_1	SEQ ID NO:125
	pm749_8	SEQ ID NO:126
	pt31_4	SEQ ID NO:127
	pv296_5	SEQ ID NO:128
10	er311_20	SEQ ID NO:129
	fh149_12	SEQ ID NO:130
	pc201_6	SEQ ID NO:131
	pl87_1	SEQ ID NO:132
	pm514_4	SEQ ID NO:133
15	co155_12	SEQ ID NO:134
	fn189_13	SEQ ID NO:135
	lv2_47	SEQ ID NO:136
	ml243_1	SEQ ID NO:137
	pm96_9	SEQ ID NO:138
20	pu261_1	SEQ ID NO:139
	pw214_15	SEQ ID NO:140
	qb56_19	SEQ ID NO:141
	qc646_1	SEQ ID NO:142
	qf116_2	SEQ ID NO:143
25	qf662_3	SEQ ID NO:144
	am748_5	SEQ ID NO:145
	cj507_1	SEQ ID NO:146
	cn922_5	SEQ ID NO:147
	cw691_11	SEQ ID NO:148
30	cw1000_2	SEQ ID NO:149
	cw1640_1	SEQ ID NO:150
	d24_1	SEQ ID NO:151
	dd426_1	SEQ ID NO:152
	di393_2	SEQ ID NO:153

	dj167_2	SEQ ID NO:154
	dw665_4	SEQ ID NO:155
	dx146_12	SEQ ID NO:156
	dx219_13	SEQ ID NO:157
5	fm3_1	SEQ ID NO:158
	h225_1	SEQ ID NO:159
	kj320_1	SEQ ID NO:160
	ml236_5	SEQ ID NO:161
	pu282_10	SEQ ID NO:162
10	at94_2	SEQ ID NO:163
	bf169_13	SEQ ID NO:164
	bl152_12	SEQ ID NO:165
	bz578_1	SEQ ID NO:166
	cb123_1	SEQ ID NO:167
15	ch245_1	SEQ ID NO:168
	cj378_3	SEQ ID NO:169
	cw1481_1	SEQ ID NO:170
	dd119_4	SEQ ID NO:171
	df202_3	SEQ ID NO:172
20	km225_1	SEQ ID NO:173
	mj301_1	SEQ ID NO:174
	ml10_7	SEQ ID NO:175
	my340_1	SEQ ID NO:176

25 In the sequences listed above which include an N at position 2, that position is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

30 The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;

- (b) It should be designed to have a  $T_m$  of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).

The oligonucleotide should preferably be labeled with  $\gamma$ -<sup>32</sup>P ATP (specific activity 6000 Ci/mmol) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a scintillation counter. Preferably, specific activity of the resulting probe should be approximately 4e+6 dpm/pmol.

The bacterial culture containing the pool of full-length clones should preferably be thawed and 100  $\mu$ l of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100  $\mu$ g/ml. The culture should preferably be grown to saturation at 37°C, and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100  $\mu$ g/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at 37°C. Other known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at 65°C for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100  $\mu$ g/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at a concentration greater than or equal to 1e+6 dpm/mL. The filter is then preferably incubated at 65°C with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at 65°C for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

5 Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as  
10 immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a  
15 decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein may be obtained by expression of the disclosed full-length  
20 polynucleotide (preferably those deposited with ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that  
25 are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can  
30 be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that

has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

The chromosomal location corresponding to the polynucleotide sequences disclosed herein may also be determined, for example by hybridizing appropriately  
5 labeled polynucleotides of the present invention to chromosomes *in situ*. It may also be possible to determine the corresponding chromosomal location for a disclosed polynucleotide by identifying significantly similar nucleotide sequences in public  
10 databases, such as expressed sequence tags (ESTs), that have already been mapped to particular chromosomal locations. For at least some of the polynucleotide sequences disclosed herein, public database sequences having at least some similarity to the  
15 polynucleotide of the present invention have been listed by database accession number. Searches using the GenBank accession numbers of these public database sequences can then be performed at an Internet site provided by the National Center for Biotechnology Information having the address <http://www.ncbi.nlm.nih.gov/UniGene/>, in order to  
20 identify "UniGene clusters" of overlapping sequences. Many of the "UniGene clusters" so identified will already have been mapped to particular chromosomal sites.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The  
25 desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, *Trends Pharmacol. Sci.* 15(7): 250-254; Lavarosky *et al.*, 1997, *Biochem. Mol. Med.* 62(1): 11-22; and Hampel, 1998, *Prog. Nucleic Acid Res. Mol. Biol.* 58: 1-39; all of which are incorporated by reference herein). Transgenic animals that have  
30 multiple copies of the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein).  
In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of



transposable elements (Plasterk, 1992, *Bioessays* **14**(9): 629-633; Zwaal *et al.*, 1993, *Proc. Natl. Acad. Sci. USA* **90**(16): 7431-7435; Clark *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* **91**(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination, preferably detected by positive/negative genetic selection strategies (Mansour *et al.*, 1988, *Nature* **336**: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of transmembrane domains in an amino acid sequence, domains which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST

version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* **266**: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of Molecular Biology* **215**: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* **3**: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* **90**: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid comparison matrices such as PAM can be utilized.

Species homologues of the disclosed polynucleotides and proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the given protein or

polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuáñez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotides which also encode proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from individuals of the appropriate species.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

5 The present invention also includes polynucleotides that hybridize under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) <sup>‡</sup>	Hybridization Temperature and Buffer <sup>†</sup>	Wash Temperature and Buffer <sup>†</sup>
5	A	≥ 50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
	B	<50	T <sub>B</sub> *; 1xSSC	T <sub>B</sub> *; 1xSSC
	C	≥ 50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC
	D	<50	T <sub>D</sub> *; 1xSSC	T <sub>D</sub> *; 1xSSC
	E	≥ 50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
	F	<50	T <sub>F</sub> *; 1xSSC	T <sub>F</sub> *; 1xSSC
10	G	≥ 50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
	H	<50	T <sub>H</sub> *; 4xSSC	T <sub>H</sub> *; 4xSSC
	I	≥ 50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
	J	<50	T <sub>J</sub> *; 4xSSC	T <sub>J</sub> *; 4xSSC
	K	≥ 50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
	L	<50	T <sub>L</sub> *; 2xSSC	T <sub>L</sub> *; 2xSSC
15	M	≥ 50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
	N	<50	T <sub>N</sub> *; 6xSSC	T <sub>N</sub> *; 6xSSC
	O	≥ 50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
	P	<50	T <sub>P</sub> *; 6xSSC	T <sub>P</sub> *; 6xSSC
	Q	≥ 50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
	R	<50	T <sub>R</sub> *; 4xSSC	T <sub>R</sub> *; 4xSSC

<sup>‡</sup>: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

<sup>†</sup>: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH<sub>2</sub>PO<sub>4</sub>, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

\*T<sub>B</sub> - T<sub>R</sub>: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T<sub>m</sub>) of the hybrid, where T<sub>m</sub> is determined according to the following equations. For hybrids less than 18 base pairs in length, T<sub>m</sub>(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T<sub>m</sub>(°C) = 81.5 + 16.6(log<sub>10</sub>[Na<sup>+</sup>]) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and [Na<sup>+</sup>] is the concentration of sodium ions in the hybridization buffer ([Na<sup>+</sup>] for 1xSSC = 0.165 M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds.,  
5 John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or  
10 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

The isolated polynucleotide of the invention may be operably linked to an  
15 expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably  
20 linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the  
25 protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

30 Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial

strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from the Eastman Kodak Company (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art



given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

### USES AND BIOLOGICAL ACTIVITY

5           The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies  
10 or vectors suitable for introduction of DNA).

#### Research Uses and Utilities

          The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express  
15 recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare  
20 with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for  
25 examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, those  
30 described in Gyuris *et al.*, 1993, *Cell* 75: 791-803 and in Rossi *et al.*, 1997, *Proc. Natl. Acad. Sci. USA* 94: 8405-8410, all of which are incorporated by reference herein) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

#### Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may

induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: *Polyclonal T cell stimulation*, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and *Measurement of mouse and human Interferon  $\gamma$* , Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: *Measurement of Human and Murine Interleukin 2 and Interleukin 4*, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., *J. Exp. Med.* 173:1205-1211, 1991; Moreau et al., *Nature* 336:690-692, 1988; Greenberger et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983; *Measurement of mouse and human interleukin 6* - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; *Measurement of human Interleukin 11* - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991;

Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

#### Immune Stimulating or Suppressing Activity

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, *Leishmania* spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease.

Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), *e.g.*, preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term

tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

5       The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as  
10 described in Lenschow *et al.*, Science 257:789-792 (1992) and Turka *et al.*, Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

15       Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms.  
20 Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from  
25 the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosis in MRL/*lpr/lpr* mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and  
30 murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune

response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain protein and  $\beta_2$

microglobulin protein or an MHC class II  $\alpha$  chain protein and an MHC class II  $\beta$  chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (*e.g.*, B7-1, B7-2, B7-3) induces a T cell mediated  
5 immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated  
10 immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without  
15 limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al.,  
20 J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnoli et al.,  
25 Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro*  
30 antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek,



D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

- 5 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995;
- 10 Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

- Assays for lymphocyte survival/apoptosis (which will identify, among others,
- 15 proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993;
- 20 Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

- Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

25

#### Hematopoiesis Regulating Activity

- A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell
- 30 lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid

cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and

Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, *et al.* eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, *et al.* eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

5

#### Tissue Growth Activity

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns,  
10 incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as  
15 well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal  
20 disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue  
25 destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in  
30 circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and

in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation

of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues,  
5 and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured  
10 by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium ).

15 Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

#### 20 Activin/Inhibin Activity

A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present  
25 invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin-  
30  $\beta$  group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

#### Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. *J. Clin. Invest.* 95:1370-1376, 1995; Lind et al.

APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

#### Hemostatic and Thrombolytic Activity

5 A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting  
10 formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

15 Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### 20 Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands,  
25 receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant  
30 receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160, 1989; Stoltenberg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

#### 10      Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

25

#### Cadherin/Tumor Invasion Suppressor Activity

Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus vulgaris and pemphigus foliaceus (auto-immune blistering skin diseases), Crohn's disease, and some developmental abnormalities.

30

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved



extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The cadherin domains bind calcium to form their tertiary structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this recognition site can change the specificity of a cadherin so that instead of recognizing only  
5 itself, the mutant molecule can now also bind to a different cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells  
10 become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal, restoring cells' adhesiveness to each other and to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas  
15 to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed  
20 in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different tissue in the body. Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the  
25 inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

Additionally, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block  
30 the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.

Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and polynucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable effects. Additionally, fragments of proteins of the present invention with cadherin activity, preferably truncated soluble cadherin fragments which have been found to be stable in the circulation of cancer patients, and polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al. Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

#### Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

#### Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s);

effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

### ADMINISTRATION AND DOSING

A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical

compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

5 The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to  
10 present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the  
15 invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers  
20 in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein  
25 by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing,  
30 prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in  
5 combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If  
10 administered sequentially, the attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical  
15 composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is  
20 administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention.  
25 When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid  
30 form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present

invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu\text{g}$  to about 100 mg (preferably about 0.1ng to about 10 mg, more preferably about 0.1  $\mu\text{g}$  to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. As used herein, the term "antibody" includes without limitation a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a single-chain antibody, a CDR-grafted antibody, a humanized antibody, or fragments thereof which bind to the indicated protein.

Such term also includes any other species derived from an antibody or antibody sequence which is capable of binding the indicated protein.

Antibodies to a particular protein can be produced by methods well known to those skilled in the art. For example, monoclonal antibodies can be produced by generation of  
5 antibody-producing hybridomas in accordance with known methods (see for example, Goding, 1983, *Monoclonal antibodies: principles and practice*, Academic Press Inc., New York; and Yokoyama, 1992, "Production of Monoclonal Antibodies" in *Current Protocols in Immunology*, Unit 2.5, Greene Publishing Assoc. and John Wiley & Sons). Polyclonal sera and antibodies can be produced by inoculation of a mammalian subject with the  
10 relevant protein or fragments thereof in accordance with known methods. Fragments of antibodies, receptors, or other reactive peptides can be produced from the corresponding antibodies by cleavage of and collection of the desired fragments in accordance with known methods (see for example, Goding, *supra*; and Andrew et al., 1992, "Fragmentation of Immunoglobulins" in *Current Protocols in Immunology*, Unit 2.8, Greene Publishing  
15 Assoc. and John Wiley & Sons). Chimeric antibodies and single chain antibodies can also be produced in accordance with known recombinant methods (see for example, 5,169,939, 5,194,594, and 5,576,184). Humanized antibodies can also be made from corresponding murine antibodies in accordance with well known methods (see for example, U.S. Patent Nos. 5,530,101, 5,585,089, and 5,693,762). Additionally, human antibodies may be  
20 produced in non-human animals such as mice that have been genetically altered to express human antibody molecules (see for example Fishwild *et al.*, 1996, *Nature Biotechnology* **14**: 845-851; Mendez *et al.*, 1997, *Nature Genetics* **15**: 146-156 (erratum *Nature Genetics* **16**: 410); and U.S. Patents 5,877,397 and 5,625,126). Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide  
25 immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, *FEBS Lett.* 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful  
30 diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where

abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

5 For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably  
10 be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the  
15 methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical  
20 applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium  
25 sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other  
30 ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.



Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions  
5 from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of  
10 carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to  
15 provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in  
20 question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to  
25 humans, are desired patients for such treatment with proteins of the present invention.

The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of  
30 a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect

the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

5 Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

10 Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

Patent and literature references cited herein are incorporated by reference as if fully set forth.

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:1;
  - (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 61 to nucleotide 366;
  - (c) the nucleotide sequence of the full-length protein coding sequence of clone bn365\_53 deposited under accession number ATCC 98752;
  - (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;
  - (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
  - (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
  - (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
  - (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:1.
2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:

- (a) growing a culture of a host cell transformed with the polynucleotide of claim 2 in a suitable culture medium; and
  - (b) purifying said protein from the culture.
6. A protein produced according to the process of claim 5.
7. An isolated polynucleotide encoding the protein of claim 6.
8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752.
9. A protein comprising an amino acid sequence selected from the group consisting of:
  - (a) the amino acid sequence of SEQ ID NO:2;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone bn365\_53 deposited under accession number ATCC 98752;the protein being substantially free from other mammalian proteins.
10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.
11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.
12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:3;
  - (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 206 to nucleotide 1915;
  - (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 1358 to nucleotide 1915;

(d) the nucleotide sequence of the full-length protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;

(f) the nucleotide sequence of a mature protein coding sequence of clone bo342\_2 deposited under accession number ATCC 98752;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone bo342\_2 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:5;

(b) the nucleotide sequence of SEQ ID NO:5 from nucleotide 749 to nucleotide 2689;

(c) the nucleotide sequence of the full-length protein coding sequence of clone dn721\_8 deposited under accession number ATCC 98752;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone dn721\_8 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:7;

(b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 20 to nucleotide 484;

(c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 18 to nucleotide 892;

(d) the nucleotide sequence of the full-length protein coding sequence of clone dn834\_1 deposited under accession number ATCC 98752;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;

(f) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

(g) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e); and

(i) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e), and that has a length that is at least 25% of the length of SEQ ID NO:7.

17. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:8;

(b) a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8; and

(c) the amino acid sequence encoded by the cDNA insert of clone dn834\_1 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins.

18. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:9;

(b) the nucleotide sequence of SEQ ID NO:9 from nucleotide 803 to nucleotide 1420;

- (c) the nucleotide sequence of SEQ ID NO:9 from nucleotide 1022 to nucleotide 1420;
  - (d) the nucleotide sequence of the full-length protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752;
  - (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;
  - (f) the nucleotide sequence of a mature protein coding sequence of clone pd278\_5 deposited under accession number ATCC 98752;
  - (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;
  - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
  - (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
  - (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
  - (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:9.
19. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:10;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pd278\_5 deposited under accession number ATCC 98752;
- the protein being substantially free from other mammalian proteins.



20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 918 to nucleotide 1295;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pe80\_1 deposited under accession number ATCC 98752;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pe80\_1 deposited under accession number ATCC 98752;
- the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:13;

(b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 189 to nucleotide 428;

(c) the nucleotide sequence of SEQ ID NO:13 from nucleotide 348 to nucleotide 428;

(d) the nucleotide sequence of the full-length protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;

(f) the nucleotide sequence of a mature protein coding sequence of clone pm113\_1 deposited under accession number ATCC 98752;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:13.

23. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:14;

(b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and

(c) the amino acid sequence encoded by the cDNA insert of clone pm113\_1 deposited under accession number ATCC 98752;

the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:15;
- (b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 108 to nucleotide 1496;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pm749\_8 deposited under accession number ATCC 98752;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm749\_8 deposited under accession number ATCC 98752;
- the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:17;
- (b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 44 to nucleotide 2023;
- (c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 137 to nucleotide 2023;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pt31\_4 deposited under accession number ATCC 98752;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:18;
- (b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and

(c) the amino acid sequence encoded by the cDNA insert of clone pt31\_4 deposited under accession number ATCC 98752;  
the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:19;
- (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 24 to nucleotide 299;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pv296\_5 deposited under accession number ATCC 98752;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:19.

29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pv296\_5 deposited under accession number ATCC 98752;
- the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone er311\_20 deposited under accession number ATCC 98781;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone er311\_20 deposited under accession number ATCC 98781;
- the protein being substantially free from other mammalian proteins.

32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 484 to nucleotide 2043;
- (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 919 to nucleotide 2043;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;
- (f) the nucleotide sequence of a mature protein coding sequence of clone fh149\_12 deposited under accession number ATCC 98781;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:24;
- (b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and

(c) the amino acid sequence encoded by the cDNA insert of clone fh149\_12 deposited under accession number ATCC 98781; the protein being substantially free from other mammalian proteins.

34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:25;
- (b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 47 to nucleotide 1099;
- (c) the nucleotide sequence of SEQ ID NO:25 from nucleotide 143 to nucleotide 1099;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pc201\_6 deposited under accession number ATCC 98781;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:26;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group consisting of:



- (a) the amino acid sequence of SEQ ID NO:26;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pc201\_6 deposited under accession number ATCC 98781;
- the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:27;
- (b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 5 to nucleotide 259;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pl87\_1 deposited under accession number ATCC 98781;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
- (b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

(c) the amino acid sequence encoded by the cDNA insert of clone pl87\_1 deposited under accession number ATCC 98781; the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:29;
- (b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 62 to nucleotide 2284;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pm514\_4 deposited under accession number ATCC 98781;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:30;
- (b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and
- (c) the amino acid sequence encoded by the cDNA insert of clone pm514\_4 deposited under accession number ATCC 98781;

the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:31;
- (b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 36 to nucleotide 1997;
- (c) the nucleotide sequence of SEQ ID NO:31 from nucleotide 135 to nucleotide 1997;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone co155\_12 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:31.

41. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:32;

- (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone co155\_12 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 21 to nucleotide 1343;
- (c) the nucleotide sequence of SEQ ID NO:33 from nucleotide 84 to nucleotide 1343;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone fn189\_13 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
- (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and
- (c) the amino acid sequence encoded by the cDNA insert of clone fn189\_13 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:35;
- (b) the nucleotide sequence of SEQ ID NO:35 from nucleotide 66 to nucleotide 557;
- (c) the nucleotide sequence of SEQ ID NO:35 from nucleotide 235 to nucleotide 899;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone lv2\_47 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;
- (f) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (g) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e); and
- (i) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e), and that has a length that is at least 25% of the length of SEQ ID NO:35.

45. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
  - (b) the amino acid sequence of SEQ ID NO:36 from amino acid 58 to amino acid 164;
  - (c) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone lv2\_47 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:37;
- (b) the nucleotide sequence of SEQ ID NO:37 from nucleotide 104 to nucleotide 499;
- (c) the nucleotide sequence of SEQ ID NO:37 from nucleotide 215 to nucleotide 499;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ml243\_1 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:37.

47. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ml243\_1 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 2172 to nucleotide 2861;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pm96\_9 deposited under accession number ATCC 98808;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:40;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pm96\_9 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:41;
- (b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 43 to nucleotide 762;
- (c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 427 to nucleotide 762;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pu261\_1 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;



- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

51. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:42;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pu261\_1 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:43;
- (b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 579 to nucleotide 824;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone pw214\_15 deposited under accession number ATCC 98808;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:44;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone pw214\_15 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:45;

(b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 6 to nucleotide 383;

(c) the nucleotide sequence of the full-length protein coding sequence of clone qb56\_19 deposited under accession number ATCC 98808;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone qb56\_19 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 170 to nucleotide 1273;
- (c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 242 to nucleotide 1273;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone qc646\_1 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

57. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone qc646\_1 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 183 to nucleotide 1097;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone qf116\_2 deposited under accession number ATCC 98808;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
- (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
- (c) the amino acid sequence encoded by the cDNA insert of clone qf116\_2 deposited under accession number ATCC 98808;

the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 160 to nucleotide 741;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 595 to nucleotide 741;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;
- (f) the nucleotide sequence of a mature protein coding sequence of clone qf662\_3 deposited under accession number ATCC 98808;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:52;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone qf662\_3 deposited under accession number ATCC 98808;
- the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:53;
- (b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 924 to nucleotide 1196;
- (c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 1002 to nucleotide 1196;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone am748\_5 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.

63. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(c) the amino acid sequence encoded by the cDNA insert of clone am748\_5 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins.

64. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:55;

(b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 51 to nucleotide 1310;

(c) the nucleotide sequence of the full-length protein coding sequence of clone cj507\_1 deposited under accession number ATCC 98817;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:55.

65. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:56;

(b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and

(c) the amino acid sequence encoded by the cDNA insert of clone cj507\_1 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:57;

(b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 195 to nucleotide 1328;

(c) the nucleotide sequence of the full-length protein coding sequence of clone cn922\_5 deposited under accession number ATCC 98817;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;



(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:57.

67. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:58;

(b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and

(c) the amino acid sequence encoded by the cDNA insert of clone cn922\_5 deposited under accession number ATCC 98817;

the protein being substantially free from other mammalian proteins.

68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:59;

(b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 76 to nucleotide 942;

(c) the nucleotide sequence of the full-length protein coding sequence of clone cw691\_11 deposited under accession number ATCC 98817;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;

- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:59.

69. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw691\_11 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 11 to nucleotide 1252;
- (c) the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 1252;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone cw1000\_2 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;

- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:61.

71. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1000\_2 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:63;
- (b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 46 to nucleotide 1296;
- (c) the nucleotide sequence of SEQ ID NO:63 from nucleotide 451 to nucleotide 1296;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;

- (f) the nucleotide sequence of a mature protein coding sequence of clone cw1640\_1 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:64;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1640\_1 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827;
- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone d24\_1 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

75. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone d24\_1 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:67;

- (b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 149 to nucleotide 529;
- (c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 413 to nucleotide 529;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone dd426\_1 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dd426\_1 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 31 to nucleotide 543;
- (c) the nucleotide sequence of SEQ ID NO:69 from nucleotide 88 to nucleotide 543;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone di393\_2 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone di393\_2 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:69.

79. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:70;
- (b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone di393\_2 deposited under accession number ATCC 98817; the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:71;
- (b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 157 to nucleotide 1356;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone dj167\_2 deposited under accession number ATCC 98818;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:72;
- (b) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and
- (c) the amino acid sequence encoded by the cDNA insert of clone dj167\_2 deposited under accession number ATCC 98818; the protein being substantially free from other mammalian proteins.



82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:73;
- (b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 1383 to nucleotide 4490;
- (c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 1485 to nucleotide 4490;
- (d) the nucleotide sequence of SEQ ID NO:73 from nucleotide 3645 to nucleotide 4343;
- (e) the nucleotide sequence of the full-length protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090;
- (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;
- (g) the nucleotide sequence of a mature protein coding sequence of clone dj167\_19 deposited under accession number ATCC 207090;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;
- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;

- (b) the amino acid sequence of SEQ ID NO:74 from amino acid 637 to amino acid 1036;
  - (c) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone dj167\_19 deposited under accession number ATCC 207090;
- the protein being substantially free from other mammalian proteins.

84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 71 to nucleotide 1441;
- (c) the nucleotide sequence of SEQ ID NO:75 from nucleotide 152 to nucleotide 1441;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone dw665\_4 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:75.

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dw665\_4 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:77;
- (b) the nucleotide sequence of SEQ ID NO:77 from nucleotide 78 to nucleotide 1592;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone dx146\_12 deposited under accession number ATCC 98818;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:77.

87. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dx146\_12 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:79;
- (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 19 to nucleotide 948;
- (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 337 to nucleotide 948;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone dx219\_13 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:80;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dx219\_13 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

90. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:81;
- (b) the nucleotide sequence of SEQ ID NO:81 from nucleotide 5 to nucleotide 286;
- (c) the nucleotide sequence of SEQ ID NO:81 from nucleotide 62 to nucleotide 286;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone fm3\_1 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:82;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:81.

91. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:82;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:82, the fragment comprising eight contiguous amino acids of SEQ ID NO:82; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone fm3\_1 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

92. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:83;
- (b) the nucleotide sequence of SEQ ID NO:83 from nucleotide 141 to nucleotide 572;
- (c) the nucleotide sequence of SEQ ID NO:83 from nucleotide 333 to nucleotide 572;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone h225\_1 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:84;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:83.

93. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:84;

(b) a fragment of the amino acid sequence of SEQ ID NO:84, the fragment comprising eight contiguous amino acids of SEQ ID NO:84; and

(c) the amino acid sequence encoded by the cDNA insert of clone h225\_1 deposited under accession number ATCC 98818;

the protein being substantially free from other mammalian proteins.

94. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:85;

(b) the nucleotide sequence of SEQ ID NO:85 from nucleotide 391 to nucleotide 3210;

(c) the nucleotide sequence of SEQ ID NO:85 from nucleotide 505 to nucleotide 3210;

(d) the nucleotide sequence of the full-length protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;

(f) the nucleotide sequence of a mature protein coding sequence of clone kj320\_1 deposited under accession number ATCC 98818;

- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:86;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:85.

95. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:86;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:86, the fragment comprising eight contiguous amino acids of SEQ ID NO:86; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone kj320\_1 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

96. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:87;
- (b) the nucleotide sequence of SEQ ID NO:87 from nucleotide 42 to nucleotide 899;
- (c) the nucleotide sequence of SEQ ID NO:87 from nucleotide 522 to nucleotide 899;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818;



- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ml236\_5 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:87.

97. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising eight contiguous amino acids of SEQ ID NO:88; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ml236\_5 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

98. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:89;
- (b) the nucleotide sequence of SEQ ID NO:89 from nucleotide 6 to nucleotide 452;

- (c) the nucleotide sequence of SEQ ID NO:89 from nucleotide 399 to nucleotide 452;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- (f) the nucleotide sequence of a mature protein coding sequence of clone pu282\_10 deposited under accession number ATCC 98818;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:90;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:89.

99. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:90;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising eight contiguous amino acids of SEQ ID NO:90; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone pu282\_10 deposited under accession number ATCC 98818;
- the protein being substantially free from other mammalian proteins.

100. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:91;
- (b) the nucleotide sequence of SEQ ID NO:91 from nucleotide 4 to nucleotide 1179;
- (c) the nucleotide sequence of SEQ ID NO:91 from nucleotide 682 to nucleotide 1179;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone at94\_2 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:91.

101. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;
- (b) a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising eight contiguous amino acids of SEQ ID NO:92; and

(c) the amino acid sequence encoded by the cDNA insert of clone at94\_2 deposited under accession number ATCC 98822; the protein being substantially free from other mammalian proteins.

102. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:93;
- (b) the nucleotide sequence of SEQ ID NO:93 from nucleotide 56 to nucleotide 2077;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone bf169\_13 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:93.

103. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
- (b) a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising eight contiguous amino acids of SEQ ID NO:94; and
- (c) the amino acid sequence encoded by the cDNA insert of clone bf169\_13 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins.

104. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:95;
- (b) the nucleotide sequence of SEQ ID NO:95 from nucleotide 124 to nucleotide 735;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone bl152\_12 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:96;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:95.

105. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:96;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:96, the fragment comprising eight contiguous amino acids of SEQ ID NO:96; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone bl152\_12 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

106. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:97;
- (b) the nucleotide sequence of SEQ ID NO:97 from nucleotide 526 to nucleotide 816;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone bz578\_1 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:98;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:97.

107. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:98;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:98, the fragment comprising eight contiguous amino acids of SEQ ID NO:98; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone bz578\_1 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

108. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:99;

- (b) the nucleotide sequence of SEQ ID NO:99 from nucleotide 597 to nucleotide 992;
- (c) the nucleotide sequence of SEQ ID NO:99 from nucleotide 765 to nucleotide 992;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone cb123\_1 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:99.

109. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising eight contiguous amino acids of SEQ ID NO:100; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cb123\_1 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

110. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:101;
- (b) the nucleotide sequence of SEQ ID NO:101 from nucleotide 181 to nucleotide 480;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone ch245\_1 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:101.

111. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising eight contiguous amino acids of SEQ ID NO:102; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ch245\_1 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

112. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:103;



- (b) the nucleotide sequence of SEQ ID NO:103 from nucleotide 281 to nucleotide 541;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone cj378\_3 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:104;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:103.

113. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:104;
- (b) a fragment of the amino acid sequence of SEQ ID NO:104, the fragment comprising eight contiguous amino acids of SEQ ID NO:104; and
- (c) the amino acid sequence encoded by the cDNA insert of clone cj378\_3 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins.

114. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:105;
- (b) the nucleotide sequence of SEQ ID NO:105 from nucleotide 586 to nucleotide 2202;

- (c) the nucleotide sequence of SEQ ID NO:105 from nucleotide 401 to nucleotide 2349;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone cw1481\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;
- (f) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:106;
- (g) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106;
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e); and
- (i) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(e), and that has a length that is at least 25% of the length of SEQ ID NO:105.

115. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:106;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:106, the fragment comprising eight contiguous amino acids of SEQ ID NO:106; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone cw1481\_1 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

116. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:107;
- (b) the nucleotide sequence of SEQ ID NO:107 from nucleotide 29 to nucleotide 2905;

- (c) the nucleotide sequence of SEQ ID NO:107 from nucleotide 146 to nucleotide 2905;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone dd119\_4 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:108;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:107.

117. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:108;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising eight contiguous amino acids of SEQ ID NO:108; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone dd119\_4 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

118. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:109;
- (b) the nucleotide sequence of SEQ ID NO:109 from nucleotide 16 to nucleotide 369;
- (c) the nucleotide sequence of SEQ ID NO:109 from nucleotide 103 to nucleotide 369;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone df202\_3 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:109.

119. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;
- (b) a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising eight contiguous amino acids of SEQ ID NO:110; and

(c) the amino acid sequence encoded by the cDNA insert of clone df202\_3 deposited under accession number ATCC 98822; the protein being substantially free from other mammalian proteins.

120. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:111;
- (b) the nucleotide sequence of SEQ ID NO:111 from nucleotide 2192 to nucleotide 2539;
- (c) the nucleotide sequence of SEQ ID NO:111 from nucleotide 2255 to nucleotide 2539;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone km225\_1 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:112;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:111.

121. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:112;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:112, the fragment comprising eight contiguous amino acids of SEQ ID NO:112; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone km225\_1 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

122. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:113;
- (b) the nucleotide sequence of SEQ ID NO:113 from nucleotide 1734 to nucleotide 2030;
- (c) the nucleotide sequence of SEQ ID NO:113 from nucleotide 1965 to nucleotide 2030;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone mj301\_1 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:114;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:113.

123. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:114;
- (b) a fragment of the amino acid sequence of SEQ ID NO:114, the fragment comprising eight contiguous amino acids of SEQ ID NO:114; and
- (c) the amino acid sequence encoded by the cDNA insert of clone mj301\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins.

124. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:115;
- (b) the nucleotide sequence of SEQ ID NO:115 from nucleotide 799 to nucleotide 1350;
- (c) the nucleotide sequence of SEQ ID NO:115 from nucleotide 925 to nucleotide 1350;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ml10\_7 deposited under accession number ATCC 98822;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:115.

125. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:116;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising eight contiguous amino acids of SEQ ID NO:116; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone ml10\_7 deposited under accession number ATCC 98822;
- the protein being substantially free from other mammalian proteins.

126. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:117;
- (b) the nucleotide sequence of SEQ ID NO:117 from nucleotide 837 to nucleotide 1094;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone my340\_1 deposited under accession number ATCC 98822;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:118;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118;



(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:117.

127. A protein comprising an amino acid sequence selected from the group consisting of:


(a) the amino acid sequence of SEQ ID NO:118;

(b) a fragment of the amino acid sequence of SEQ ID NO:118, the fragment comprising eight contiguous amino acids of SEQ ID NO:118; and

(c) the amino acid sequence encoded by the cDNA insert of clone my340\_1 deposited under accession number ATCC 98822;

the protein being substantially free from other mammalian proteins.

FIGURE 1A




**Plasmid name:** pED6dpc2

**Plasmid size:** 5374 bp

**Comments/References:** pED6dpc2 is derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning. SST cDNAs are cloned between EcoRI and NotI. pED vectors are described in Kaufman et al.(1991), NAR 19: 4485-4490.

FIGURE 1B



**Plasmid name:** pNOTs

**Plasmid size:** 4529 bp

**Comments/References:** pNOTs is a derivative of pMT2 (Kaufman et al, 1989. Mol. Cell. Biol. 9:1741-1750). DHFR was deleted and a new polylinker was inserted between EcoRI and HpaI. M13 origin of replication was inserted in the Clal site. SST cDNAs are cloned between EcoRI and NotI.

## SEQUENCE LISTING

<110> Jacobs, Kenneth  
 McCoy, John M.  
 LaVallie, Edward R.  
 Collins-Racie, Lisa A.  
 Evans, Cheryl  
 Merberg, David  
 Treacy, Maurice  
 Agostino, Michael J.  
 Steininger II, Robert J.  
 Bowman, Michael R.  
 DiBlasio-Smith, Elizabeth  
 Widom, Angela  
 Genetics Institute, Inc.

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6069-74A

<140>  
 <141>

<160> 180

<170> PatentIn Ver. 2.0

<210> 1  
 <211> 571  
 <212> DNA  
 <213> Homo sapiens

<400> 1  
 ttcttcgccca ggctctctgc tgactcaagt tcttcagttc acgatcttct agttgcagcg 60  
 atgagtgcac gagtgagatc aagatccaga ggaagaggag atggtcagga ggctcccgat 120  
 gtggttgcat tcgtggctcc cgggtgaatct cagcaagagg aaccaccaac tgacaatcag 180  
 gatattgaac ctggacaaga gagagaagga acacctccga tcgaagaacg taaagtagaa 240  
 ggtgattgcc aggaaatgga tctggaaaag actcggagtg agcgtggaga tggctctgat 300  
 gtaaaagaga agactccacc taatcctaag catgctaaga ctaaagaagc aggagatggg 360  
 cagccataag ttaaaaagaa gacaagctga agctacacac atggctgatg tcacattgaa 420  
 aatgtgactg aaaatttgaa aattctctca ataaagtttg agttttctct gaaaaaaaaa 480  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 540  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a a 571

<210> 2  
 <211> 102  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met Ser Ala Arg Val Arg Ser Arg Ser Arg Gly Arg Gly Asp Gly Gln  
 1 5 10 15  
 Glu Ala Pro Asp Val Val Ala Phe Val Ala Pro Gly Glu Ser Gln Gln  
 20 25 30  
 Glu Glu Pro Pro Thr Asp Asn Gln Asp Ile Glu Pro Gly Gln Glu Arg  
 35 40 45  
 Glu Gly Thr Pro Pro Ile Glu Glu Arg Lys Val Glu Gly Asp Cys Gln

50

55

60

Glu Met Asp Leu Glu Lys Thr Arg Ser Glu Arg Gly Asp Gly Ser Asp  
65 70 75 80

Val Lys Glu Lys Thr Pro Pro Asn Pro Lys His Ala Lys Thr Lys Glu  
85 90 95

Ala Gly Asp Gly Gln Pro  
100

&lt;210&gt; 3

&lt;211&gt; 2709

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

```

gaggaaacct ctcgctgggg ctaggagttc ggcgggggcgc gcgcgggcgg ctgcggagct 60
ggcaggtgcg aagcgtctgc acctggcggg cgatggcgcc cgatgcgggc gccccgggat 120
agcgtgggcg agcgtgcggg gccccggcgc gcacgcccgc acctctcccc agccctggcg 180
tgggcccagc ccggcccagg cagcaatggg gttcctgcag ctgctggtcg tagcgggtgct 240
ggcatccgaa caccgggtgg ctggtgcagc cgaggtcttc gggaattcca gcgaggggtct 300
tattgaattt tctgtgggga aatttagata cttcgagctc aataggccct ttccagagga 360
agctattttg catgatattt caagcaatgt gacttttctt attttccaaa tacactcaca 420
gtatcagaat acaactgttt ccttttctcc gactctcctt tccaattcct cggaacacagg 480
cactgccagt ggactggttt tcatccttag accagagcag agtacatgca cttggtactt 540
ggggacttca ggcatacagc ctgtccagaa tatggctatc ctactctcct actcagaaaag 600
agatcctgtc cctggagggt gtaatttgga gttcgattta gatattgatc ccaacattta 660
cttggagtat aatttctttg aaacgactat caagtttgcc ccagcaaacc taggctatgc 720
gagaggcgta gatccccac catgtgacgc tgggacagac caggactcca ggtggaggtt 780
gcagtatgat gtctatcagt attttctgcc tgagaatgac ctactgagg agatgttgct 840
gaagcatctg cagaggatgg tcagtgtgcc ccagggtgaag gccagtgtc tcaagggtgt 900
taccctaaca gctaatagata agacaagtgt ttccttctcc tccctcccgg gacaagggtgt 960
catatacaat gtcattgttt gggaccggtt tctaaataca tctgctgcct acattcctgc 1020
tcacacatac gcttgagctt ttgaggcagg agagggtagt tgtgcttccc taggaagagt 1080
gtcttccaaa gtgttcttca ctctttttgc cctgcttggt ttcttcattt gtttcttttg 1140
acacagcctt tccaaaacag aattattctt cataggcttt atcatcatgg gattcttctg 1200
ttatatactg attacaagac tgacacctat caagtatgat gtgaatctga ttctgacagc 1260
tgtcactgga agcgtcggtg gaatgttctt ggtagctgtg tgggtggcgat ttggaatcct 1320
ctcgcactgc atgctctgtg ttggactagt gctgggggtc ctcatctcgt cagtgaactt 1380
ctttactcca ctgggaaacc taaagatttt tcatgatgat ggtgtattct gggtcacttt 1440
ctcttgcata gctatcctca ttccagtagt tttcatgggc tgcctaagaa tactgaacat 1500
actgacttgt ggagtcattg gctcctattc ggtggtttta gccattgaca gttactgggtc 1560
cacaagcctt tccatcatca ctttgaacgt actcaagaga gcgctcaaca aggatttcca 1620
cagagctttc acaaattgtc cttttcaaac taatgacttc attatcctgg cagtatgggg 1680
catgctggct gtaagtggaa ttacgttaca gattcgaaga gagagaggac gaccgttctt 1740
ccctccccac ccatacaagt tatggaagca agagagagag cgccgagtga caaacattct 1800
ggaccctagc taccacattc ctccattgag agagaggctc tatggccgat taaccagat 1860
taaagggtc ttccagaagg agcagccagc tggagagaga acgcctttgc ttctgtagat 1920
gcccaggggc ttggtcagtg tgccctcagct ttggagttca tgccctggagt ggttcaacag 1980
tctctggtgc aagtctaata agagatcagg catatatatc tgttctttgc ataataattat 2040
ggtgccctta ttgatatatg gtaagggtgt actaggggat taggatgatt gtaagagaat 2100
gagaaagatg accaaaaggt tgggtggtag agggcttttt cttatttcca aatacttgag 2160
aaattacctt ttggtttaca aatctatgat caacttatc cattaaatag atacattaaa 2220
aaaattaaaa actgattctt ctgcagagca ctggtgttct tttttataac cccttgaaac 2280
aagtctctca cstgagcctg tctaaacttt cggagggagt ttattattga gtctttatct 2340
gtgacagtat ttggagattt agggatttga tacttaggcc tttgaatttt agaatacaaa 2400
aagagaagca agccagacat ggtggctcac acctgtaatc ccaatactgg gaggccaaag 2460
tggtgagatc gcttgagccc aggagtttga gaccgacatg ggcaacatga caagacccca 2520

```

```

tctctgcaaa aagattaaaa agttggccag gcatggtggc acatgcctgc tcccagctcc 2580
cggggagact gagatggggg gatccctggg agccctgaag attgaggctg cagtgagcct 2640
tgattgtgtc actgcactcc agcttgggtg acagagaccc tgtctcgaga aattaaaaaa 2700
aaaaaaaaa
2709

```

<210> 4  
 <211> 570  
 <212> PRT  
 <213> Homo sapiens

```

<400> 4
Met Gly Phe Leu Gln Leu Leu Val Val Ala Val Leu Ala Ser Glu His
  1             5             10             15
Arg Val Ala Gly Ala Ala Glu Val Phe Gly Asn Ser Ser Glu Gly Leu
             20             25             30
Ile Glu Phe Ser Val Gly Lys Phe Arg Tyr Phe Glu Leu Asn Arg Pro
             35             40             45
Phe Pro Glu Glu Ala Ile Leu His Asp Ile Ser Ser Asn Val Thr Phe
             50             55             60
Leu Ile Phe Gln Ile His Ser Gln Tyr Gln Asn Thr Thr Val Ser Phe
             65             70             75             80
Ser Pro Thr Leu Leu Ser Asn Ser Ser Glu Thr Gly Thr Ala Ser Gly
             85             90             95
Leu Val Phe Ile Leu Arg Pro Glu Gln Ser Thr Cys Thr Trp Tyr Leu
             100            105            110
Gly Thr Ser Gly Ile Gln Pro Val Gln Asn Met Ala Ile Leu Leu Ser
             115            120            125
Tyr Ser Glu Arg Asp Pro Val Pro Gly Gly Cys Asn Leu Glu Phe Asp
             130            135            140
Leu Asp Ile Asp Pro Asn Ile Tyr Leu Glu Tyr Asn Phe Phe Glu Thr
             145            150            155            160
Thr Ile Lys Phe Ala Pro Ala Asn Leu Gly Tyr Ala Arg Gly Val Asp
             165            170            175
Pro Pro Pro Cys Asp Ala Gly Thr Asp Gln Asp Ser Arg Trp Arg Leu
             180            185            190
Gln Tyr Asp Val Tyr Gln Tyr Phe Leu Pro Glu Asn Asp Leu Thr Glu
             195            200            205
Glu Met Leu Leu Lys His Leu Gln Arg Met Val Ser Val Pro Gln Val
             210            215            220
Lys Ala Ser Ala Leu Lys Val Val Thr Leu Thr Ala Asn Asp Lys Thr
             225            230            235            240
Ser Val Ser Phe Ser Ser Leu Pro Gly Gln Gly Val Ile Tyr Asn Val
             245            250            255
Ile Val Trp Asp Pro Phe Leu Asn Thr Ser Ala Ala Tyr Ile Pro Ala

```

260										265					270				
His	Thr	Tyr	Ala	Cys	Ser	Phe	Glu	Ala	Gly	Glu	Gly	Ser	Cys	Ala	Ser				
		275						280					285						
Leu	Gly	Arg	Val	Ser	Ser	Lys	Val	Phe	Phe	Thr	Leu	Phe	Ala	Leu	Leu				
	290					295					300								
Gly	Phe	Phe	Ile	Cys	Phe	Phe	Gly	His	Arg	Phe	Trp	Lys	Thr	Glu	Leu				
305					310					315				320					
Phe	Phe	Ile	Gly	Phe	Ile	Ile	Met	Gly	Phe	Phe	Phe	Tyr	Ile	Leu	Ile				
				325					330				335						
Thr	Arg	Leu	Thr	Pro	Ile	Lys	Tyr	Asp	Val	Asn	Leu	Ile	Leu	Thr	Ala				
			340					345					350						
Val	Thr	Gly	Ser	Val	Gly	Gly	Met	Phe	Leu	Val	Ala	Val	Trp	Trp	Arg				
	355						360					365							
Phe	Gly	Ile	Leu	Ser	Ile	Cys	Met	Leu	Cys	Val	Gly	Leu	Val	Leu	Gly				
	370					375				380									
Phe	Leu	Ile	Ser	Ser	Val	Thr	Phe	Phe	Thr	Pro	Leu	Gly	Asn	Leu	Lys				
385					390					395				400					
Ile	Phe	His	Asp	Asp	Gly	Val	Phe	Trp	Val	Thr	Phe	Ser	Cys	Ile	Ala				
				405					410					415					
Ile	Leu	Ile	Pro	Val	Val	Phe	Met	Gly	Cys	Leu	Arg	Ile	Leu	Asn	Ile				
			420					425					430						
Leu	Thr	Cys	Gly	Val	Ile	Gly	Ser	Tyr	Ser	Val	Val	Leu	Ala	Ile	Asp				
	435						440					445							
Ser	Tyr	Trp	Ser	Thr	Ser	Leu	Ser	Tyr	Ile	Thr	Leu	Asn	Val	Leu	Lys				
	450					455					460								
Arg	Ala	Leu	Asn	Lys	Asp	Phe	His	Arg	Ala	Phe	Thr	Asn	Val	Pro	Phe				
465					470					475				480					
Gln	Thr	Asn	Asp	Phe	Ile	Ile	Leu	Ala	Val	Trp	Gly	Met	Leu	Ala	Val				
				485					490				495						
Ser	Gly	Ile	Thr	Leu	Gln	Ile	Arg	Arg	Glu	Arg	Gly	Arg	Pro	Phe	Phe				
			500					505					510						
Pro	Pro	His	Pro	Tyr	Lys	Leu	Trp	Lys	Gln	Glu	Arg	Glu	Arg	Arg	Val				
		515					520					525							
Thr	Asn	Ile	Leu	Asp	Pro	Ser	Tyr	His	Ile	Pro	Pro	Leu	Arg	Glu	Arg				
	530					535					540								
Leu	Tyr	Gly	Arg	Leu	Thr	Gln	Ile	Lys	Gly	Leu	Phe	Gln	Lys	Glu	Gln				
545					550					555				560					
Pro	Ala	Gly	Glu	Arg	Thr	Pro	Leu	Leu	Leu										
				565				570											

<210> 5  
<211> 3063  
<212> DNA  
<213> Homo sapiens

<400> 5

```
cgaggcgccg  gtggtgcccg  tggcgccg  gggggagcgc  gggacaggag  gcttcgggga  60
agatggaccc  ggcgcctcgc  ctgggctgca  gcctcaagga  tgtgaagtgg  agctcggtgg  120
ccgtgccgct  cgacctcctg  gtcagcactt  accggctgcc  ccagatcgcg  cgcttgga  180
acggagagtg  cgtagaaggg  ctgccccgaa  atgactatct  gctgattcat  tcctgccgcc  240
agtggaccac  catcactgcc  cacagcttgg  aggagggtca  ctatgtcatt  ggcccaaaga  300
tagagattcc  ggtacattat  gcaggccaat  tcaagctgct  ggaacaagac  cgagatataa  360
aggagccagt  gcaatatttc  aacagtgtgg  aggaggtggc  taaggcattt  cctgaacgcg  420
tgtacgtcat  ggaggatata  acattcaacg  tgaaggttgc  ttcaggtgaa  tgcaatgaag  480
acactgaagt  ttacaacatc  accctgtgta  ctggggatga  actcactcta  atggggggcag  540
gcagaaatcc  ttcatgcaaa  gacattcaag  gaaaagtcac  gactcaacac  aatcttcaaa  600
aagattggga  agctcaattc  catcagcaag  ctgggaaaag  gcaaaatgcc  gtgcctcatt  660
tgtatgaatc  accggacca  cgaaagcatt  agccttccat  tccagtgcga  gggcagattt  720
agcaccgccg  agtccccctg  aacttcagat  gcaaaagggc  gaacacacca  tccgccacat  780
tgtggagaaa  accaggcttc  ctgtgaatgt  gactgtgcca  agcctccac  cgagaaaccc  840
atacgacctc  cacttcaccc  gtgaggggca  ccgctataag  tttgtgaaca  tccagacca  900
gacggtgggtg  gtttgctgtg  tgctgcggga  caacaagatc  ctccccatgc  actttccttt  960
gcacttgact  gtccccaaat  tcagcctccc  agaacacctg  gtgaaggagg  agagctggcc  1020
cgaaaccctg  gtccatcact  ggctagggtat  ctggcaagaa  cagttcgaca  tcgatgagta  1080
ttcacgggct  gtccgtgatg  tgaaaaccga  ctggaatgaa  gaatgcaaga  gccccagaa  1140
gggtcggtgc  tctggccaca  accacgtgcc  caattcgctc  agctacgccc  gcgatgagct  1200
caccagctcc  ttccaccgac  tctcggtctg  tgtgtatggc  aacaatctcc  atggcaacag  1260
tgaggtgaac  ctcatggtt  gcagggacct  ggggggagat  tgggctccct  ttctcatga  1320
catcctgccc  tatcaggact  ctggagatag  tgggagcgac  taccttttcc  cagaagctag  1380
tgaagaatca  gcaggcatcc  cgggaaagtc  agaacttccc  tacgaagagc  tgtggctgga  1440
ggaaggcaag  cccagccatc  agcctctcac  ctgctctctg  agcgagaaga  acagatgtga  1500
tcagtttaga  ggttctgtcc  gatccaaatg  tgcgacttct  cctcttccca  tccctgggac  1560
tctgggagca  gcagtgaagt  ctccagatac  tgccctacct  ccacctccag  tgcctcccaa  1620
atctgaagcc  gtcagagaag  aatgccggct  cctgaacgcc  ccacctgttc  caccgccgaag  1680
cgcaaaacct  ttgtccacca  gtccctccat  ccctcctcgc  acagtcaagc  cagcgcgga  1740
acagactcgc  tctcccagcc  ccacctgtgc  ctactattct  tcagggctac  acaacatcgt  1800
cactaaaact  gacacaaatc  cttctgaaag  cactcctggt  tctgtctatc  catgtaaccg  1860
agtgaaaact  gattctgtgg  acctgaaatc  accgtttgga  agtccttctg  ctgaagctgt  1920
gtcctctcgg  ctctcatggc  ctaaccatta  ttcaggagca  tcagaaagcc  agaccaggag  1980
tgacttctcg  ctggatccaa  gcaggagtta  tagttacct  agacaaaaga  cgccaggcac  2040
accaagaga  aactgccag  caccttttga  ttttgatggc  tgtgagctcc  tggccagccc  2100
cactagccca  gtcactgcag  aattcagtag  cagcgtctct  ggttggtcca  agtcagccag  2160
ctactctctg  gagagcacag  atgtgaaatc  tcttgcaagt  ggtgtgacaa  agcagagtac  2220
gtcatgccc  gccttcccc  ccagggtccc  aaaactagt  gaagagaagg  tgcctccga  2280
aacatctcct  ttgcctctga  aaattgatgg  tgctgaggaa  gacccaagt  ctgggtcacc  2340
agatctctcg  gaggaccagt  attttgttaa  aaagggcag  caggacatct  tctctgcctc  2400
ctacctttc  tcatctccgc  tccatctcca  gctggcccc  agatcctgtg  gcgacggttc  2460
cccatggcag  ccacctgctg  acctatcagg  actctctata  gaggaagtgt  ccaagtcact  2520
acggttcatt  ggtttgtccg  aagatgtcat  atcattcttt  gttactgaaa  agattgatgg  2580
gaacctgctt  gttcagctaa  cggaagaaat  cctctcagag  gatttcaaat  tgagcaaat  2640
gcagggtgaag  aagataatgc  aattcattaa  tggttgagg  cccaaaatat  agccaaataa  2700
cccccgcca  gcatggaaca  aaactgatca  atgcgtgtgc  tagaaggggt  gggctgggac  2760
acaatttcat  gtttttgac  taaaaacctt  ctctgtaaat  agggataaga  gaaactctta  2820
ctatgcagat  tacgtttttg  aatgggtgaac  aggtatattt  gtacatcaat  aaaaatgctg  2880
tacagaacac  ttggaggtgt  gccttgtagc  tcaactcaaca  aacactcagc  agctgctaaa  2940
agaaaaaaag  gcatgtgcag  agaaatcatt  cttacccaag  taggtttatg  tgagaaggta  3000
tgatatttat  taaaaaatag  ccaaagctga  aagacataaa  aatctttaaa  aaaaaaaaaa  3060
aaa  3063
```

<210> 6



&lt;211&gt; 647

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

```

Met Gln Lys Gly Glu His Thr Ile Arg His Ile Val Glu Lys Thr Arg
  1           5           10           15

Leu Pro Val Asn Val Thr Val Pro Ser Pro Pro Pro Arg Asn Pro Tyr
          20           25           30

Asp Leu His Phe Ile Arg Glu Gly His Arg Tyr Lys Phe Val Asn Ile
  35           40           45

Gln Thr Lys Thr Val Val Val Cys Cys Val Leu Arg Asp Asn Lys Ile
  50           55           60

Leu Pro Met His Phe Pro Leu His Leu Thr Val Pro Lys Phe Ser Leu
  65           70           75           80

Pro Glu His Leu Val Lys Gly Glu Ser Trp Pro Glu Thr Leu Val His
          85           90           95

His Trp Leu Gly Ile Cys Gln Glu Gln Phe Asp Ile Asp Glu Tyr Ser
  100           105           110

Arg Ala Val Arg Asp Val Lys Thr Asp Trp Asn Glu Glu Cys Lys Ser
  115           120           125

Pro Lys Lys Gly Arg Cys Ser Gly His Asn His Val Pro Asn Ser Leu
  130           135           140

Ser Tyr Ala Arg Asp Glu Leu Thr Gln Ser Phe His Arg Leu Ser Val
  145           150           155           160

Cys Val Tyr Gly Asn Asn Leu His Gly Asn Ser Glu Val Asn Leu His
          165           170           175

Gly Cys Arg Asp Leu Gly Gly Asp Trp Ala Pro Phe Pro His Asp Ile
          180           185           190

Leu Pro Tyr Gln Asp Ser Gly Asp Ser Gly Ser Asp Tyr Leu Phe Pro
  195           200           205

Glu Ala Ser Glu Glu Ser Ala Gly Ile Pro Gly Lys Ser Glu Leu Pro
  210           215           220

Tyr Glu Glu Leu Trp Leu Glu Glu Gly Lys Pro Ser His Gln Pro Leu
  225           230           235           240

Thr Arg Ser Leu Ser Glu Lys Asn Arg Cys Asp Gln Phe Arg Gly Ser
          245           250           255

Val Arg Ser Lys Cys Ala Thr Ser Pro Leu Pro Ile Pro Gly Thr Leu
          260           265           270

Gly Ala Ala Val Lys Ser Ser Asp Thr Ala Leu Pro Pro Pro Pro Val
  275           280           285

Pro Pro Lys Ser Glu Ala Val Arg Glu Glu Cys Arg Leu Leu Asn Ala

```

290	295	300
Pro Pro Val Pro Pro Arg Ser Ala Lys Pro Leu Ser Thr Ser Pro Ser		
305	310	315 320
Ile Pro Pro Arg Thr Val Lys Pro Ala Arg Gln Gln Thr Arg Ser Pro		
	325	330 335
Ser Pro Thr Leu Ser Tyr Tyr Ser Ser Gly Leu His Asn Ile Val Thr		
	340	345 350
Lys Thr Asp Thr Asn Pro Ser Glu Ser Thr Pro Val Ser Cys Tyr Pro		
	355	360 365
Cys Asn Arg Val Lys Thr Asp Ser Val Asp Leu Lys Ser Pro Phe Gly		
	370	375 380
Ser Pro Ser Ala Glu Ala Val Ser Ser Arg Leu Ser Trp Pro Asn His		
	385	390 395 400
Tyr Ser Gly Ala Ser Glu Ser Gln Thr Arg Ser Asp Phe Leu Leu Asp		
	405	410 415
Pro Ser Arg Ser Tyr Ser Tyr Pro Arg Gln Lys Thr Pro Gly Thr Pro		
	420	425 430
Lys Arg Asn Cys Pro Ala Pro Phe Asp Phe Asp Gly Cys Glu Leu Leu		
	435	440 445
Ala Ser Pro Thr Ser Pro Val Thr Ala Glu Phe Ser Ser Ser Val Ser		
	450	455 460
Gly Cys Pro Lys Ser Ala Ser Tyr Ser Leu Glu Ser Thr Asp Val Lys		
	465	470 475 480
Ser Leu Ala Ala Gly Val Thr Lys Gln Ser Thr Ser Cys Pro Ala Leu		
	485	490 495
Pro Pro Arg Ala Pro Lys Leu Val Glu Glu Lys Val Ala Ser Glu Thr		
	500	505 510
Ser Pro Leu Pro Leu Lys Ile Asp Gly Ala Glu Glu Asp Pro Lys Ser		
	515	520 525
Gly Ser Pro Asp Leu Ser Glu Asp Gln Tyr Phe Val Lys Lys Gly Met		
	530	535 540
Gln Asp Ile Phe Ser Ala Ser Tyr Pro Phe Ser Ser Pro Leu His Leu		
	545	550 555 560
Gln Leu Ala Pro Arg Ser Cys Gly Asp Gly Ser Pro Trp Gln Pro Pro		
	565	570 575
Ala Asp Leu Ser Gly Leu Ser Ile Glu Glu Val Ser Lys Ser Leu Arg		
	580	585 590
Phe Ile Gly Leu Ser Glu Asp Val Ile Ser Phe Phe Val Thr Glu Lys		
	595	600 605
Ile Asp Gly Asn Leu Leu Val Gln Leu Thr Glu Glu Ile Leu Ser Glu		

610

615

620

Asp Phe Lys Leu Ser Lys Leu Gln Val Lys Lys Ile Met Gln Phe Ile  
 625 630 635 640

Asn Gly Trp Arg Pro Lys Ile  
 645

&lt;210&gt; 7

&lt;211&gt; 892

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

```

ggcacgagct cgtgcactca tggcgaccgc gaacccccct cccaagact atgaaagtga 60
tgacgactct tatgaagtgt tggatttaac tgagtatgcc agaagacacc agtggtggaa 120
tcgagtgttt ggccacagtt cgggacctat ggtagaaaaa tactcagtag ctaccagat 180
tgtaatgggt ggcgttactg gctggtgtgc aggatttctg ttccagaaag ttggaaaact 240
tgcagcaact gcagtaggtg gtggttttct tcttcttcag attgctagtc atagtggcta 300
tgtgcagatt gactggaaga gagttgaaaa agatgtaaat aaagcaaaaa gacagattaa 360
gaaacgagcg aacaaagcag cacctgaaat caacaattta attgaagaag caacagaatt 420
tatcaagcag aacattgtga tatccagtgg atttgtggga ggctttttgc tcggacttgc 480
atcttaagga catgaatatt ctcccataac ggattcaact atgagaagag aagtggcagc 540
aataaggcag tctctcaaaa gtcatactgc cagagtctct agggcaagga gaaacaacta 600
gctggacaat actcaattca caacttagca ttttgccatc tgaagcttgg caaactagta 660
tctgctgtaa aacaacctat atggtatgtg aaccgtagta ttcctgagca aaacgtggct 720
ttcatcgctt tgtaaaaatt tgcattctgt tagaaactag cctataaaat atcaccattg 780
gatgtagata tggagagaaa agaaatatgt tgggtttatt gcttagcgaa atattctctt 840
tttattttaa taaaatgttc ttcattgtgt tttaaaaaaa aaaaaaaaaa aa 892

```

&lt;210&gt; 8

&lt;211&gt; 155

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

```

Met Ala Thr Arg Asn Pro Pro Pro Gln Asp Tyr Glu Ser Asp Asp Asp
  1           5           10           15

Ser Tyr Glu Val Leu Asp Leu Thr Glu Tyr Ala Arg Arg His Gln Trp
      20           25           30

Trp Asn Arg Val Phe Gly His Ser Ser Gly Pro Met Val Glu Lys Tyr
      35           40           45

Ser Val Ala Thr Gln Ile Val Met Gly Gly Val Thr Gly Trp Cys Ala
      50           55           60

Gly Phe Leu Phe Gln Lys Val Gly Lys Leu Ala Ala Thr Ala Val Gly
      65           70           75           80

Gly Gly Phe Leu Leu Leu Gln Ile Ala Ser His Ser Gly Tyr Val Gln
      85           90           95

Ile Asp Trp Lys Arg Val Glu Lys Asp Val Asn Lys Ala Lys Arg Gln
      100          105          110

Ile Lys Lys Arg Ala Asn Lys Ala Ala Pro Glu Ile Asn Asn Leu Ile
      115          120          125

```

Glu Glu Ala Thr Glu Phe Ile Lys Gln Asn Ile Val Ile Ser Ser Gly  
 130 135 140

Phe Val Gly Gly Phe Leu Leu Gly Leu Ala Ser  
 145 150 155

<210> 9  
 <211> 1850  
 <212> DNA  
 <213> Homo sapiens

<400> 9  
 cactcctact gcggtctgcta tgaagcttac tgggtgtgat gtgttataat ttagtctgtt 60  
 tttttgattg aatgcagttt aatgtttcca gaaagccaaa gtaattttct ttccagatat 120  
 gcaaggcttt ggtgggtcca aaaaatgtct atcacaagcc attttttccct ttccctctct 180  
 cgaaaagtta aaatatctat gtgttattcc caaacctctt tacctatgta tctgcctgtc 240  
 tgtccatcat ctcccttccct ccctatctct gtgtatctgg atggcagccg ctgcccargg 300  
 gagtggctgt ggggagggca ggtactgtct ttgcctgtgg gtccagctga gccatccctg 360  
 ctgggtgatg ctgggcaaga cccttggccc gtctgggcct tggtctctct acttgtgaaa 420  
 tgagcgggaa gatgactctc agttccctcc acctcttaga catggtgagg taacagacat 480  
 caaaagcttt tctgaaatct tcagaagaaa tagttccatt acagaaaact cttcaaaata 540  
 aatagtagtg aaaactttta aaaactctca ttggagtaag tcttttcaag atgacctcc 600  
 acaatggagg cagcgttccct acttgtcatc acacagctga agacattgtt tcttaggtgt 660  
 gaaatcgggg acaaaggaca aacagagaca cagggcattg ttcattgggag gcacgtcac 720  
 cctcctgggt gttctgtggg aatttccctgt gtgaggaaaa cgtggccaca ggggtgtgct 780  
 gtaccacccc ttccccggcg agatggccct cggcctgtgc cgtgcttcc acctcgcga 840  
 ctccatggca gcttttggtc tgtttccggc tctgccctct gccctgaact ctcatccggc 900  
 ttgtacctgc ctgctggacc cctccacctg gaggccagcc catgtctcag gccagccct 960  
 agcctcttct cctcaaattc taagtgtttt ctctttaggt ttccctggct ttgtgaatgg 1020  
 atcatgtgtc tctaggtata aacctgacat catctctcca cccggcttac ctccaccaga 1080  
 tctcccaggt tctgtctcca tcttctacct gcagctgtc tgttctcatg gtcactgtct 1140  
 catcactgag tctggacctt gtgtatcatt ttcaaaactgg cctccttccc tegtcccca 1200  
 cttcttaaag tcacctgtcc attgccacca gattaagctt tctccagcca gatcacctct 1260  
 ctctgagaaa cctccattga catggaaaca ccattgtctg gcacacatac tcacatactc 1320  
 accttcccggt cttgatcccc acacatcttt ccagcctccc ctccactcc actccttgc 1380  
 cctcctcca cctccccatc ctcttgtctc cctccccctc tgaatccagc ccagcggggc 1440  
 ttctcctgcc tccatcacat cacagaagta cctcctgctt ctggttttaa ttagagcctt 1500  
 ccccgattac attttctct gaattttttc ctatctacat ttgatctgtc atgtttaaac 1560  
 cccctacttc taagggaact tctctaactc cttatcctca tccccaaata gtgttttctt 1620  
 cctctgggtt cttataatgt tgggtatcaat ctcacagcat ttagtgcttc ctgectgggt 1680  
 tgacagttac ctgtgtgcat gtgcaatttc taatttccca cgctagactg tgagcttcc 1740  
 aaggcaagaa tcatgccttg ttggtttctg tattcctcat ggtgccaaac acagtgcctt 1800  
 ctacattgca ggcgctgaat aaacattttt aaagcaaaaa aaaaaaaaaa 1850

<210> 10  
 <211> 206  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Met Ala Leu Gly Leu Cys Arg Cys Phe His Pro Arg His Ser Met Ala  
 1 5 10 15  
 Ala Phe Gly Leu Phe Pro Ala Leu Pro Ser Ala Leu Asn Ser His Pro  
 20 25 30  
 Ala Cys Thr Cys Leu Leu Asp Pro Ser Thr Trp Arg Pro Ala His Val  
 35 40 45

Ser Gly Pro Ala Leu Ala Ser Ser Pro Gln Ile Leu Ser Val Phe Ser  
 50 55 60  
 Leu Gly Phe Pro Gly Phe Val Asn Gly Ser Cys Val Ser Arg Tyr Lys  
 65 70 75 80  
 Pro Asp Ile Ile Ser Pro Pro Gly Leu Pro Pro Pro Asp Leu Pro Ser  
 85 90 95  
 Ser Val Ser Ile Phe Tyr Leu Gln Leu Leu Cys Ser His Gly His Cys  
 100 105 110  
 Cys Ile Thr Glu Ser Gly Pro Leu Leu Ser Phe Ser Asn Trp Pro Pro  
 115 120 125  
 Ser Leu Val Pro His Phe Leu Lys Ser Pro Val His Cys His Gln Ile  
 130 135 140  
 Lys Leu Ser Pro Ala Arg Ser Pro Leu Ser Glu Lys Pro Pro Leu Thr  
 145 150 155 160  
 Trp Lys His His Cys Leu Ala His Ile Leu Thr Tyr Ser Pro Ser Arg  
 165 170 175  
 Leu Asp Pro His Thr Ser Phe Gln Pro Pro Leu Pro Leu His Ser Leu  
 180 185 190  
 Leu Pro Pro Pro Pro Pro His Pro Leu Val Ser Pro Pro Leu  
 195 200 205

&lt;210&gt; 11

&lt;211&gt; 2216

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

cttgtaagtt actgtagtg aattgttttt tacgtttcat ttaataattg ctgctaaagg 60  
 tgatgttttac tgataaatca ttttaaaatt tttttgtttt gaaaagtaaa tttatccccc 120  
 atgatgttag atacatttaa attattaagt cttttcagag atgagatggg gacaggaagt 180  
 tatttttgagc cttacaatat tatttagccc aataaaagat gcattgaagc tcttatatat 240  
 tatgagtttg aaaaattttg aaggtagcat attgaagtga tctataaata tcttcagtc 300  
 tctctgaagt gtgggtattt cttctatcta aaaaatacat acagtgactg tcttcaaate 360  
 tacttggttc ttgaccaaata argagctaata gggtaatgaa tacctttttg tttgtttgtt 420  
 tgtttggttg tttttgtttt ttttttttaa ggggtctcact cttttgcccc ggctggagtg 480  
 cagtggcaca atcacggctc ccaggctaata gtttttattt ttaatttgta attttttttt 540  
 tatttttttt gttgagatgg agttgctcca tgttgcacag gctgttctca aactcctaag 600  
 ctcaagccat ctgcctgcst tggcctcccc aagtgstggg attgtagaca taagccacct 660  
 caccagcct atgaatatct ttctaataatg gtaagaatga ggtaatgttt ccatcagtc 720  
 aatacagata tatttcttcc ctccaaaaca gtttatattt attgtttatt ttattttgat 780  
 tgtaactccg tcataactyg acatggaaaa tgctatatac tatgaaaact tagctgaaag 840  
 ggaagaattg ttttagaaaag acaatatttta aaacaccgca ctgccaatat attgatcctt 900  
 tatagttatt tcctaaaatg ctgttttcga aacattcctt tttcaccctg ttgtgtggct 960  
 tagaccatc tcgtaatctg ttaattggaa agaggctaca gacaccagca gtgtgcgttc 1020  
 tgcagggtaca cgctgcaaaa gtaattcctg ctcattccatg ccctgtctct gtctctttta 1080  
 gagtcatacc ttatttgagt ataggtggct taattttgct agacttctctg aaaacactaa 1140  
 ggtggagtat cagaagtgat ttagtcaca gttctgctgg agagcttaga ataacatcct 1200  
 cctttgggag gtggtcttgg gtgcgtggat cttggtatac agtctttatt gtaagtcctga 1260  
 tacaaaatgc taataaattt aatgtttttc ttccttaatt tattggcata gttcttcagg 1320

```

tagcacctca tttttattaa tgatattggg attaactatg aacaagctat atgtagacat 1380
ttgcatttaa ggacattgca gtgtttcaaa gatcccatca ttgcagcttg tatccttttag 1440
atccaatcgg aaactttctgg agtcttacat taatgctcat ttgagctaata tagtaatctg 1500
tttaaacaga tttggcaata ctttaaagat actgtagact atttatgtat agatagatca 1560
tattacccat taaaagtctg ggggaaaaaa ttttttaatt ttactcttct tatgtactga 1620
aaactttttt taaaaaagggt gatgatgaag ttcattctgt agcagcagcg cagctatgct 1680
ttaaaccaca caaaaggctg tgtccagggtg cagcctcctt cacccttctt gcccacggtg 1740
aggattgaat aaccaggact tggggatatk gtttggtgtc aggggttattc tgtgtggtaa 1800
ggaatatttg tttcacattt atacattttc tttttccact cacgtaagtt tctatcttga 1860
gagcatagtc caaagtgcaa aacttggtgt ttacaaggaa aattgtcttc cagaactcca 1920
ctgtcatcac tttcaccaaa gtggaagttt gcatgaatat gctcagaatc taatattcaa 1980
tgttctgtta cattgtaagt gaagtccagc tacaaaatag atttaataata ttgaatttat 2040
ttgtacatat gcagagtacg gtatttctgt atggaatctg ctttattcct atttttccca 2100
actctgatga gtagaatatt aaatgtgttg ttatggaaat acagattatt gcttctatag 2160
gaagataatt atgaaaataa aacctgaaac tatataaata taaaaaaaaa aaaaaa 2216

```

&lt;210&gt; 12

&lt;211&gt; 126

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

```

Met Leu Phe Ser Lys His Ser Phe Phe Thr Leu Leu Cys Gly Leu Asp
  1                      5                      10                      15

Pro Ser Arg Asn Leu Leu Ile Gly Lys Arg Leu Gln Thr Pro Ala Val
          20                      25                      30

Cys Val Leu Gln Val His Ala Ala Lys Val Ile Pro Ala His Pro Cys
          35                      40                      45

Pro Val Ser Val Ser Phe Arg Val Ile Pro Tyr Leu Ser Ile Gly Gly
          50                      55                      60

Leu Ile Leu Leu Asp Phe Leu Lys Thr Leu Arg Trp Ser Ile Arg Ser
          65                      70                      75                      80

Asp Phe Ser His Ser Ser Ala Gly Glu Leu Arg Ile Thr Ser Ser Phe
          85                      90                      95

Gly Arg Trp Ser Trp Val Arg Gly Ser Trp Tyr Thr Val Phe Ile Val
          100                      105                      110

Ser Leu Ile Gln Asn Ala Asn Lys Phe Asn Val Phe Leu Pro
          115                      120                      125

```

&lt;210&gt; 13

&lt;211&gt; 1426

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

```

ctgggtctcc agggggagag cctggccctg tccttttgcta cccagggctg cccccaggcc 60
catgaagcca ataggagagc gtgtggcact ggccccacaaa ctgtccctgt cctgtcttcc 120
tcccagacca tggcctctgc tagctccacc ttgaaggagc cccccacatc ctcccctaca 180
tcccagagat gccaccactt gtgtctccac aatgtgtctc tgcccaccog ggttcgcgac 240
tgtccgaccc ctgcacacca ctcatgtcac caggcgctgc atcatgttca tccccatcta 300
tttatttaag ctttctttg cttgtagggc attttgtatg tagagcagtt gaaaacagaa 360
cctcagaact taacatctgt cctgatgtta aagtgccttt catgaccacc ctgttatcta 420

```

```

tgtatatgta aagttaagga tgagatctta agttttacaat taaaaaactca gtactcaata 480
tttaatatcc tactcgagct ttatggaagc caaatcatgc atgtgtgtgt gtgcgtgtgt 540
gcaagctttg aacctccttc cacagccgca tcttctcatg acacaaagct tttgataagt 600
actttcctgt gggctcgtca gggcctcata gcatctcatt caattacaag aatagaggcc 660
agacacgggtg gcgcattgctt gtagtcccag ctaactggga ggctgaggca ggaggatcac 720
ttgagcccag gagattgagg ctgcagttag catgatcgcg aactgcaact ccagcctggg 780
tgacgggtgag actttgtctc aaaaaaaaaa aaaaaaaca tggaaggcag acagcaagtc 840
cctgaggaca catcacacag tgtcctgtag ctaagtgtct aggaaaaaac aaaaactcca 900
aacccttcag tggatgagga caaggtcgca gaaaggcatt ctggtgacag atgaacagcc 960
gaaagctggc cagaccctcc tgtatgcctc tgcccttgct ctgtgggttg agggggtctg 1020
accaggaggc cacctacagc aggaagttag gctgccatgt ttccttgaga cacagctgcc 1080
tctccccagc tctgtccctg tagtcacctg ccggtgggag aggatcctct ccctgggata 1140
agcactccca gccccgttta tcagaaacac aggaaggaa attggaactg ccaccagcc 1200
cagcattggg gctcaattgg ttggttgctg tgtcagttgt ctcttcgttt tgttaagggt 1260
tttaataagt acgtttggca taatgtcttt tacttaggga aaaaaacaat ttgtaaatac 1320
agcagcctat aacttttcag ctggtgcttt tacttaggga aaaaaacaat ttgtaaatac 1380
agaacattgt ttaaaagaca taaccataga aaaaaaaaaa aaaaaa 1426

```

&lt;210&gt; 14

&lt;211&gt; 80

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

```

Met Pro Pro Leu Val Ser Pro Gln Cys Ala Pro Ala His Pro Gly Ser
  1                      5                      10                      15

```

```

Ala Leu Ser Asp Pro Cys Thr Pro Leu Met Ser Pro Arg Arg Ala Ser
          20                      25                      30

```

```

Cys Ser Ser Pro Ser Ile Tyr Leu Ser Leu Ser Leu Leu Val Gly His
          35                      40                      45

```

```

Phe Val Cys Arg Ala Val Glu Asn Arg Thr Ser Glu Leu Asn Ile Cys
          50                      55                      60

```

```

Pro Asp Val Lys Val Leu Phe Met Thr Thr Leu Leu Ser Met Tyr Met
          65                      70                      75                      80

```

&lt;210&gt; 15

&lt;211&gt; 2364

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

```

gaagcggctg ctgtagggcg cgacggagcg agcggggcgtg cggagcgggc gacagtggcg 60
tgggatctgc ctctctgcga gcagctggga gcggcgcccg cggcgccatg agcgggggca 120
ccccctacat cggcagcaag atcagcctca tctccaaggc ggagatccgc tacgagggca 180
tcctctacac catcgacacc gaaaactcca ccgtagccct tgccaaagt ttcgattctt 240
gtacagaaga cagaccgaca gatcgtccaa taccacctcg agatgaagtc tttgaataca 300
ttatattccg tgggagtgac attaaagacc ttactgtttg tgagccacca aaaccacagt 360
gttctttgcc tcaagaccca gctattgttc agtcctcact aggtcctatg acttcttcat 420
tccagtccat gggttcttat ggacctttcg gcaggatgcc cacatacagt cagttcagtc 480
cgagttcctt agttgggcag cagtttggtg ctgttggtgt tgctggaagc tctttgacat 540
cctttggaac agaaacatca aacagtggta ccttacccca aagtagtgcg gttgggttctg 600
cctttacaca ggatacaaga tctctaaaaa cacagttatc tcaaggctcg tcaagccctc 660
agttagaccc tttgagaaaa agcccaacca tggaacaagc agtgacagcc gcctcagccc 720
acttacctgc tccagcagct gttgggagaa ggagtcctgt atcaaccagg cctttgccat 780
ctgccagcca aaaggcagga gagaatcagg agcacaggca agctgaagta cacaagttt 840

```

```

caaggccaga aatgagcaa ctcagaaatg ataacaagag acaagtagct ccagggtgctc 900
cttcagctcc aaggagaggg cgtggggggtc atcggggttg caggggaaga tttggtattc 960
ggcgagatgg gccaatgaaa tttgagaaaag actttgactt tgaaagtgca aatgcacaat 1020
tcaacaagga agagattgac agagagtttc ataataaact taaattaaaa gaagataaac 1080
ttgagaaaaca ggagaagcct gtaaatggtg aagataaagg agactcagga gttgataccc 1140
aaaacagtga aggaatgcc gatgaagaag atccacttgg acctaattgc tattatgaca 1200
aaactaaatc cttctttgat aatatttctt gtgatgacaa tagagaacgg agaccaacct 1260
gggctgaaga aagaagatta aatgctgaaa catttggaat ccacttcgt ccaaaccgtg 1320
gccgtggggg atacagaggg agaggaggtc ttggtttccg tggtggcaga gggcgtggtg 1380
gtggcagagg tggtagcttc actgcccctc gaggatttcg cgggtggattc agaggaggtc 1440
gtggggggcgg ggagtttgcg gattttgaat ataggaaaac cacagctttt ggaccctaaa 1500
aggtctggat tgatcgtact gctttctgaa agaaaagacaa caaagttgct gcatagtcta 1560
caaacaagtc tctgaaaata ggtgaatttc tagctcttca tggctcctgaa cattgatttc 1620
agtctttgca aagaatgaag aagtgaattc gctgtacatt tgtcaccagc actgggtttt 1680
tggtttttgt ttgtttttcc gcttaatttc aaagataaaa tgcagttact tttgggggtg 1740
gaaggctcat cttaaaacat gagcattaaa tatatttggg atagcagaag gttaagtaat 1800
ttcttatgta tagttaaact aaagcagtac ttcagtggga cttaacaagt atttttcat 1860
cactgaaagg tttttttttt tttatcacta aattgtattt ggcaattgca agttgcctgc 1920
agatagggcc gtgatactgt gttttgagcc acagaagggt gtgtgtgtgt gtgtgtgtgt 1980
gtgtgtgtgt gtgtgtgtgt gtatgtgtgt gtctttttcc tcctttcttt tggggaatcc 2040
tgtaatatga ggtagcttat ttcgtcaatt aattaggggt ctggatggta gagaattttg 2100
tcagtcaact atgtacacac agtaataact gtttcttagg caaaggtaac ttttttatat 2160
agttgtaaaa ttccattata ttccattgcc aaagaaacat taagaccttt gtatagctgt 2220
ataaaaagca actaattttt taaagaaata aacattttta agtccaaaaa aaaaaaaa 2280
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2340
aaaaaaaaa aaaaaaaaaa aaaa                                     2364

```

&lt;210&gt; 16

&lt;211&gt; 463

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

```

Met Ser Gly Gly Thr Pro Tyr Ile Gly Ser Lys Ile Ser Leu Ile Ser
  1              5              10              15

Lys Ala Glu Ile Arg Tyr Glu Gly Ile Leu Tyr Thr Ile Asp Thr Glu
      20              25              30

Asn Ser Thr Val Ala Leu Ala Lys Val Arg Ser Phe Gly Thr Glu Asp
      35              40              45

Arg Pro Thr Asp Arg Pro Ile Pro Pro Arg Asp Glu Val Phe Glu Tyr
      50              55              60

Ile Ile Phe Arg Gly Ser Asp Ile Lys Asp Leu Thr Val Cys Glu Pro
      65              70              75              80

Pro Lys Pro Gln Cys Ser Leu Pro Gln Asp Pro Ala Ile Val Gln Ser
      85              90              95

Ser Leu Gly Ser Ser Thr Ser Ser Phe Gln Ser Met Gly Ser Tyr Gly
      100              105              110

Pro Phe Gly Arg Met Pro Thr Tyr Ser Gln Phe Ser Pro Ser Ser Leu
      115              120              125

Val Gly Gln Gln Phe Gly Ala Val Gly Val Ala Gly Ser Ser Leu Thr
      130              135              140

```



Ser Phe Gly Thr Glu Thr Ser Asn Ser Gly Thr Leu Pro Gln Ser Ser  
 145 150 155 160  
 Ala Val Gly Ser Ala Phe Thr Gln Asp Thr Arg Ser Leu Lys Thr Gln  
 165 170 175  
 Leu Ser Gln Gly Arg Ser Ser Pro Gln Leu Asp Pro Leu Arg Lys Ser  
 180 185 190  
 Pro Thr Met Glu Gln Ala Val Gln Thr Ala Ser Ala His Leu Pro Ala  
 195 200 205  
 Pro Ala Ala Val Gly Arg Arg Ser Pro Val Ser Thr Arg Pro Leu Pro  
 210 215 220  
 Ser Ala Ser Gln Lys Ala Gly Glu Asn Gln Glu His Arg Gln Ala Glu  
 225 230 235 240  
 Val His Lys Val Ser Arg Pro Glu Asn Glu Gln Leu Arg Asn Asp Asn  
 245 250 255  
 Lys Arg Gln Val Ala Pro Gly Ala Pro Ser Ala Pro Arg Arg Gly Arg  
 260 265 270  
 Gly Gly His Arg Gly Gly Arg Gly Arg Phe Gly Ile Arg Arg Asp Gly  
 275 280 285  
 Pro Met Lys Phe Glu Lys Asp Phe Asp Phe Glu Ser Ala Asn Ala Gln  
 290 295 300  
 Phe Asn Lys Glu Glu Ile Asp Arg Glu Phe His Asn Lys Leu Lys Leu  
 305 310 315 320  
 Lys Glu Asp Lys Leu Glu Lys Gln Glu Lys Pro Val Asn Gly Glu Asp  
 325 330 335  
 Lys Gly Asp Ser Gly Val Asp Thr Gln Asn Ser Glu Gly Asn Ala Asp  
 340 345 350  
 Glu Glu Asp Pro Leu Gly Pro Asn Cys Tyr Tyr Asp Lys Thr Lys Ser  
 355 360 365  
 Phe Phe Asp Asn Ile Ser Cys Asp Asp Asn Arg Glu Arg Arg Pro Thr  
 370 375 380  
 Trp Ala Glu Glu Arg Arg Leu Asn Ala Glu Thr Phe Gly Ile Pro Leu  
 385 390 395 400  
 Arg Pro Asn Arg Gly Arg Gly Gly Tyr Arg Gly Arg Gly Gly Leu Gly  
 405 410 415  
 Phe Arg Gly Gly Arg Gly Arg Gly Gly Arg Gly Gly Thr Phe Thr  
 420 425 430  
 Ala Pro Arg Gly Phe Arg Gly Gly Phe Arg Gly Gly Arg Gly Gly Arg  
 435 440 445  
 Glu Phe Ala Asp Phe Glu Tyr Arg Lys Thr Thr Ala Phe Gly Pro  
 450 455 460

<210> 17  
 <211> 2760  
 <212> DNA  
 <213> Homo sapiens

<400> 17  
 tgaagatgcc tcctctgatg cctactgctt tgagctgctc tctatggttt tagcactgag 60  
 tggctctaac gttggcggc aatatctggc tcaacagcta accctgcttc aggatctctt 120  
 ctcgctgctt cacacagcct ctccatagagt ccagagacag gtaacctctt tactaagaag 180  
 agttttgcct gaagtaacct ctagtctgtt ggccagcatc ataggagtga aatccctccc 240  
 cccagcagat atcagtgata tcattcactc aacagagaaa ggagactgga ataagctggg 300  
 tatcttgagc atgtttctag gatgcattgc caaagcactc actgtacagc taaaagccaa 360  
 aggaaccacc atcactggaa cagctgggtac cactgtgggc aaaggagtta caacagttac 420  
 tcttccgatg attttcaatt ccagttatct ccgacgaggt gaaagtcatt ggtggatgaa 480  
 gggctcaacc cctaccaga tctcagagat catcattaaa cttatcaagg atatggcagc 540  
 aggtcatctg tcagaagctt ggtcccgagt gacaaaaaat gctattgcag aaaccatcat 600  
 tgccttgacc aagatggaag aagaatttag gtctccagtg agatgtattg caacaactag 660  
 actctggctt gctctcgcat ccctatgtgt tcttgatcag gaccacgtag atcgtctctc 720  
 ctggtggaga tggatgggaa aggatggaca acaaaaacaa atgcctatgt gtgataacca 780  
 tgatgatggt gaaactgcag caatcatttt atgcaatgtc tgtggaaatt tatgtacaga 840  
 ctgtgacaga ttccttcacc ttcacgaag aacaaaaact catcaaagac aggtcttcaa 900  
 agaagaagaa gaagctataa aggttgacct tcatgaagg tgtggtagaa ccaaattggt 960  
 ctggttgatg gcaactggcag attctaaaac aatgaaggca atggtggaat tccgagaaca 1020  
 cacaggcaaa cccaccacga gtagctcaga agcatgtcgc ttctgtggtt ccaggagtgg 1080  
 aacagagtta tctgctgttg gcagtgtttg ttctgatgca gattgccagg aatacgttaa 1140  
 gatagcctgt agtaagacgc atcctgtgtg ccatccatgc gggggtgtta aaaacgaaga 1200  
 gcactgtctg ccctgtctac acggctgtga caaaagtgc acaagcctga agcaagacgc 1260  
 cgatgacatg tgcagtatat gtttcaccga agcgctctcg gcagcaccag ccattcagct 1320  
 ggattgtagt cacatattcc acttacagtg ctgtcggcga gtattagaaa atcgatggct 1380  
 tggcccaagg ataactttg gattttatct ttgtccatt tgcaagaaca aaattaatca 1440  
 catagtacta aaagacctac ttgatccaat aaaagaactc tatgaggatg tcagaagaaa 1500  
 agccttaatg agattggaat atgaaggtct gcataagagt gaagctatca caactcctgg 1560  
 tgtgaggttt tataatgacc cagctggcta tgcaatgaat agatatgcat attatgtgtg 1620  
 ctacaaatgc agaaaggcat attttggtgg tgaagctcgc tgcgatgctg aggtctggacg 1680  
 ggggatgat tatgatccca gagagctcat ttgtggtgcc tgttctgatg tttccagggc 1740  
 tcagatgtgt cccaaacatg gcacagactt ttgggaatat aaatgtcgtc actgctgttc 1800  
 agtggctgtt tttttctgtt ttggaacaac acatttttgt aatgcttgtc atgatgattt 1860  
 tcaaagaatg actagcattc ctaaggaaga actaccacac tgtcctgcag gtcccaaagg 1920  
 caagcagtta gaaggaactg aatgtccact ccatgttgtt catccacca ctggggaaga 1980  
 gtttgcctct ggatgtggag tgtgcagaaa tgcccacact ttttagaaca cgcagatcct 2040  
 ttgtctacag agagaaaaat tgccttcac ccccaaggag atgctgtgaa gtttaaaactc 2100  
 tgctcaggat aaggacggga ccatttttac atccatgaaa atgaaccatt cacagtgcaa 2160  
 gaaggatacc aaataccatg tacataattc ttgctatgaa agttttcccc attatttttg 2220  
 tttatcttct tttgaacaaa tgacatcaaa cttgtgaggt gtttgcattg ggccattacc 2280  
 gtcattggcc tgtgaagcat tggacattta tagataattg atataaaaga atcgccatgc 2340  
 ccatggacta agaacgatgc tggctttcaa gcaaaaaaga aaaataatca ttgtttattg 2400  
 tatactgcct ttttgaatc ctgtacaatt gcacacggg tggggataaa aagaggaata 2460  
 ttctggttta tttcctagac tgttatttaa aaaaaaaaaa acattgtgtt aggacagcat 2520  
 ataaatgtaa taagtatcac actgtatata aacatatcaa tgtttgtcct gtataagaat 2580  
 tactaaatta caaatgcaat ttcattttaa cttctagggt aagtttgagc ctgaaatttt 2640  
 aatgaagtgc aatactgagt gtgcctcatt atcttgcagc tgtaaacata ttggaatgta 2700  
 catgtcaata aaaccactgt acatttttat acagtgataa agtctaaaaa aaaaaaaaaa 2760

<210> 18  
 <211> 660  
 <212> PRT  
 <213> Homo sapiens

<400> 18

```

Met Val Leu Ala Leu Ser Gly Ser Asn Val Gly Arg Gln Tyr Leu Ala
 1              5              10              15

Gln Gln Leu Thr Leu Leu Gln Asp Leu Phe Ser Leu Leu His Thr Ala
          20              25              30

Ser Pro Arg Val Gln Arg Gln Val Thr Ser Leu Leu Arg Arg Val Leu
          35              40              45

Pro Glu Val Thr Pro Ser Arg Leu Ala Ser Ile Ile Gly Val Lys Ser
 50              55              60

Leu Pro Pro Ala Asp Ile Ser Asp Ile Ile His Ser Thr Glu Lys Gly
 65              70              75              80

Asp Trp Asn Lys Leu Gly Ile Leu Asp Met Phe Leu Gly Cys Ile Ala
          85              90              95

Lys Ala Leu Thr Val Gln Leu Lys Ala Lys Gly Thr Thr Ile Thr Gly
          100              105              110

Thr Ala Gly Thr Thr Val Gly Lys Gly Val Thr Thr Val Thr Leu Pro
          115              120              125

Met Ile Phe Asn Ser Ser Tyr Leu Arg Arg Gly Glu Ser His Trp Trp
 130              135              140

Met Lys Gly Ser Thr Pro Thr Gln Ile Ser Glu Ile Ile Ile Lys Leu
 145              150              155              160

Ile Lys Asp Met Ala Ala Gly His Leu Ser Glu Ala Trp Ser Arg Val
          165              170              175

Thr Lys Asn Ala Ile Ala Glu Thr Ile Ile Ala Leu Thr Lys Met Glu
          180              185              190

Glu Glu Phe Arg Ser Pro Val Arg Cys Ile Ala Thr Thr Arg Leu Trp
          195              200              205

Leu Ala Leu Ala Ser Leu Cys Val Leu Asp Gln Asp His Val Asp Arg
 210              215              220

Leu Ser Ser Gly Arg Trp Met Gly Lys Asp Gly Gln Gln Lys Gln Met
 225              230              235              240

Pro Met Cys Asp Asn His Asp Asp Gly Glu Thr Ala Ala Ile Ile Leu
          245              250              255

Cys Asn Val Cys Gly Asn Leu Cys Thr Asp Cys Asp Arg Phe Leu His
          260              265              270

Leu His Arg Arg Thr Lys Thr His Gln Arg Gln Val Phe Lys Glu Glu
          275              280              285

Glu Glu Ala Ile Lys Val Asp Leu His Glu Gly Cys Gly Arg Thr Lys
          290              295              300

Leu Phe Trp Leu Met Ala Leu Ala Asp Ser Lys Thr Met Lys Ala Met
 305              310              315              320

```

Val Glu Phe Arg Glu His Thr Gly Lys Pro Thr Thr Ser Ser Ser Glu  
 325 330 335  
 Ala Cys Arg Phe Cys Gly Ser Arg Ser Gly Thr Glu Leu Ser Ala Val  
 340 345 350  
 Gly Ser Val Cys Ser Asp Ala Asp Cys Gln Glu Tyr Ala Lys Ile Ala  
 355 360 365  
 Cys Ser Lys Thr His Pro Cys Gly His Pro Cys Gly Gly Val Lys Asn  
 370 375 380  
 Glu Glu His Cys Leu Pro Cys Leu His Gly Cys Asp Lys Ser Ala Thr  
 385 390 395 400  
 Ser Leu Lys Gln Asp Ala Asp Asp Met Cys Met Ile Cys Phe Thr Glu  
 405 410 415  
 Ala Leu Ser Ala Ala Pro Ala Ile Gln Leu Asp Cys Ser His Ile Phe  
 420 425 430  
 His Leu Gln Cys Cys Arg Arg Val Leu Glu Asn Arg Trp Leu Gly Pro  
 435 440 445  
 Arg Ile Thr Phe Gly Phe Ile Ser Cys Pro Ile Cys Lys Asn Lys Ile  
 450 455 460  
 Asn His Ile Val Leu Lys Asp Leu Leu Asp Pro Ile Lys Glu Leu Tyr  
 465 470 475 480  
 Glu Asp Val Arg Arg Lys Ala Leu Met Arg Leu Glu Tyr Glu Gly Leu  
 485 490 495  
 His Lys Ser Glu Ala Ile Thr Thr Pro Gly Val Arg Phe Tyr Asn Asp  
 500 505 510  
 Pro Ala Gly Tyr Ala Met Asn Arg Tyr Ala Tyr Tyr Val Cys Tyr Lys  
 515 520 525  
 Cys Arg Lys Ala Tyr Phe Gly Gly Glu Ala Arg Cys Asp Ala Glu Ala  
 530 535 540  
 Gly Arg Gly Asp Asp Tyr Asp Pro Arg Glu Leu Ile Cys Gly Ala Cys  
 545 550 555 560  
 Ser Asp Val Ser Arg Ala Gln Met Cys Pro Lys His Gly Thr Asp Phe  
 565 570 575  
 Leu Glu Tyr Lys Cys Arg Tyr Cys Cys Ser Val Ala Val Phe Phe Cys  
 580 585 590  
 Phe Gly Thr Thr His Phe Cys Asn Ala Cys His Asp Asp Phe Gln Arg  
 595 600 605  
 Met Thr Ser Ile Pro Lys Glu Glu Leu Pro His Cys Pro Ala Gly Pro  
 610 615 620  
 Lys Gly Lys Gln Leu Glu Gly Thr Glu Cys Pro Leu His Val Val His  
 625 630 635 640

Pro Pro Thr Gly Glu Glu Phe Ala Leu Gly Cys Gly Val Cys Arg Asn  
 645 650 655

Ala His Thr Phe  
 660

<210> 19  
 <211> 1649  
 <212> DNA  
 <213> Homo sapiens

<400> 19  
 gattgtacat agtcttgtgg ggcattggggg agccggctgg aggtgagaac cctccccctct 60  
 cccccacccc cccggggaga gcaaatgtaa aactactaat ttttgtgctt tatatattct 120  
 atataaatat atctattttc tttttacaaa accagtttat aaatggtagg ggggtgtggg 180  
 gcggacacat ggagctcccc ttgtgggggg gccccctcca ttacccgacc taccgcccct 240  
 ttcttcacccc cccaccccac tccccacccc ctggctgtga ctgctgtaag atgggggtat 300  
 agaggtctgg caattcccac cccctgttgt atagttggac tatgttataa cgcacaaaag 360  
 agagctgacc ccagggggag ccagaggggtg atgggttctt tgcctcccct tccttcccct 420  
 ttttgcctaa gcttgtgctg cagttgaacc tcttctctgg ggtgggagta ggtaaggggt 480  
 ggggtgaggg ccaaacccct ctctggtagg gaaccgtggg gatgaagatg aagcttatat 540  
 gcagttctct tctaggggct gtgggcaaa ggcattttgt aattaatatt ttcaagaatc 600  
 agatgtctgg agtgtagggg tgggcttggg ggtggtggac gggcgggcct gctggagggg 660  
 gagcttggtc gctgttgtga ttttaggttt gtttttgttt tgttttgaat ttgggggggt 720  
 gtggattgtt gggggttaggg agattttttt ttttttaaag ctgcttcctc aactgtttca 780  
 agctgcaaat gtttaagaga ataacagccc ccactcccac aggaaccgct gtaattaaat 840  
 cagacagtag gaagactggg ctgctgccct caaagccaca gcccttgat gttccttttc 900  
 cgagagcaga aggtctaggc tacagggagg gggagattgg ctcccgtgag tcaggctgtg 960  
 tttggggcct gggccctggg attgggaaaa ggggatgggg cagactttgt aagcatatgc 1020  
 taggtatccg atagtcctgt agaatttagt gaagaaacct tatacagttt ttaattttta 1080  
 tataaactat aactcagacc caagctacaa ggttggaatt ttggttggtt ttttttttaa 1140  
 gtaccctgcc tgtataattg catcagaatc cccaccccca ccccsgcc csgtgtttgt 1200  
 attttgggtt ggtttacact cgcacatact cagttttcag ttttcccctt tacagtcttc 1260  
 tcccctcacc tccaggaccc tccccctttt taaaaataa atcgtgaca agtgtgaatc 1320  
 ccgtgaagac tttattttgt gttgtgtgta tctgttacag caaggttggt ccttcgtaac 1380  
 aacggatgaa atggttccct tttttaaaag gccctctctc cctccaccct cagcgcccct 1440  
 gtccttggca tgttttgtat cagcgatcat tctgaactgt acataattat gttgcgagag 1500  
 gcaaagggca agttttggat tttgttctt ccaagtttgt ttttaaacga caaataaaaa 1560  
 aagaacattt taataaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1620  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1649

<210> 20  
 <211> 92  
 <212> PRT  
 <213> Homo sapiens

<400> 20  
 Met Gly Glu Pro Ala Gly Gly Glu Asn Pro Pro Leu Ser Pro His Pro  
 1 5 10 15  
 Pro Gly Arg Ala Asn Val Lys Leu Leu Ile Phe Val Leu Tyr Ile Phe  
 20 25 30  
 Tyr Ile Asn Ile Ser Ile Phe Phe Leu Gln Asn Gln Phe Ile Asn Gly  
 35 40 45  
 Arg Gly Val Trp Gly Gly His Met Glu Leu Pro Leu Trp Gly Gly Pro  
 50 55 60

Leu His Tyr Pro Thr Tyr Arg Pro Phe Pro His Pro Pro Pro His Ser  
 65 70 75 80

Pro Pro Pro Gly Cys Asp Cys Cys Lys Met Gly Val  
 85 90

<210> 21  
 <211> 2644  
 <212> DNA  
 <213> Homo sapiens

<400> 21  
 gttgaggatg gctgacattc tctctcagtc agagaccctg gcgtcgcaag acctcagtgg 60  
 ggacttcaag aagccagctc tgccggtgtc cccagcggcg cggagtaagg ccccgccag 120  
 cagttcttca aaccctgagg aggtacagaa ggaagggccc actgcgttgc aggactccaa 180  
 ttctggggag cccgacatcc ctctctctca gccggactgc ggtgatttta ggagtctaca 240  
 ggaggagcag tcgcgccccca cgacagcggg ttcttcccct ggcggtccag cccgggctcc 300  
 cccctaccaa gagcctccat ggggtggccc tgccacagcc ccctacagct tagagacct 360  
 gaagggcggc actatccttg gcacccgtag cttgaaaggg acgagttact gccttttcgg 420  
 gaggctgtct ggctgcgacg tgtgcctgga gcaccttcg gtgtctcgg accacgcagt 480  
 gctgcagcac agggcgctccg gccctgacgg agaatgcgac agcaacgggc cgggcttcta 540  
 cctctacgat ctgggaagca cccatggcac tttctcaac aaaactcgca tcccacctcg 600  
 cacctactgt cgagtccacg ttgggcatgt tgttcgcttt ggaggcagca cccggctctt 660  
 tctctgcag ggaccagagg aagaccgaga ggcagaatcc gagttaacag taacacagtt 720  
 gaaggaattg cgcaagcagc agcaaatatt gttggrgaag aagatgctag gagaagactc 780  
 agatgaagaa gaggaatgg atacctctga aaggaagata aatgctggt gccaagatga 840  
 tgagatgggt tgcacctggg gaatgggaga agatgcagta gaggatgatg ctgaagagaa 900  
 ccctattgtc ttagagtttc agcaggaaag ggaggccttt tatataaagg atccccaaaa 960  
 ggctctccaa ggcttttttt accgagaagg agaagaatta gaatatgaat ttgatgaaca 1020  
 gggacatagc acttggtctc gcaggggtgag attacctgtg gacgattcaa ctggaaaaca 1080  
 actggtggct gaggccattc actcaggaaa gaaaaaagaa gcaatgatcc agtgctcatt 1140  
 ggaagcttgt cggattcttg acactttggg attgcttcgg caggaagcag tatctcggaa 1200  
 aaggaagcc aagaactggg aagatgaaga cttttatgat agtgatgatg acacatttct 1260  
 tgataggact ggcctgattg agaagaagcg tctgaacaga atgaagaagg ctggcaagat 1320  
 tgatgagaag ccagagacct ttgaatcatt ctcaagccaa gttctatcag agtctccat 1440  
 actttctgaa atttctgaga gattgaaagc aatgaaatca ggcagtacat tagatggtgt 1500  
 tcaggattct ttagatgcgt tcatgtcaga aatgaaatca ggcagtacat tagatggtgt 1500  
 gtcccgaag aaacttcacc tgagaacttt tgaactgagg aaagaacaac agagacttaa 1560  
 agggttaata aaaattgtaa agccagcaga gattccagaa ctaaaaaaga ctgaaactca 1620  
 gactacaggt gcagaaaaca aagctaaaaa gcttacattg cctctatttg gtgccatgaa 1680  
 aggaggaagc aaattcaaat taaaaactgg aacagtaggg aagttacccc ccaagcgtcc 1740  
 agaactccct ccaactctaa tgagaatgaa agatgagcct gaagtagaag aggaggagga 1800  
 agaggaagag gaagaagaga aagaaaagga ggagcatgaa aagaaaaaac tggaggatgg 1860  
 aagcctcagt aggccacagc cagagataga gccagaagca gcagtgcagg aaatgaggcc 1920  
 tcccacagat ctacacatt ttaaagaaac ccaaaccat ggtaatatct ttcttctcct 1980  
 tctgtgttg ttcagtgggc agttacattg attgtggata ggttttaaaa agcaaggcca 2040  
 gttcttgtct gtgcatttga ctttgtatgt gatatactga ctctgtagca aggaaacata 2100  
 ctttcttggg cttcttcctt tgaccgccag tcattatttg tcttcattgc aaattaaggg 2160  
 cagttatttc caatccattc cagaattaca gaaaattgaa gggctatgga atctgaaacc 2220  
 atagctgctg tggataatc ctgagctgct gccactgtgt gagggtggag gcagtggaa 2280  
 agggtagatg atggggcctg atcaggtggt ctccggataa gtcaaccctt attcattttt 2340  
 tctccatcc ctaaaacaga ggccaaacca taattgtact cattggacta aagttctcaa 2400  
 gaaggatctt gcttcattca tttttgtgtg tttggaacct agcacaaaac ctgacacata 2460  
 tccacccgcc tcagcaaata tttgatgaaa aatgttgaaa gacggaatag attgatattc 2520  
 atatagatat atgcatacat taattctgta ttttctatat atatattcta attacaaagg 2580  
 gttatatggt cattttagaa actatagatc atacataaaa gtccaaagga aaaaaaaaaa 2640  
 aaaa 2644

<210> 22

<211> 667  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (250)

<400> 22

```

Met Ala Asp Ile Leu Ser Gln Ser Glu Thr Leu Ala Ser Gln Asp Leu
 1              5              10              15

Ser Gly Asp Phe Lys Lys Pro Ala Leu Pro Val Ser Pro Ala Ala Arg
      20              25              30

Ser Lys Ala Pro Ala Ser Ser Ser Ser Asn Pro Glu Glu Val Gln Lys
      35              40              45

Glu Gly Pro Thr Ala Leu Gln Asp Ser Asn Ser Gly Glu Pro Asp Ile
      50              55              60

Pro Pro Pro Gln Pro Asp Cys Gly Asp Phe Arg Ser Leu Gln Glu Glu
      65              70              75              80

Gln Ser Arg Pro Thr Thr Ala Val Ser Ser Pro Gly Gly Pro Ala Arg
      85              90              95

Ala Pro Pro Tyr Gln Glu Pro Pro Trp Gly Gly Pro Ala Thr Ala Pro
      100             105             110

Tyr Ser Leu Glu Thr Leu Lys Gly Gly Thr Ile Leu Gly Thr Arg Ser
      115             120             125

Leu Lys Gly Thr Ser Tyr Cys Leu Phe Gly Arg Leu Ser Gly Cys Asp
      130             135             140

Val Cys Leu Glu His Pro Ser Val Ser Arg Tyr His Ala Val Leu Gln
      145             150             155             160

His Arg Ala Ser Gly Pro Asp Gly Glu Cys Asp Ser Asn Gly Pro Gly
      165             170             175

Phe Tyr Leu Tyr Asp Leu Gly Ser Thr His Gly Thr Phe Leu Asn Lys
      180             185             190

Thr Arg Ile Pro Pro Arg Thr Tyr Cys Arg Val His Val Gly His Val
      195             200             205

Val Arg Phe Gly Gly Ser Thr Arg Leu Phe Ile Leu Gln Gly Pro Glu
      210             215             220

Glu Asp Arg Glu Ala Glu Ser Glu Leu Thr Val Thr Gln Leu Lys Glu
      225             230             235             240

Leu Arg Lys Gln Gln Gln Ile Leu Leu Xaa Lys Lys Met Leu Gly Glu
      245             250             255

Asp Ser Asp Glu Glu Glu Glu Met Asp Thr Ser Glu Arg Lys Ile Asn
      260             265             270
  
```

Ala Gly Ser Gln Asp Asp Glu Met Gly Cys Thr Trp Gly Met Gly Glu  
 275 280 285  
 Asp Ala Val Glu Asp Asp Ala Glu Glu Asn Pro Ile Val Leu Glu Phe  
 290 295 300  
 Gln Gln Glu Arg Glu Ala Phe Tyr Ile Lys Asp Pro Lys Lys Ala Leu  
 305 310 315 320  
 Gln Gly Phe Phe Asp Arg Glu Gly Glu Glu Leu Glu Tyr Glu Phe Asp  
 325 330 335  
 Glu Gln Gly His Ser Thr Trp Leu Cys Arg Val Arg Leu Pro Val Asp  
 340 345 350  
 Asp Ser Thr Gly Lys Gln Leu Val Ala Glu Ala Ile His Ser Gly Lys  
 355 360 365  
 Lys Lys Glu Ala Met Ile Gln Cys Ser Leu Glu Ala Cys Arg Ile Leu  
 370 375 380  
 Asp Thr Leu Gly Leu Leu Arg Gln Glu Ala Val Ser Arg Lys Arg Lys  
 385 390 395 400  
 Ala Lys Asn Trp Glu Asp Glu Asp Phe Tyr Asp Ser Asp Asp Asp Thr  
 405 410 415  
 Phe Leu Asp Arg Thr Gly Leu Ile Glu Lys Lys Arg Leu Asn Arg Met  
 420 425 430  
 Lys Lys Ala Gly Lys Ile Asp Glu Lys Pro Glu Thr Phe Glu Ser Leu  
 435 440 445  
 Val Ala Lys Leu Asn Asp Ala Glu Arg Glu Leu Ser Glu Ile Ser Glu  
 450 455 460  
 Arg Leu Lys Ala Ser Ser Gln Val Leu Ser Glu Ser Pro Ser Gln Asp  
 465 470 475 480  
 Ser Leu Asp Ala Phe Met Ser Glu Met Lys Ser Gly Ser Thr Leu Asp  
 485 490 495  
 Gly Val Ser Arg Lys Lys Leu His Leu Arg Thr Phe Glu Leu Arg Lys  
 500 505 510  
 Glu Gln Gln Arg Leu Lys Gly Leu Ile Lys Ile Val Lys Pro Ala Glu  
 515 520 525  
 Ile Pro Glu Leu Lys Lys Thr Glu Thr Gln Thr Thr Gly Ala Glu Asn  
 530 535 540  
 Lys Ala Lys Lys Leu Thr Leu Pro Leu Phe Gly Ala Met Lys Gly Gly  
 545 550 555 560  
 Ser Lys Phe Lys Leu Lys Thr Gly Thr Val Gly Lys Leu Pro Pro Lys  
 565 570 575  
 Arg Pro Glu Leu Pro Pro Thr Leu Met Arg Met Lys Asp Glu Pro Glu  
 580 585 590



Val Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu Lys Glu Lys Glu  
 595 600 605

Glu His Glu Lys Lys Lys Leu Glu Asp Gly Ser Leu Ser Arg Pro Gln  
 610 615 620

Pro Glu Ile Glu Pro Glu Ala Ala Val Gln Glu Met Arg Pro Pro Thr  
 625 630 635 640

Asp Leu Thr His Phe Lys Glu Thr Gln Thr His Gly Asn Ile Phe Leu  
 645 650 655

Leu Leu Pro Val Leu Phe Ser Gly Gln Leu His  
 660 665

&lt;210&gt; 23

&lt;211&gt; 2402

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

gatcgagag accaaggagg cgcccgcggc tgcagagctg cagagcgagg tctcttcgag 60  
 ctgtctgtgt ccgggcagcc ggcgcgcaac tgagccagag gacagcgcat cctttcggcg 120  
 cgggcccggca gggcccctgc ggctcggaag ctggctcccc ggggtggccac cgggaccccc 180  
 gagcccaatg gcgggggagg cggaacaaatc cacaacactg tagagatcac cccacacctc 240  
 aacggacagg tcgggaccct cggagatgag gtgcccacgg agcagctgca gggtagagcg 300  
 gagcgcgagc gggaggggga gggagacgag ggcggcgagc gactgggagc cagcctgtcg 360  
 ctggccgtgc cccagggccc cctcagcttt agtgctctgc cggtgctctt cgtggctctg 420  
 ggcggcgagg agcagctgca gctcgccctc tgctgctctg cggtgctctt cgtggctctg 480  
 ggcattggcct cggaccccat cttcacgctg gcgccccgc tgcattgcca ctacggggcc 540  
 tcccccccta atgctctgag ctgggagcag cctcccaatg ccagcggcgt cagcgtcgcc 600  
 agcgtgccc tagcagccag gcgcgcccag cgtgtcgcca ccagtagcga cccctcgtgc 660  
 agcggcttcg ccccgccgga cttcaacatg tgctcaagg attgggacta taatggcctt 720  
 cctgtgctca ccaccaacgc catcgccagc tgggagctgt tgtgtgacct gggctggcag 780  
 gtgatccagg acagatctt cttcatcttg ggctttgctt ccggctacct gttcctgggt 840  
 taccgagcag acagatttgg ccgtcgcgag attgtgctgc tgaccttggg gctgggtggg 900  
 cctgtggag taggaggggc tgctgcaggc tctccacag gcgtcatggc cctccgattc 960  
 ctcttgggct ttctgctgag cgggtgtgac ctgggtgtct acctgatgag cctggagctg 1020  
 tgcgacccaa cccagaggct tcgggtggcc ctggcagggg agttgggtgg ggtgggaggg 1080  
 cacttctgtg tcttgggcct ggcccttgct tctaaggatt ggcgattcct acagcgaatg 1140  
 atcacgctc cctgcatcct cttcctgttt tatggctggc ctggtttgtt cctggagtc 1200  
 gcacggtggc tgatagtga ggcgcagatt gaggaggctc agtctgtgct gaggatcctg 1260  
 gctgagcgaa accggcccca tgggcagatg ctgggggagg agggccagga ggccctgcag 1320  
 gacctggaga atacctgccc tctccctgca acatcctcct tttcctttgc tccctcctc 1380  
 aactaccgca acatctggaa aaatctgctt atcctgggct tcaccaactt cattgccc 1440  
 gccattcgcc actgctacca gcctgtggga ggaggaggga gcccatcgga cttctacctg 1500  
 tgctctctgc tggccagcgg caccgcagcc ctggcctgtg tcttctctgg ggtcaccgtg 1560  
 gaccgatttg gccgcgggg catccttctt cttccatga ccttaccgg cattgcttcc 1620  
 ctggctctgc tgggcctgtg ggattatctg aacgaggctg ccatcaccac tttctctgtc 1680  
 cttggctctt tctcctcca agctgcccgc atcctcagca cctccttgc tgcagagtc 1740  
 atccccacca ctgtccgggg ccgtggcctg ggctgatga tggctctagg ggcgcttggg 1800  
 ggactgagcg gcccgccca gcgcctccac atggggccat gagccttctt gcagcacgtg 1860  
 gtgctggcgg cctgcgccct cctctgcatt ctcagcatta tgctgctgcc ggagaccaag 1920  
 cgcaagctcc tgcccagagt gtcgggggac ggggagctgt gtcgcccggc ttcctctgtg 1980  
 cggcagccac cccctaccg ctgtgaccac gtcccgtgc ttgccacccc caaccctgcc 2040  
 ctctgagcgg cctctgagta ccctggcggg aggctggccc acacagaaag gtggcaagaa 2100  
 gatcggaag actgagtagg gaaggcaggg ctgcccagaa gtctcagagg cacctcacgc 2160  
 cagccatcgc ggagagctca gaggccgctc cccaccctgc ctctccctg ctgctttgca 2220  
 ttcacttctt tggccagagt caggggacag ggagagagct ccacactgta accactgggt 2280

ctgggctcca tcttgccccc aaagacatcc acccagacct cattatttct tgctctatca 2340  
 ttctgtttca ataaagacat ttggaataaa cgaaaaaaaa aaaaaaaaaa 2400  
 aa 2402

<210> 24

<211> 520

<212> PRT

<213> Homo sapiens

<400> 24

Met Ala Ser Asp Pro Ile Phe Thr Leu Ala Pro Pro Leu His Cys His  
 1 5 10 15  
 Tyr Gly Ala Phe Pro Pro Asn Ala Ser Gly Trp Glu Gln Pro Pro Asn  
 20 25 30  
 Ala Ser Gly Val Ser Val Ala Ser Ala Ala Leu Ala Ala Ser Ala Ala  
 35 40 45  
 Ser Arg Val Ala Thr Ser Thr Asp Pro Ser Cys Ser Gly Phe Ala Pro  
 50 55 60  
 Pro Asp Phe Asn His Cys Leu Lys Asp Trp Asp Tyr Asn Gly Leu Pro  
 65 70 75 80  
 Val Leu Thr Thr Asn Ala Ile Gly Gln Trp Asp Leu Val Cys Asp Leu  
 85 90 95  
 Gly Trp Gln Val Ile Leu Glu Gln Ile Leu Phe Ile Leu Gly Phe Ala  
 100 105 110  
 Ser Gly Tyr Leu Phe Leu Gly Tyr Pro Ala Asp Arg Phe Gly Arg Arg  
 115 120 125  
 Gly Ile Val Leu Leu Thr Leu Gly Leu Val Gly Pro Cys Gly Val Gly  
 130 135 140  
 Gly Ala Ala Ala Gly Ser Ser Thr Gly Val Met Ala Leu Arg Phe Leu  
 145 150 155 160  
 Leu Gly Phe Leu Leu Ala Gly Val Asp Leu Gly Val Tyr Leu Met Arg  
 165 170 175  
 Leu Glu Leu Cys Asp Pro Thr Gln Arg Leu Arg Val Ala Leu Ala Gly  
 180 185 190  
 Glu Leu Val Gly Val Gly Gly His Phe Leu Phe Leu Gly Leu Ala Leu  
 195 200 205  
 Val Ser Lys Asp Trp Arg Phe Leu Gln Arg Met Ile Thr Ala Pro Cys  
 210 215 220  
 Ile Leu Phe Leu Phe Tyr Gly Trp Pro Gly Leu Phe Leu Glu Ser Ala  
 225 230 235 240  
 Arg Trp Leu Ile Val Lys Arg Gln Ile Glu Glu Ala Gln Ser Val Leu  
 245 250 255  
 Arg Ile Leu Ala Glu Arg Asn Arg Pro His Gly Gln Met Leu Gly Glu  
 260 265 270

Glu Ala Gln Glu Ala Leu Gln Asp Leu Glu Asn Thr Cys Pro Leu Pro  
 275 280 285  
 Ala Thr Ser Ser Phe Ser Phe Ala Ser Leu Leu Asn Tyr Arg Asn Ile  
 290 295 300  
 Trp Lys Asn Leu Leu Ile Leu Gly Phe Thr Asn Phe Ile Ala His Ala  
 305 310 315 320  
 Ile Arg His Cys Tyr Gln Pro Val Gly Gly Gly Gly Ser Pro Ser Asp  
 325 330 335  
 Phe Tyr Leu Cys Ser Leu Leu Ala Ser Gly Thr Ala Ala Leu Ala Cys  
 340 345 350  
 Val Phe Leu Gly Val Thr Val Asp Arg Phe Gly Arg Arg Gly Ile Leu  
 355 360 365  
 Leu Leu Ser Met Thr Leu Thr Gly Ile Ala Ser Leu Val Leu Leu Gly  
 370 375 380  
 Leu Trp Asp Tyr Leu Asn Glu Ala Ala Ile Thr Thr Phe Ser Val Leu  
 385 390 395 400  
 Gly Leu Phe Ser Ser Gln Ala Ala Ala Ile Leu Ser Thr Leu Leu Ala  
 405 410 415  
 Ala Glu Val Ile Pro Thr Thr Val Arg Gly Arg Gly Leu Gly Leu Ile  
 420 425 430  
 Met Ala Leu Gly Ala Leu Gly Gly Leu Ser Gly Pro Ala Gln Arg Leu  
 435 440 445  
 His Met Gly His Gly Ala Phe Leu Gln His Val Val Leu Ala Ala Cys  
 450 455 460  
 Ala Leu Leu Cys Ile Leu Ser Ile Met Leu Leu Pro Glu Thr Lys Arg  
 465 470 475 480  
 Lys Leu Leu Pro Glu Val Leu Arg Asp Gly Glu Leu Cys Arg Arg Pro  
 485 490 495  
 Ser Leu Leu Arg Gln Pro Pro Pro Thr Arg Cys Asp His Val Pro Leu  
 500 505 510  
 Leu Ala Thr Pro Asn Pro Ala Leu  
 515 520

&lt;210&gt; 25

&lt;211&gt; 2377

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

ttcattcttc agtggaatc catcagttga aatagttcat ggtattatgc acctatataa 60  
 gacaaataag atgacctcct taaaagaaga tgtgcggcgc agtgccatgc tgtgtattct 120  
 cacagtcctt gctgcaatga ccagtcacga ccttatgaag tttgttgccc catttaacga 180  
 agtaattgaa caaatgaaaa ttatcagaga ctctactccc aaccaatata tgggtgctgat 240

```

aaggtttcgt gcacaggctg atgcggatag tttttatatg acatgcaatg gccgccagtt 300
caactcaata gaagatgacg tttgccagct agtgtatgtg gaaagagctg aagtgtctaa 360
atctgaagat ggcgccagcc tcccagtgat ggacctgact gaactcccca agtgcacggg 420
gtgtctggag cgcattggacg agtctgtgaa tggcatcctc acaacgttat gtaaccacag 480
cttccacagc cagtgtctac agcgtctggga cgataccacg tgtcctgttt gccgggtactg 540
tcaaacgccc gagccagtag aagaaaaataa gtgtttttgag tgtggtgttc aggaaaaatct 600
ttggattttgt ttaatatgcg gccacatagg atgtggacgg tatgtcagtc gacatgctta 660
taagcacttt gaggaaacgc agcacacgta tgccatgcag cttaccaacc atcgagtctg 720
ggactatgct ggagataact atgttcatcg actggttgca agtaaaacag atggaaaaat 780
agtacagtat gaatgtgagg gggatacttg ccaggaagag aaaatagatg ccttacagtt 840
agagtattca tatttactaa caagccagct ggaatctcag cgaatctact gggaaaaacaa 900
gatagtctcg atagagaagg acacagcaga ggaaattaac aacatgaaga ccaagtttaa 960
agaaacaatt gagaagtgtg ataattctaga gcacaaacta aatgatctcc taaaagaaaa 1020
gcagtctgtg gaaagaaaag gcactcagct aaacacaaaa gtggccaaac tcaaattctca 1080
gagtgggtat cctagcatct agcaagactg agtggggaga tttctcatcc gtgtgaaaaat 1140
gtagagtgtg gcctctgact agctaattgt gtattttgtt gggtttagta ttttctaaat 1200
gtttacaaaa tattgggctg catgttcagg ttgcagctag agggagcttg ggcagatttt 1260
caattacgct ttcaagatat aaccaaaagc tgtttctaaa tcctaaaatt agaatttcaa 1320
cagagccccc tttagaacag tcatataacg cttgtgtggg ccaacagagg ggctgtgtac 1380
tctctctgga accataaatg tcaaataatt tataacctgc agtaattgag caaacttaaa 1440
ataagacctg tggttgaatt tagtttcttg aagaggtaga gggatagggt agtaagatgt 1500
attgttaaac aacaggtttt agtttttctg ttataattag ccacagggtt tcaaattgat 1560
acatttcaga atagggtttt agcctgtaat taggcctcat cccctttgac cttaatgtct 1620
gacatgttac ttgttagcac atcaactgta tcaactaatca ccatctgttt ttgtgggag 1680
tgctgcagca tttcccaaaa aactttacgt gtaattgttc aaaatgaatg tactcagaca 1740
ttcttaattt ttacttaggg cagaccaact ctttgagtct ctcttggaact tatatataca 1800
gatattctaa gagtgggaat gtaaagcata acctaatctt ctttctctata gagattctat 1860
tttattttaa atctattttt acactagtta gaatcctgct gttttggcca agtacttgct 1920
ttgcatgtct caccttgtag aagctggggg ggatcatagc atactaatga agagaattag 1980
aagtagttta caaagctcgc tcaactccta tttctctgtg atcccttcta tccagtggcc 2040
ccaccaccac ctgggaaaac agattttttca gtacagggtg gataaatgct ttgaaaggct 2100
gtgcccagag caatgagcaa ataggcaagt gtttccaaac tamttggagg tttacaaaaa 2160
atatgtccca gaaaaaaaaa aaatcttacc aagatacgt aagaaaaaaaa aatttttttt 2220
taaacagtca aagagtcatt tttgaatttc acaaaatcac atcagacaga agttgttttc 2280
ttcaggaggg aaatgaacca cttaatatat ccatactacc ttgaacaatg aaattgaatt 2340
aaaaatagcca aactttgaaa ttaaaaaaaaa aaaaaaa 2377

```

&lt;210&gt; 26

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

```

Met His Leu Tyr Lys Thr Asn Lys Met Thr Ser Leu Lys Glu Asp Val
  1                      5                      10          15
Arg Arg Ser Ala Met Leu Cys Ile Leu Thr Val Pro Ala Ala Met Thr
          20                      25          30
Ser His Asp Leu Met Lys Phe Val Ala Pro Phe Asn Glu Val Ile Glu
          35                      40          45
Gln Met Lys Ile Ile Arg Asp Ser Thr Pro Asn Gln Tyr Met Val Leu
          50                      55          60
Ile Lys Phe Arg Ala Gln Ala Asp Ala Asp Ser Phe Tyr Met Thr Cys
          65                      70          75          80
Asn Gly Arg Gln Phe Asn Ser Ile Glu Asp Asp Val Cys Gln Leu Val
          85                      90          95

```

Tyr Val Glu Arg Ala Glu Val Leu Lys Ser Glu Asp Gly Ala Ser Leu  
 100 105 110  
 Pro Val Met Asp Leu Thr Glu Leu Pro Lys Cys Thr Val Cys Leu Glu  
 115 120 125  
 Arg Met Asp Glu Ser Val Asn Gly Ile Leu Thr Thr Leu Cys Asn His  
 130 135 140  
 Ser Phe His Ser Gln Cys Leu Gln Arg Trp Asp Asp Thr Thr Cys Pro  
 145 150 155 160  
 Val Cys Arg Tyr Cys Gln Thr Pro Glu Pro Val Glu Glu Asn Lys Cys  
 165 170 175  
 Phe Glu Cys Gly Val Gln Glu Asn Leu Trp Ile Cys Leu Ile Cys Gly  
 180 185 190  
 His Ile Gly Cys Gly Arg Tyr Val Ser Arg His Ala Tyr Lys His Phe  
 195 200 205  
 Glu Glu Thr Gln His Thr Tyr Ala Met Gln Leu Thr Asn His Arg Val  
 210 215 220  
 Trp Asp Tyr Ala Gly Asp Asn Tyr Val His Arg Leu Val Ala Ser Lys  
 225 230 235 240  
 Thr Asp Gly Lys Ile Val Gln Tyr Glu Cys Glu Gly Asp Thr Cys Gln  
 245 250 255  
 Glu Glu Lys Ile Asp Ala Leu Gln Leu Glu Tyr Ser Tyr Leu Leu Thr  
 260 265 270  
 Ser Gln Leu Glu Ser Gln Arg Ile Tyr Trp Glu Asn Lys Ile Val Arg  
 275 280 285  
 Ile Glu Lys Asp Thr Ala Glu Glu Ile Asn Asn Met Lys Thr Lys Phe  
 290 295 300  
 Lys Glu Thr Ile Glu Lys Cys Asp Asn Leu Glu His Lys Leu Asn Asp  
 305 310 315 320  
 Leu Leu Lys Glu Lys Gln Ser Val Glu Arg Lys Cys Thr Gln Leu Asn  
 325 330 335  
 Thr Lys Val Ala Lys Leu Lys Ser Gln Ser Gly Tyr Pro Ser Ile  
 340 345 350

&lt;210&gt; 27

&lt;211&gt; 460

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

cgagatgaag cccggcgggtgg acgagatggt ccccgagggc gccggggccct acgtggacct 60  
 ggacgaggcg ggaggcagca ccgggctctt gatggacttg gcagccaatg aaaagccgtt 120  
 catgcagact tttttaacga ttttgaagat ctttttgatg atgatgacat ccagtgaatg 180  
 gccctctggc tgcaggcggg gcccaagccct tggtagagag ccgcagtgtg agcctgcgca 240

```

ggacagtttc aggtggtttt aaagaacacg tggaaatccc ttgaatttag gacctgggta 300
accagaaaga taagactggt cttaacgacc tagatgattc tgttcacatc tgaacgggat 360
cagggttttgt cctcactcca attaaaagaa agcaatgtca catgaaaaaa aaaaaaaaaa 420
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa

```

460

&lt;210&gt; 28

&lt;211&gt; 85

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

```

Met Lys Pro Ala Val Asp Glu Met Phe Pro Glu Gly Ala Gly Pro Tyr
  1             5             10             15

```

```

Val Asp Leu Asp Glu Ala Gly Gly Ser Thr Gly Leu Leu Met Asp Leu
          20             25             30

```

```

Ala Ala Asn Glu Lys Pro Phe Met Gln Thr Phe Leu Thr Ile Leu Lys
      35             40             45

```

```

Ile Phe Leu Met Met Met Thr Ser Ser Glu Met Pro Ser Gly Cys Arg
      50             55             60

```

```

Arg Gly Gln Ala Leu Gly Thr Glu Pro Gln Cys Glu Pro Ala Gln Asp
      65             70             75             80

```

```

Ser Phe Arg Trp Phe
          85

```

&lt;210&gt; 29

&lt;211&gt; 3204

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 29

```

gtttggcatc tgtggccgag ttgctgttgc cgggtgatag ttggagcggg gacttagcac 60
aatggcagaa cctgtttctc cactgaagca ctttgtgctg gctaagaagg cgattactgc 120
aatctttgac cagttactgg agtttggttac tgaaggatca cattttgttg aagcaacata 180
taagaatccg gaacttgatc gaatagccac tgaagatgat ctggtagaaa tgcaaggata 240
taaagacaag ctttccatca ttggtgaggt gctatctcgg agacacatga aggtggcatt 300
ttttggcagg acaagcagtg ggaagagctc tgttatcaat gcaatgttgt gggataaagt 360
tctccctagt gggattggcc atataaccaa ttgcttcceta agtggtgaag gaactgatgg 420
agataaagcc tatcttatga cagaaggatc agatgaaaaa aagagtgtga agacagttaa 480
tcaactggcc catgcccttc acatggacaa agatttgaaa gctggctgtc ttgtacgtgt 540
gttttgacca aaagcaaaat gtgccctctt gagagatgac ctggtgttag tagacagtcc 600
aggcacagat gtcactacag agctggatag ctggattgat aagttttgcc tagatgctga 660
tgtctttgtt ttggtcgcaa actctgaatc aacactaatg aatacggaaa aacacttttt 720
tcacaagggtg aatgagcggc tttccaagcc taatattttc attctcaata atcgttggga 780
tgctcttgca tcagagccag aatatatgga agacgtacgc agacagcaca tggaaagatg 840
cctgatttcc ttggtggagg agctcaaagt tgtaaatgct ttagaagcac agaatcgtat 900
cttctttgtt tcagcaaagg aagttcttag tgctagaaag caaaaagcac aggggatgcc 960
agaaagtggg gtggcacttg ctgaaggatt tcatgcaaga ttacaggaat ttcagaattt 1020
tgaacaaatc tttgaggagt gtatctcgca gtcagcagtg aaaacaaagt tcgaacagca 1080
cactatcaga gctaaacaga tactagctac tgtgaaaaac ataatggatt cagtaaacct 1140
ggcagctgaa gataaaaggc attattcagt ggaagagagg gaagaccaa ttgatagact 1200
ggactttatt cgaaaccaga tgaacctttt aacactggat gttaagaaaa aaatcaagga 1260
ggttaccgag gaggtggcaa acaaagtttc atgtgcaatg acagatgaaa tttgtcgact 1320
gtctgttttg gttgatgaat tttgttcaga gtttcacatc aatccagatg tattaaaaat 1380
atataaaagt gaattaaata agcacataga ggatggtatg ggaagaaatt tggctgatcg 1440

```

```

atgcaccgat gaagtaaacg ccttagtgcc tcagaccag caagaaatta ttgaaaattt 1500
gaagccatta cttccagctg gtatacagga taaactacat acactgatcc cttgcaagaa 1560
at ttgatctc agttataatc taaattacca caagttatgt tcagattttc aagaggatat 1620
tgtatttcgt ttttccctgg gctgggtcttc ccttgtagat cgatttttgg gccctagaaa 1680
tgctcaaagg gtgctcctag gattatcaga gcctatcttt cagctcccta gatcttttagc 1740
ttctactccc actgctccta ccactccagc aacgccagat aatgcatcac aggaagaact 1800
catgattaca ttagtaacag gattggcgct cgttacatct agaacttcta tgggcatcat 1860
tattgttggg ggagtgattt ggaaaactat aggctggaaa ctctatctcg tttcatatac 1920
tatgtatgga gctttgtatc tttatgaaag actgagctgg accacccatg ccaaggagcg 1980
agcctttaaag cagcagtttg taaactatgc aactgaaaaa ctgaggatga ttgttagctc 2040
cacgagtgca aactgcagtc accaagtaaa acaacaaata gctaccactt ttgctcgcct 2100
gtgccaacaa gttgatatta ctcaaaaaa gctggaagaa gaaattgcta gattacccaa 2160
agaaatagat cagttggaga aaatacaaaa caattcaaag ctcttaagaa ataaagctgt 2220
tcaacttgaa aatgagctgg agaattttac taagcagttt ctaccttcaa gcaatgaaga 2280
ctcctaacaa tagagattgc tttggtgacc atgtaggag gaaacgaaac ttgtaagatt 2340
ggaacagttg ttatttttat gaaattactt taaatatgaa ttgtactaac tgtacctaaa 2400
tagcaaagcc ctgtgtagat tctggtaatg atctgtctca gggtagtgtt atttttgaag 2460
agtgttatgt ccttagtttt aattttgagt aaagaaaagg cttaatcatg aattagttac 2520
aagcaacagt accaacttat gtgacccctg aggggtgggg ctgtgagctc ttaatttggt 2580
tttgattctg aaaaactctg ctctctggca tccaggagtt agagattgag cctttcatct 2640
tctttctcaa aactagtttt tgatgcttct tttcatggga atagtcactt ttttatttag 2700
taaatacgcat tgctggaacc accaaggagt gtggaatgtc cttgagtgta ttatttatgc 2760
aagtcacagt cacgttgcca tcatggcagc tatgtgaaac actaataaat gtgtttttac 2820
tttttattcc cgtaaaaact gatgtaaaac aggataaagg cttgttatag tcacttataa 2880
gtatctgggt ctaagtaatt tccttagatg tttctaaaga aacattttca gctttgctcc 2940
cattatgatt ccaataagga acgctttcct agtgcaattt taggagtaaa gtttgaagag 3000
ataaaaaatag ccaaagatag gagacgtctg aattttgaat gataaacagt gatgttttaa 3060
aaaagctggt gttcttcagg aggcatttgc ctaggatatt gctggattat accccattgg 3120
aggcttttaa ttttatttgt atgaattttc caggatttca ttaaaaatta ttattgtatt 3180
ttttacctta aaaaaaaaaa aaaa

```

&lt;210&gt; 30

&lt;211&gt; 741

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

```

Met Ala Glu Pro Val Ser Pro Leu Lys His Phe Val Leu Ala Lys Lys
  1                      5                      10                      15

Ala Ile Thr Ala Ile Phe Asp Gln Leu Leu Glu Phe Val Thr Glu Gly
          20                      25                      30

Ser His Phe Val Glu Ala Thr Tyr Lys Asn Pro Glu Leu Asp Arg Ile
          35                      40                      45

Ala Thr Glu Asp Asp Leu Val Glu Met Gln Gly Tyr Lys Asp Lys Leu
          50                      55                      60

Ser Ile Ile Gly Glu Val Leu Ser Arg Arg His Met Lys Val Ala Phe
          65                      70                      75                      80

Phe Gly Arg Thr Ser Ser Gly Lys Ser Ser Val Ile Asn Ala Met Leu
          85                      90                      95

Trp Asp Lys Val Leu Pro Ser Gly Ile Gly His Ile Thr Asn Cys Phe
          100                     105                     110

Leu Ser Val Glu Gly Thr Asp Gly Asp Lys Ala Tyr Leu Met Thr Glu
          115                     120                     125

```

Gly Ser Asp Glu Lys Lys Ser Val Lys Thr Val Asn Gln Leu Ala His  
 130 135 140  
 Ala Leu His Met Asp Lys Asp Leu Lys Ala Gly Cys Leu Val Arg Val  
 145 150 155 160  
 Phe Trp Pro Lys Ala Lys Cys Ala Leu Leu Arg Asp Asp Leu Val Leu  
 165 170 175  
 Val Asp Ser Pro Gly Thr Asp Val Thr Thr Glu Leu Asp Ser Trp Ile  
 180 185 190  
 Asp Lys Phe Cys Leu Asp Ala Asp Val Phe Val Leu Val Ala Asn Ser  
 195 200 205  
 Glu Ser Thr Leu Met Asn Thr Glu Lys His Phe Phe His Lys Val Asn  
 210 215 220  
 Glu Arg Leu Ser Lys Pro Asn Ile Phe Ile Leu Asn Asn Arg Trp Asp  
 225 230 235 240  
 Ala Ser Ala Ser Glu Pro Glu Tyr Met Glu Asp Val Arg Arg Gln His  
 245 250 255  
 Met Glu Arg Cys Leu His Phe Leu Val Glu Glu Leu Lys Val Val Asn  
 260 265 270  
 Ala Leu Glu Ala Gln Asn Arg Ile Phe Phe Val Ser Ala Lys Glu Val  
 275 280 285  
 Leu Ser Ala Arg Lys Gln Lys Ala Gln Gly Met Pro Glu Ser Gly Val  
 290 295 300  
 Ala Leu Ala Glu Gly Phe His Ala Arg Leu Gln Glu Phe Gln Asn Phe  
 305 310 315 320  
 Glu Gln Ile Phe Glu Glu Cys Ile Ser Gln Ser Ala Val Lys Thr Lys  
 325 330 335  
 Phe Glu Gln His Thr Ile Arg Ala Lys Gln Ile Leu Ala Thr Val Lys  
 340 345 350  
 Asn Ile Met Asp Ser Val Asn Leu Ala Ala Glu Asp Lys Arg His Tyr  
 355 360 365  
 Ser Val Glu Glu Arg Glu Asp Gln Ile Asp Arg Leu Asp Phe Ile Arg  
 370 375 380  
 Asn Gln Met Asn Leu Leu Thr Leu Asp Val Lys Lys Lys Ile Lys Glu  
 385 390 395 400  
 Val Thr Glu Glu Val Ala Asn Lys Val Ser Cys Ala Met Thr Asp Glu  
 405 410 415  
 Ile Cys Arg Leu Ser Val Leu Val Asp Glu Phe Cys Ser Glu Phe His  
 420 425 430  
 Pro Asn Pro Asp Val Leu Lys Ile Tyr Lys Ser Glu Leu Asn Lys His  
 435 440 445



Ile Glu Asp Gly Met Gly Arg Asn Leu Ala Asp Arg Cys Thr Asp Glu  
 450 455 460  
 Val Asn Ala Leu Val Pro Gln Thr Gln Gln Glu Ile Ile Glu Asn Leu  
 465 470 475 480  
 Lys Pro Leu Leu Pro Ala Gly Ile Gln Asp Lys Leu His Thr Leu Ile  
 485 490 495  
 Pro Cys Lys Lys Phe Asp Leu Ser Tyr Asn Leu Asn Tyr His Lys Leu  
 500 505 510  
 Cys Ser Asp Phe Gln Glu Asp Ile Val Phe Arg Phe Ser Leu Gly Trp  
 515 520 525  
 Ser Ser Leu Val His Arg Phe Leu Gly Pro Arg Asn Ala Gln Arg Val  
 530 535 540  
 Leu Leu Gly Leu Ser Glu Pro Ile Phe Gln Leu Pro Arg Ser Leu Ala  
 545 550 555 560  
 Ser Thr Pro Thr Ala Pro Thr Thr Pro Ala Thr Pro Asp Asn Ala Ser  
 565 570 575  
 Gln Glu Glu Leu Met Ile Thr Leu Val Thr Gly Leu Ala Ser Val Thr  
 580 585 590  
 Ser Arg Thr Ser Met Gly Ile Ile Ile Val Gly Gly Val Ile Trp Lys  
 595 600 605  
 Thr Ile Gly Trp Lys Leu Leu Ser Val Ser Leu Thr Met Tyr Gly Ala  
 610 615 620  
 Leu Tyr Leu Tyr Glu Arg Leu Ser Trp Thr Thr His Ala Lys Glu Arg  
 625 630 635 640  
 Ala Phe Lys Gln Gln Phe Val Asn Tyr Ala Thr Glu Lys Leu Arg Met  
 645 650 655  
 Ile Val Ser Ser Thr Ser Ala Asn Cys Ser His Gln Val Lys Gln Gln  
 660 665 670  
 Ile Ala Thr Thr Phe Ala Arg Leu Cys Gln Gln Val Asp Ile Thr Gln  
 675 680 685  
 Lys Gln Leu Glu Glu Glu Ile Ala Arg Leu Pro Lys Glu Ile Asp Gln  
 690 695 700  
 Leu Glu Lys Ile Gln Asn Asn Ser Lys Leu Leu Arg Asn Lys Ala Val  
 705 710 715 720  
 Gln Leu Glu Asn Glu Leu Glu Asn Phe Thr Lys Gln Phe Leu Pro Ser  
 725 730 735  
 Ser Asn Glu Asp Ser  
 740

&lt;210&gt; 31

&lt;211&gt; 2483

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

```

cacatgttgc cccaaataca agcacaaatc taaccatgag cttcagcaat cagctcaata 60
cagtgcacaa tcaggccagt gttctagctt ccagttctac tgcagcagct gctactcttt 120
ctctggctaa ttcagatgtc tcaactactaa actaccagtc agctttgtac ccatcatctg 180
ctgcaccagt tcttgaggtt gccacgcagg gtgtttcctt gcagcctgga accaccaga 240
tttgcaactca gacagatcca ttccaacaga catttatagt atgtccacct gcgtttcaaa 300
ctggactaca agcaacaaca aagcattctg gattccctgt gaggatggat aatgctgtac 360
cgattgtacc ccaggcacca gctgctcagc ccactacaga ttcagtcagg agttctcacg 420
cagacttgca gggaaaaaat atccagacat tcttgagaaa tggctctctg aggaagctgt 480
acaccactaa tggtagcaac tctccaccct caagtagcca catcacaccg cagtatgcgg 540
tgccctttac tctgagctgc gcagccggcc ggccggcgct ggttgaacag actgccgctg 600
tactgcaggg gtggcctgga gggactcagc aaattctcct gccttcaact tggcaacagt 660
tgcttggggt agctctacac aactctgtcc agccacagc aatgattcca gaggccatgg 720
ggagtggaca gcagctagct gactggagga atgccactc tcatggcaac cagtacagca 780
ctatcatgca gcagccatcc ttgctgacta accatgtgac attggccact gctcagcctc 840
tgaatgttgg tgttgcccat gttgtcagac aacaacaatc cagtccctc ccttcgaaga 900
agaataagca gtcagctcca gtctcttcca agtctctct agatgttctg ccttcccaag 960
tctattctct ggttgggagc agtcccctcc gcaccacatc ttctataat tcttgggtcc 1020
ctgtccaaga tcagcatcag cccatcatca ttccagatac tcccagccct cctgtgagtg 1080
tcactactat ccgaagtgc actgatgagg aagaggacaa caaatacaag cccagtagct 1140
ctggactgaa gccaaaggtct aatgtcatca gttatgtcac tgtcaatgat tctccagact 1200
ctgactcttc tttgagcagc ccttattcca ctgataccct gagtgtcttc cgaggcaata 1260
gtggatccgt tttggagggg cctggcagag ttgtggcaga tggcactggc accgcacta 1320
tcattgtgcc tccactgaaa actcagcttg gtgactgcac tgtagcaacc caggcctcag 1380
gtctcctgag caataagact aagccagtcg cttcagtgag tgggcagtca tctggatgct 1440
gtatcacccc cacaggggat cgagctcaac gcggggggac cagtgcagca caaccactca 1500
atcttagcca gaaccagcag tcacgcggcg ctccaacctc acaggagaga agcagcaacc 1560
cagcccccgc caggcagcag gcgtttgtgg cccctctctc ccaagcccc tacaccttcc 1620
agcatggcag cccgtacac tcgacagggc acccacacct tgccccggcc cctgtctacc 1680
tgccaagcca ggctcatctg tatacgtatg ctgccccgac ttctgtgtgt gcactgggct 1740
caaccagctc cattgtcat cttttctccc cacagggttc ctcaaggcat gctgcagcct 1800
ataccactca ccctagcact ttggtgcacc aggtccctgt cagtgttggg cccagcctcc 1860
tcacttctgc cagcgtggcc cctgtcagt accaacacca gtttgccacc caatcctaca 1920
ttgggtcttc ccgaggctca acaatttaca ctggataccc gctgagtcct accaagatca 1980
gccagtattc ctactatag ttggtgagca tgagggagga ggaatcatgg ctaccttctc 2040
ctggccctgc gttcttaata ttgggctatg gagagatcct cctttacctc cttgaaattt 2100
cttagccagc aacttgttct gcaggggccc actgaagcag aagggtttttc tctgggggaa 2160
cctgtctcag tgttgactgc attgtttag tcttcccaa gtttgcccta tttttaaat 2220
cattattttt gtgacagtaa ttttggtact tggaagagtt cagatgcca tcttctgcag 2280
ttaccaagga agagagattg ttctgaagtt accctctgaa aaatattttg tctctctgac 2340
ttgatttcta taaatgcttt taaaaacaag tgaagcccct ctttatttca ttttgtgtta 2400
ttgtgattgc tggtcaggaa aaatgctgat agaaggagtt gaaatctgat gacaaaaaaa 2460
aaaaaaaaa aaaaaaaaaa aaa

```

2483

&lt;210&gt; 32

&lt;211&gt; 654

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 32

```

Met Ser Phe Ser Asn Gln Leu Asn Thr Val His Asn Gln Ala Ser Val
  1                      5                      10                     15
Leu Ala Ser Ser Ser Thr Ala Ala Ala Ala Thr Leu Ser Leu Ala Asn
                20                      25                      30

```

Ser Asp Val Ser Leu Leu Asn Tyr Gln Ser Ala Leu Tyr Pro Ser Ser  
 35 40 45  
 Ala Ala Pro Val Pro Gly Val Ala Gln Gln Gly Val Ser Leu Gln Pro  
 50 55 60  
 Gly Thr Thr Gln Ile Cys Thr Gln Thr Asp Pro Phe Gln Gln Thr Phe  
 65 70 75 80  
 Ile Val Cys Pro Pro Ala Phe Gln Thr Gly Leu Gln Ala Thr Thr Lys  
 85 90 95  
 His Ser Gly Phe Pro Val Arg Met Asp Asn Ala Val Pro Ile Val Pro  
 100 105 110  
 Gln Ala Pro Ala Ala Gln Pro Thr Thr Asp Ser Val Arg Ser Ser His  
 115 120 125  
 Ala Asp Leu Gln Gly Lys Asn Ile Gln Thr Phe Leu Arg Asn Gly Leu  
 130 135 140  
 Leu Arg Lys Leu Tyr Thr Thr Asn Gly Ser Asn Ser Pro Pro Ser Ser  
 145 150 155 160  
 Ser His Ile Thr Pro Gln Tyr Ala Val Pro Phe Thr Leu Ser Cys Ala  
 165 170 175  
 Ala Gly Arg Pro Ala Leu Val Glu Gln Thr Ala Ala Val Leu Gln Ala  
 180 185 190  
 Trp Pro Gly Gly Thr Gln Gln Ile Leu Leu Pro Ser Thr Trp Gln Gln  
 195 200 205  
 Leu Pro Gly Val Ala Leu His Asn Ser Val Gln Pro Thr Ala Met Ile  
 210 215 220  
 Pro Glu Ala Met Gly Ser Gly Gln Gln Leu Ala Asp Trp Arg Asn Ala  
 225 230 235 240  
 His Ser His Gly Asn Gln Tyr Ser Thr Ile Met Gln Gln Pro Ser Leu  
 245 250 255  
 Leu Thr Asn His Val Thr Leu Ala Thr Ala Gln Pro Leu Asn Val Gly  
 260 265 270  
 Val Ala His Val Val Arg Gln Gln Gln Ser Ser Ser Leu Pro Ser Lys  
 275 280 285  
 Lys Asn Lys Gln Ser Ala Pro Val Ser Ser Lys Ser Ser Leu Asp Val  
 290 295 300  
 Leu Pro Ser Gln Val Tyr Ser Leu Val Gly Ser Ser Pro Leu Arg Thr  
 305 310 315 320  
 Thr Ser Ser Tyr Asn Ser Leu Val Pro Val Gln Asp Gln His Gln Pro  
 325 330 335  
 Ile Ile Ile Pro Asp Thr Pro Ser Pro Pro Val Ser Val Ile Thr Ile  
 340 345 350

Arg Ser Asp Thr Asp Glu Glu Glu Asp Asn Lys Tyr Lys Pro Ser Ser  
 355 360 365  
 Ser Gly Leu Lys Pro Arg Ser Asn Val Ile Ser Tyr Val Thr Val Asn  
 370 375 380  
 Asp Ser Pro Asp Ser Asp Ser Ser Leu Ser Ser Pro Tyr Ser Thr Asp  
 385 390 395 400  
 Thr Leu Ser Ala Leu Arg Gly Asn Ser Gly Ser Val Leu Glu Gly Pro  
 405 410 415  
 Gly Arg Val Val Ala Asp Gly Thr Gly Thr Arg Thr Ile Ile Val Pro  
 420 425 430  
 Pro Leu Lys Thr Gln Leu Gly Asp Cys Thr Val Ala Thr Gln Ala Ser  
 435 440 445  
 Gly Leu Leu Ser Asn Lys Thr Lys Pro Val Ala Ser Val Ser Gly Gln  
 450 455 460  
 Ser Ser Gly Cys Cys Ile Thr Pro Thr Gly Tyr Arg Ala Gln Arg Gly  
 465 470 475 480  
 Gly Thr Ser Ala Ala Gln Pro Leu Asn Leu Ser Gln Asn Gln Gln Ser  
 485 490 495  
 Ser Ala Ala Pro Thr Ser Gln Glu Arg Ser Ser Asn Pro Ala Pro Arg  
 500 505 510  
 Arg Gln Gln Ala Phe Val Ala Pro Leu Ser Gln Ala Pro Tyr Thr Phe  
 515 520 525  
 Gln His Gly Ser Pro Leu His Ser Thr Gly His Pro His Leu Ala Pro  
 530 535 540  
 Ala Pro Ala His Leu Pro Ser Gln Ala His Leu Tyr Thr Tyr Ala Ala  
 545 550 555 560  
 Pro Thr Ser Ala Ala Ala Leu Gly Ser Thr Ser Ser Ile Ala His Leu  
 565 570 575  
 Phe Ser Pro Gln Gly Ser Ser Arg His Ala Ala Ala Tyr Thr Thr His  
 580 585 590  
 Pro Ser Thr Leu Val His Gln Val Pro Val Ser Val Gly Pro Ser Leu  
 595 600 605  
 Leu Thr Ser Ala Ser Val Ala Pro Ala Gln Tyr Gln His Gln Phe Ala  
 610 615 620  
 Thr Gln Ser Tyr Ile Gly Ser Ser Arg Gly Ser Thr Ile Tyr Thr Gly  
 625 630 635 640  
 Tyr Pro Leu Ser Pro Thr Lys Ile Ser Gln Tyr Ser Tyr Leu  
 645 650

&lt;210&gt; 33

&lt;211&gt; 2731

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (2173)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (2700)

&lt;400&gt; 33

```
ggcactccac ggctgtgaag atggcggcgg ctgcgtggct tcaggtgttg cctgtcattc 60
ttctgcttct gggagctcac ccgtcaccac tgctcgtttt cagtgcggga cgggcaaccg 120
tagctgctgc cgaccggtcc aaatggcaca ttccgatacc gtcggggaaa aattatttta 180
gttttggaaa gatcctcttc agaaatacca ctatcttctt gaagtttgat ggagaacctt 240
gtgacctgtc tttgaatata acctggtatc tgaaaagcgc tgattgttac aatgaaatct 300
ataacttcaa ggcagaagaa gtagagttgt atttggaaaa acttaaggaa aaaagaggct 360
tgtctgggaa atatcaaaca tcatcaaaat tgttccagaa ctgcagtga cttcttataa 420
cacagacctt ttctggagat tttatgcac gactgcctct tttaggagaa aaacaggagg 480
ctaaggagaa tggaacaaac cttaccttta ttggagacaa aaccgcaatg catgaaccat 540
tgcaaaactg gcaagatgca ccatacattt ttattgtaca tattggcatt tcacacctaa 600
aggaatcatc aaaagaaaat tcaactgagta atctttttac catgactgtt gaagtgaagg 660
gtccctatga atacctcaca cttgaagact atcccttgat gatttttttc atggtgatgt 720
gtattgtata tgtcctgttt ggtgttctgt ggctggcatg gtctgcctgc tactggagag 780
atctcctgag aattcagttt tggattgggt ctgtcatctt cctgggaatg cttgagaaag 840
ctgtcttcta tgcggaattt cagaatatcc gacacaaagg agaactctgc caggggtgctt 900
tgatccttgc agagctgctt tcagcagtga aacgctcact ggctcgaacc ctggtcatca 960
tagtcagtct gggatatggc atcgtcaagc cagccttgg agtactctt cataaggttg 1020
tagtagcagg agcctctat ctttgttct ctggcatgga aggggtctct agagtacttg 1080
ggccctagac tgatcttgc tcttggcct ttatccctt ggctttccta gacctgcct 1140
tgtgctggtg gatatttatt agcctgactc aaacaatgaa gctattaaaa cttcggagga 1200
acattgtaaa actctctttg tatcggcatt tcaccaacac gcttattttg gcagtggcag 1260
catccattgt gtttatcatc tggacaacca tgaagttcag aatagtgaac tgtcagtcgg 1320
tgagttataa gcacatttat gaataatgta ctgtcttata aacaactgat ggtgttgatg 1380
acagtggtaa ggttcttcta agttatatac cttataaaaa attagagcta ggtctctact 1440
ctgaggggtg tgatacttcc ctctctctaa gtattctgta ctatcatggt gcttgggtata 1500
gtactttttt gttgttttt tctgactgta ttctccagtt tttgggagag aattttgtaa 1560
gttataacta cagtgtgcta taaccagtc ttattttaac taaaaatctt aagaagtcca 1620
gagtactaaa tattaagtac catatgtgta aataatacta atctgaatag aagccacatc 1680
cttaagatct gagctcaacg actgtgacag taggatttct tcagaagcag ctaaggctct 1740
tattttgttc aataaataat gaaaatgaaa attataaagt ataccaacct aatgtaactt 1800
tctcttacac tgtataaggt aactttctct taaccctgta taaaaccctt tcttaaagct 1860
tctcagaggg atgatgaagc tttgacaaat actctgttcc gttgatgcat tttctttaac 1920
aacagtaagc actacaaggg caaaaactac attcattcac tttgtttccc cacacttacc 1980
acagtactga gcacgtagca ggctcttagt aaacataact tgaatgaaca aataagtgat 2040
ttttgttgta tgccaaaggc tttatgaaca aggggttaag ataattgtga tgaatgttgt 2100
acttctcccc tgtattgtag gactggcggg agctgtgggt agacgatgcc atctggcgct 2160
tgctgttctc cangatectc tttgtcatca tgggttctctg gcgaccatct gcaaacaacc 2220
agagggttct ggactcttct gtttactctg ctaacatgag atgaccatgt catcaattag 2280
gggtggtgca ttgggggaca gtatcagggc tgtgtcatat agtggaagga aactgggcc 2340
tggaatcaga agaactgggt tcctatctca gctctctctt taacttcatg atttttggca 2400
tgcggcctct ccacctctct ggcttagtt tctttctat atactgaggg ggaattaaac 2460
ccagcaacat gaagttcctt tcagctctga cattttgtga taaatacaca ggcatactat 2520
ggaaataaat tgcaagtttg gtttcagacc atcacgataa agcagatatt gagttacata 2580
catattttgt ttttccagtt gcatatcaaa gttatgttta cactattctg tagtctacta 2640
tgtgtgcaat agcattatgt ctaaaaaata tatgcacgtt aattttaaaa cactttgtn 2700
ccaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2731
```

&lt;210&gt; 34

&lt;211&gt; 441

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Met Ala Ala Ala Ala Trp Leu Gln Val Leu Pro Val Ile Leu Leu Leu
  1              5              10              15

Leu Gly Ala His Pro Ser Pro Leu Ser Phe Phe Ser Ala Gly Pro Ala
              20              25              30

Thr Val Ala Ala Ala Asp Arg Ser Lys Trp His Ile Pro Ile Pro Ser
              35              40              45

Gly Lys Asn Tyr Phe Ser Phe Gly Lys Ile Leu Phe Arg Asn Thr Thr
  50              55              60

Ile Phe Leu Lys Phe Asp Gly Glu Pro Cys Asp Leu Ser Leu Asn Ile
  65              70              75              80

Thr Trp Tyr Leu Lys Ser Ala Asp Cys Tyr Asn Glu Ile Tyr Asn Phe
              85              90              95

Lys Ala Glu Glu Val Glu Leu Tyr Leu Glu Lys Leu Lys Glu Lys Arg
              100              105              110

Gly Leu Ser Gly Lys Tyr Gln Thr Ser Ser Lys Leu Phe Gln Asn Cys
              115              120              125

Ser Glu Leu Phe Lys Thr Gln Thr Phe Ser Gly Asp Phe Met His Arg
              130              135              140

Leu Pro Leu Leu Gly Glu Lys Gln Glu Ala Lys Glu Asn Gly Thr Asn
              145              150              155              160

Leu Thr Phe Ile Gly Asp Lys Thr Ala Met His Glu Pro Leu Gln Thr
              165              170              175

Trp Gln Asp Ala Pro Tyr Ile Phe Ile Val His Ile Gly Ile Ser Ser
              180              185              190

Ser Lys Glu Ser Ser Lys Glu Asn Ser Leu Ser Asn Leu Phe Thr Met
              195              200              205

Thr Val Glu Val Lys Gly Pro Tyr Glu Tyr Leu Thr Leu Glu Asp Tyr
              210              215              220

Pro Leu Met Ile Phe Phe Met Val Met Cys Ile Val Tyr Val Leu Phe
              225              230              235              240

Gly Val Leu Trp Leu Ala Trp Ser Ala Cys Tyr Trp Arg Asp Leu Leu
              245              250              255

Arg Ile Gln Phe Trp Ile Gly Ala Val Ile Phe Leu Gly Met Leu Glu
              260              265              270

Lys Ala Val Phe Tyr Ala Glu Phe Gln Asn Ile Arg His Lys Gly Glu
              275              280              285

Ser Val Gln Gly Ala Leu Ile Leu Ala Glu Leu Leu Ser Ala Val Lys

```

290                      295                      300  
 Arg Ser Leu Ala Arg Thr Leu Val Ile Ile Val Ser Leu Gly Tyr Gly  
 305                      310                      315                      320  
 Ile Val Lys Pro Arg Leu Gly Val Thr Leu His Lys Val Val Val Ala  
 325                      330                      335  
 Gly Ala Leu Tyr Leu Leu Phe Ser Gly Met Glu Gly Val Leu Arg Val  
 340                      345                      350  
 Thr Gly Ala Gln Thr Asp Leu Ala Ser Leu Ala Phe Ile Pro Leu Ala  
 355                      360                      365  
 Phe Leu Asp Thr Ala Leu Cys Trp Trp Ile Phe Ile Ser Leu Thr Gln  
 370                      375                      380  
 Thr Met Lys Leu Leu Lys Leu Arg Arg Asn Ile Val Lys Leu Ser Leu  
 385                      390                      395                      400  
 Tyr Arg His Phe Thr Asn Thr Leu Ile Leu Ala Val Ala Ala Ser Ile  
 405                      410                      415  
 Val Phe Ile Ile Trp Thr Thr Met Lys Phe Arg Ile Val Thr Cys Gln  
 420                      425                      430  
 Ser Val Ser Tyr Lys His Ile Tyr Glu  
 435                      440

&lt;210&gt; 35

&lt;211&gt; 1670

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 35

aatcgggctc accccaagt tggggcgggtc attgacaagt cgaagagttg ggtccttgtg 60  
 tatgcatggg tgggatggta agggagaaga ccttggcctg gatgtgccgg gaaccccgga 120  
 aagccttctc agccattgtt gggcctagcc tgggaccoga cagcactcct ggggtggggga 180  
 ctggggagtg ggcaacaggt ggagccatcc ttggcagacc gaccccatgt gcagtccttg 240  
 ggacaggttt ctccctcctg agcacttgta gctccctcctg agggccagtt ccagagacag 300  
 gccgagggtg gcgagtcccc accccatgct ctcttccaga cctcctacga gatgatgatg 360  
 cagtgtgtgt cccgcatgtt ggccccaccc ctgcatgtca tctcaatgcg ctgcatggtc 420  
 cagtttgtgg gacgggaggg caagtacagt ggtgtgctga gctccattgg gaagattttc 480  
 aaagaggaag ggctgctggg attcttctgt ggattaatcc ctcacctcct gggcgatgtg 540  
 gttttcttgt ggggctgtaa cctgctggcc cacttcatca atgcctacct ggtggatgac 600  
 agcttcagcc aggcctgggc catccggagc tataccaagt tcgtgatggg gattgcagtg 660  
 agcatgctga cctacccctt cctgctagtt ggcgacctca tggctgtgaa caactgcggg 720  
 ctgcaagctg ggctcccccc ttactcccca gtgttcaaat cctggattca ctgctggaag 780  
 tacctgagtg tgcagggcca gctcttccga ggctccagcc tgcttttccg cggggtgtca 840  
 tcaggatcat gctttgccct ggagtaacct gaatcatcta aaaaacacgg tctcaacctg 900  
 gccaccgtgg gtgaggcctg accaccttgg gacacctgca agacgactcc aacccaacaa 960  
 caaccagatg tgctccagcc cagccgggct tcagttccat atttgccatg tgtctgtcca 1020  
 gatgtggggt tgagcggggg tggggctgca ccagtggtat tgggtcacc ggcagacctg 1080  
 ggggaagtgga ggcgaggtgg ggagttggca gaatcccat acctgcaga tttgctgagt 1140  
 ctgtcttgtg cagagggcca gagaatggct tatgggggcc caggttggat ggggaaaggc 1200  
 taatggggtc agacccacc ccgtctaccc ctccagtcag ccagcgccc atcctgcagc 1260  
 tcagctggga gcatcattct cctgctttgt acataggggtg tgggtccctg gcacgtggcc 1320  
 accatcatgt ctaggcctat gctaggaggc aaatggccag gctctgcctg tgtttttctc 1380  
 aacactactt ttctgatatg agggcagcac ctgcctctga atgggaaatc atgcaactac 1440

tcagaatgtg tctctctcat ctaatgetca tctgtttaat ggtgatgcct cgcgtacagg 1500  
 atctggttac ctgtgcagtt gtgaataccc agaggttggg cagatcagtg tctctagtcc 1560  
 taccagttt taaagtccat ggtaagattt gacctcatct cccgcaaata aatgtattgg 1620  
 tgatttgtaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1670

<210> 36

<211> 164

<212> PRT

<213> Homo sapiens

<400> 36

Met Gly Gly Met Val Arg Glu Glu Ala Leu Ala Trp Met Cys Arg Glu  
 1 5 10 15  
 Pro Arg Lys Ala Phe Ser Ala Ile Val Gly Pro Ser Leu Gly Pro Asp  
 20 25 30  
 Ser Thr Pro Gly Trp Gly Thr Gly Glu Trp Ala Thr Gly Gly Ala Ile  
 35 40 45  
 Leu Gly Arg Pro Thr Pro Cys Ala Val Pro Gly Thr Gly Phe Ser Leu  
 50 55 60  
 Leu Ser Thr Cys Ser Ser Pro Arg Gly Pro Val Pro Glu Thr Gly Arg  
 65 70 75 80  
 Gly Trp Arg Val Pro Thr Pro Cys Ser Leu Pro Asp Leu Leu Arg Asp  
 85 90 95  
 Asp Asp Ala Val Cys Val Pro His Val Gly Pro Pro Pro Ala Cys His  
 100 105 110  
 Leu Asn Ala Leu His Gly Pro Val Cys Gly Thr Gly Gly Gln Val Gln  
 115 120 125  
 Trp Cys Ala Glu Leu His Trp Glu Asp Phe Gln Arg Gly Arg Ala Ala  
 130 135 140  
 Gly Ile Leu Arg Trp Ile Asn Pro Ser Pro Pro Gly Arg Cys Gly Phe  
 145 150 155 160  
 Leu Val Gly Leu

<210> 37

<211> 1493

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (1415)

<400> 37

ggatggcgcg cgcggggccc gcacgtggag gccggcgcg gggcgcgggc agggccggct 60  
 gctgagacgc gctgctgccc cccgcgcggg cgccgcggct tcaatggcgc catcgcccag 120  
 gaccggcagc cggcaagatg cgaccgccct gccagcatg tcctcaactt tctgggcgct 180  
 catgatcctg gccagcctgc tcatcgcccta ctgcagtcag ctggccgccc gcacctgtga 240  
 gattgtgacc ttggaccggg acagcagcca gcctcggagg acgatcgccc ggcagaccgc 300



```

ccgctgtgcg tgtagaaagg ggcagatcgc cggcaccacg agagcccggc ccgcctgtgt 360
ggacgcaaga atcatcaaga ccaagcagtg gtgtgacatg cttccgtgtc tggaggggga 420
aggctgcgac ttgttaatca accggtcagg ctggacgtgc acgcagcccg gggggaggat 480
aaagaccacc acggtctcct gacaaacaca gccctgagg ggccccggga gtggccttgg 540
ctccctggag agcccacgtc tcagccacag ttctccactc gcctcggact tca:ccgttc 600
tctgccgccc gccactccg tttccctgtg gtccgtgaag gacggcctca ggccttggca 660
tcctgagctt ctgtctgtcc agccgaccg aggaggccg actcagacac ataggcgggg 720
ggcggcacct ggcacagca atacgcagtc tgtgggagcc cggccgcgcc cagccccgc 780
cgaccgtggc gttggccctg ctgtcctcag aggaggagga ggaggaggca gctccggcag 840
ccacagaagg ctgcagccca gcccgctga gacacgacgc ctgcccagg ggactgtcag 900
gcacagaagc ggcctcctcc cgtgccccag actgtccgaa ttggttttat tttcttatac 960
tttcagtata ctccatagac caaagagcaa aatctatctg aacctggacg caccctcact 1020
gtcagggtcc ctggggtcgc ttgtgcgggc gggagggcaa tgggtggcaga gacatgctgt 1080
ggccccggcg gacggagag ggcgggcctg gtggaggcct ccaccccagg agcaccgcc 1140
gcaccctcgg aggacgggtt tcggctgcgc ggaggccgtg gcacacctgc gggaggcagc 1200
gacggcccc cgcagacgc cgggaacgca ggccgcttta ttcctctgta cttagatcaa 1260
cttgaccgta ctaaaatccc tttctgtttt aaccagttaa acatgcctct tctacagtc 1320
catttttgat agttggataa tccagtatct gccaaagca tgttgggtct cccgtgactg 1380
ctgcctcatc gatacccat ttagctccag aaagnaaaga aaactcgagt aacacttggt 1440
tgaaagagat cattaatgt attttgcaaa gcctaaaaaa aaaaaaaaaa aaa 1493

```

&lt;210&gt; 38

&lt;211&gt; 132

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

```

Met Ala Pro Ser Pro Arg Thr Gly Ser Arg Gln Asp Ala Thr Ala Leu
  1              5              10              15

Pro Ser Met Ser Ser Thr Phe Trp Ala Phe Met Ile Leu Ala Ser Leu
          20              25              30

Leu Ile Ala Tyr Cys Ser Gln Leu Ala Ala Gly Thr Cys Glu Ile Val
      35              40              45

Thr Leu Asp Arg Asp Ser Ser Gln Pro Arg Arg Thr Ile Ala Arg Gln
  50              55              60

Thr Ala Arg Cys Ala Cys Arg Lys Gly Gln Ile Ala Gly Thr Thr Arg
  65              70              75              80

Ala Arg Pro Ala Cys Val Asp Ala Arg Ile Ile Lys Thr Lys Gln Trp
          85              90              95

Cys Asp Met Leu Pro Cys Leu Glu Gly Glu Gly Cys Asp Leu Leu Ile
      100              105              110

Asn Arg Ser Gly Trp Thr Cys Thr Gln Pro Gly Gly Arg Ile Lys Thr
      115              120              125

Thr Thr Val Ser
      130

```

&lt;210&gt; 39

&lt;211&gt; 3693

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (108)

&lt;400&gt; 39

```

cgtggccgaa ggatgcccgt ttgtgtctaa acggaggctc ggccacaacg cccctggatt 60
gggtgtaggg cggggcgggc cacagtctcc accctgaagc ggaagtgnag gaaagatgga 120
ggaccatcac cacgtgcccc tgcacatcca gaccagccag ctgctcgatt ggctggtgga 180
cagaaggcac tgcagcctga aatggcagag tctggtgctg acgatccgcy agaagatcaa 240
tgctgccatc caggacatgc cagagagcga agagatcgcc cagctgctgt ctgggtcctg 300
tgagtgttta ggggctgtca ctggagtccc ctctttgctg aggtagtatg tatcagctga 360
gctactctct tgttatttag ccaccagggc tggcctttag agaggggagc atttttattt 420
gaggtataga tgctggcttc ttccactatg aaatgattta gggaaaagct acacactaaa 480
gttactggca gtatattata catgaatcag ccacctgcag tgcccagagt aggtgaagaa 540
gatacttttag ttctcgaaat ccgctctctt gctttctaga cattcactac tttcactgcc 600
taagaatcct ggaccttctc aaaggcacag aggcctccac gaagaatatt tttggccgat 660
actcttcaca gcgatgaag gattggcagg agattatagc tctgtatgag aaggacaaca 720
cctacttagg taaagtggcc cggcctggga gccctggtat ccattgggaa gccactctc 780
agagtcttga gataccaggc ttataggagg cacagtctgt gagtgggaag agactggagt 840
gtagatgttg cccatttgta ggtggtaaaa tcaattgttt ttgatggaat tgattttccc 900
tgagtggagt gctgggggaa ggaggaggct caggccggta gtggccattc gccgtgctc 960
agcgagcagg tgtgtgtggg tcctccacca ctacctctt ggttagcggg agtgtgctgc 1020
ccccaccccc accccgtac cccatttgta cacaaggcag aagggcacg ggttttctctg 1080
ggagcgaata tcaagtgcct gagagcaact acaggactaa ctgtgtttgg gttgggtgta 1140
gtataaataa taataatggc taatatctcc tgagcatcta ctaaatgcaa ggaattgtgc 1200
ttggtgtgtc atgtggattc tctcttgcac ctctcatgata aatgttattg tcgctgtttt 1260
accgatgagg gttggattag aggggttaaa caacttgtct taggctccac agctgggaac 1320
aagtggggct gggaagctga ctctgtgctc ttaccacca caaaggatgt gtgtgcatcc 1380
tggggcatgc ctgcctcatg tgggggtgtc ctgggctgaa tttcctgggc acttctcagt 1440
ggaactctct agcctcctgg ttcggaatgt caactatgag atccctcac tgaagaagca 1500
gattgccaag tgccagcagc tgcagcaaga atacagccgc aaggaggagg agtgccaggc 1560
aggggctgcc gagatgcggg agcagttcta ccactcctgc aagcagtatg gcatcacggt 1620
gagcgggcggc agcctctctg cagccagagg ccctgttgtt cctcctctga 1680
ccccgtctga cccctcagcc tgggtgccc ccctttgggc cagtgtctta ctttctctg 1740
gtctttggat gttttcttca atctgttgga ctccacctct tctccctct ctaggcgaa 1800
aatgtccgag gagaactgct ggccctggtg aaggacctgc cgagtcagct ggctgagatt 1860
ggggcagcgg ctacgagtc cctgggggaa gccattgacg tgtaccaggc gtctgtgggg 1920
ttgtgtgtg agaggtagag aggcctcagc ttctcctggt ggggggtgctt tgcctgtgtt 1980
ccccagtgca tgaccctct ccagttgtct tgttccata taacatttga actctttaca 2040
cacctgaacc tgtggggggc ttgccattt ttgacctgtg gttgtgtgct cccaggccaa agcccagtgt 2100
tggccttacg catggctggc agggagtgca gttgtgtgct ctgttgaagc cccacagagc 2160
aggtgttgcc aatgctgcgg ttcgtgcaga agcggggaaa ctcaacggtg tacgagtga 2220
ggacagggac agagccctct gtggtggaac gacccacct cgaggagctt cctgagcagg 2280
tggcagaaga tgcgattgac tggggcgact ttggggtaga ggcagtgtct gaggggactg 2340
actctggcat ctctgccgag gctgctggaa tcgactgggg catcttcccg gaatcagatt 2400
caaaggatcc tggaggtgat gggatagact ggggagacga tgctgttgct ttgcagatca 2460
cagtgtgga agcaggaacc caggctccag aaggtgttgc cagggggcca gatgccctga 2520
cactgcttga atacactgag acccggaatc agttccttga tgagctcatg gagcttgaga 2580
tcttcttagc ccagagagca gtggagttga gtgaggaggc agatgtcctg tctgtgagcc 2640
agttccagct ggctccagcc atcctgcagg gccagaccaa agagaagatg gttaccatgg 2700
tgtcagtgtc ggaggatctg attggcaagc ttaccagtct tcagctgcaa cacctgttta 2760
tgatcctggc ctccaccaag tctggtctcc ccttgatgca aggtctgtcc atcttgagca 2820
gctctgcctc cttgtattcc tcctctgtt ccatgacccc ttaaacccca tcctgcctc 2880
ctggccattg ccatccactg gggatagggg ttctctttgg gacaagaggg ggagggttta 2940
catatacagg aagaatctgc ttgcttctg agtaggacag gggaaactggg agtgggtttt 3000
ccttaaaagg aaagggttta aggatgtgag ggtaagcggc cagttggggg tttgttccc 3060
gagcctctca cctccccagc agctgaatgg gaatgctcag gatgcacagc taaccagca 3120
ctcacctgag tgccccgcac aggtatgtgg accgagtgac tgaattcctc cagcaaaagc 3180
tgaagcagtc ccagctgctg gctttgaaga aagagctgat ggtgcagaag cagcaggagg 3240
cacttgagga gcaggcggct ctggagccta agctggacct gctactggag aagaccaagg 3300

```

```

agctgcagaa gctgattgaa gctgacatct ccaagaggta cagcgggagc cctgtgaacc 3360
tgatgggaac ctctctgtga caccctccgt gttcttgcct gcccatcttc tccgcttttg 3420
ggatgaagat gatagccagg gctgttggtt tggggccctt caaggcaaaa gaccaggctg 3480
actggaagat ggaaagccac aggaaggaag cggcacctga tggatgatctt ggcactctcc 3540
atgttctcta caagaagctg tgggtgattgg ccctgtgggtc taccaggcga aaaccacaga 3600
ttctccttct agttagtata gcggacttaa taaaagagga aaaaactctt gcttcaaaaa 3660
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa                               3693

```

&lt;210&gt; 40

&lt;211&gt; 230

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 40

```

Met Leu Arg Phe Val Gln Lys Arg Gly Asn Ser Thr Val Tyr Glu Trp
  1              5              10              15

Arg Thr Gly Thr Glu Pro Ser Val Val Glu Arg Pro His Leu Glu Glu
      20              25              30

Leu Pro Glu Gln Val Ala Glu Asp Ala Ile Asp Trp Gly Asp Phe Gly
      35              40              45

Val Glu Ala Val Ser Glu Gly Thr Asp Ser Gly Ile Ser Ala Glu Ala
      50              55              60

Ala Gly Ile Asp Trp Gly Ile Phe Pro Glu Ser Asp Ser Lys Asp Pro
      65              70              75              80

Gly Gly Asp Gly Ile Asp Trp Gly Asp Asp Ala Val Ala Leu Gln Ile
      85              90              95

Thr Val Leu Glu Ala Gly Thr Gln Ala Pro Glu Gly Val Ala Arg Gly
      100             105             110

Pro Asp Ala Leu Thr Leu Leu Glu Tyr Thr Glu Thr Arg Asn Gln Phe
      115             120             125

Leu Asp Glu Leu Met Glu Leu Glu Ile Phe Leu Ala Gln Arg Ala Val
      130             135             140

Glu Leu Ser Glu Glu Ala Asp Val Leu Ser Val Ser Gln Phe Gln Leu
      145             150             155             160

Ala Pro Ala Ile Leu Gln Gly Gln Thr Lys Glu Lys Met Val Thr Met
      165             170             175

Val Ser Val Leu Glu Asp Leu Ile Gly Lys Leu Thr Ser Leu Gln Leu
      180             185             190

Gln His Leu Phe Met Ile Leu Ala Ser Pro Arg Ser Gly Phe Pro Leu
      195             200             205

Met Gln Gly Ser Ala Ile Leu Ser Ser Ser Ala Ser Leu Tyr Ser Ser
      210             215             220

Ser Cys Ser Met Thr Pro
      225             230

```

<210> 41  
 <211> 1701  
 <212> DNA  
 <213> Homo sapiens

<400> 41  
 cccttgagat gattttctct tttcaacttc ttgaacttgg acatgaagga tgtgggcccc 60  
 gaatcatgtg gccagcccac cccctgttgg ccctcaccag ccttggagtc tgttctaggg 120  
 aaggcctccc agcatctggg actcgagagt gggcagcccc tctacctcct ggagctgaac 180  
 tggggtggaa ctgagtgtgt tcttagctct accgggagga cagctgcctg tttcctcccc 240  
 accagcctcc tccccacatc cccagctgcc tggctgggtc ctgaagccct ctgtctacct 300  
 gggagaccag ggaccacagg ccttagggat acagggggtc cccttctgtt accaccccc 360  
 accctcctcc aggacaccac taggtggtgc tggatgcttg ttctttggcc agccaagggtt 420  
 cacggcgatt ctcccatgg gatcttgagg gaccaagctg ctgggattgg gaaggagttt 480  
 caccctgacc gttgccctag ccaggttccc aggaggcctc accatactcc ctttcagggc 540  
 cagggctcca gcaagcccag ggcaaggatc ctgtgctgct gtctggttga gacctgcca 600  
 ccgtgtgtcg ggagtgtggg ccaggctgag tgcataagtg acagggcctg gagcatgggc 660  
 ctgggtgtgt gtgagctcag gcctaggtgc gcagtgtgga gacgggtgtt gtcggggaag 720  
 aggtgtggct tcaaagtgtg tgtgtgcagg ggggtgggtgt gtttagcgtgg gttaggggaa 780  
 cgtgtgtgctg cgtgctgggtg ggcatgtgag atgagtgaat gccggtgaat gtgtccacag 840  
 ttgagagggtt ggagcaggat gagggaatcc tgtcaccatc aataatcact tgtggagcgc 900  
 cagctctgcc caagacgcca cctgggcgga cagccaggag ctctccatgg ccaggctgcc 960  
 tgtgtgcatg ttccctgtct ggtgccctt tgcctgcctc ctgcaaacct cacagggtcc 1020  
 ccacacaaca gtgccctcca gaagcagccc ctcgaggcca gaggaaggaa aatggggatg 1080  
 gctggggctc tctccatcct ccttttctcc ttgccttcgc atggctggcc ttccctcca 1140  
 aaacctccat tccccgtctg ccagcccctt tgccatagcc tgattttggg gaggaggaag 1200  
 gggcgatttg agggagaagg ggagaaagct tatggctggg tctggtttct tcccttcccc 1260  
 gagggtctta ctgttccagg gtgccccag ggcaggcagg ggccacacta tgctgcgccc 1320  
 ctggtaaagg tgaccctgc catttaccag cagccctggc atgttctctg cccacaggaa 1380  
 tagaatggag ggagctccag aaactttcca tcccaaaggc agtctccgtg gttgaagcag 1440  
 actggatttt tgcctctgcc ctgaccctt gtccctcttt gagggagggg agctatgcta 1500  
 ggactccaac ctcagggact cgggtggcct gcgctagctt cttttgatac tgaaaacttt 1560  
 taagggtggg ggggtggcaag ggatgtgctt aataaatcaa ttccaagcct caaaaaaaaa 1620  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1680  
 aaaaaaaaaa aaaaaaaaaa a a 1701

<210> 42  
 <211> 240  
 <212> PRT  
 <213> Homo sapiens

<400> 42  
 Met Lys Asp Val Gly Pro Glu Ser Cys Gly Gln Pro Thr Pro Cys Trp  
 1 5 10 15  
 Pro Ser Pro Ala Leu Glu Ser Val Leu Gly Lys Ala Ser Gln His Leu  
 20 25 30  
 Gly Leu Glu Ser Gly Gln Pro Leu Tyr Leu Leu Glu Leu Asn Trp Gly  
 35 40 45  
 Gly Thr Glu Cys Val Leu Ser Ser Thr Gly Arg Thr Ala Ala Cys Phe  
 50 55 60  
 Leu Pro Thr Ser Leu Leu Pro Thr Ser Pro Ala Ala Trp Leu Gly Pro  
 65 70 75 80  
 Glu Ala Leu Cys Leu Pro Gly Arg Pro Gly Thr Thr Gly Leu Arg Asp  
 85 90 95

Thr Gly Gly Pro Leu Leu Leu Pro Pro Pro Thr Leu Leu Gln Asp Thr  
 100 105 110  
 Thr Arg Trp Cys Trp Met Leu Val Leu Trp Pro Ala Lys Val His Gly  
 115 120 125  
 Asp Ser Pro His Gly Ile Leu Arg Asp Gln Ala Ala Gly Ile Gly Lys  
 130 135 140  
 Glu Phe His Pro Asp Arg Cys Pro Ser Gln Val Pro Arg Arg Pro His  
 145 150 155 160  
 His Thr Pro Phe Gln Gly Gln Gly Ser Ser Lys Pro Arg Ala Arg Ile  
 165 170 175  
 Leu Cys Cys Cys Leu Val Glu Ser Leu Pro Pro Cys Val Gly Ser Val  
 180 185 190  
 Gly Gln Ala Glu Cys Ile Gly Asp Arg Ala Val Ser Met Gly Leu Gly  
 195 200 205  
 Val Cys Glu Leu Arg Pro Arg Cys Ala Val Trp Arg Arg Val Leu Ser  
 210 215 220  
 Gly Lys Arg Cys Gly Phe Lys Val Cys Val Cys Arg Gly Trp Val Cys  
 225 230 235 240

&lt;210&gt; 43

&lt;211&gt; 1784

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

aggtctagaa ttcaatcggg aatatctttt aagtttttaa aaaactggaa taattatatc 60  
 tatctttttt gccgtttata tttaggggtt tttgttgata aaatcaagtc ttgggtgtgg 120  
 cttgctgaat taaatattta tgagtgggtc atttttaagt atagtgaaca agacaccata 180  
 ttaagtacag tgataaagca tctatattct gtaaaaaaaaa aaaaaatctg cctatgcatg 240  
 ttttttaaga aaaaaaaaaat ggctgtatcg gctgttatgg gactgtaatg cgcttagtgg 300  
 tctgacatat actggaaatg tatgtatact ggcgtacttt atattctcta aaatgcttaa 360  
 tgcctttgaa attttgtaat caaaaaaaaa ctttgaaaaa tctaaagggg agagtattct 420  
 ttaaagtttt taacataagc ttgtcaatgc acatgtagat ggtagcatg ttagcaaac 480  
 cttgtgaaat tataataagt ttgtagtac atgtgaaact ctaaatgcat ggcaactgtt 540  
 aatgtcataa cagtttagtt attttgttct gttctgtcat gtgccacaaa atatgtactt 600  
 ttttcacttt tttccctttg tatatcagtt acgggttaca actggttcat tctgaaaaca 660  
 acaacaacaa aagtccattc atatttttta acaattgtat aagtgcccaa gtaattcact 720  
 acagcctaaa gccttgctt tgtaatttga cttctgacat gttggcaatc aaagcatgca 780  
 cttgtaacaa tgaaaaagaa aaagcatttt atattactac tcaataaaat gtgcatgaac 840  
 ttacagaatt ctcatccttc cactgagtcc gctgaaggga tttatgtgca caaccaccat 900  
 gtgtcttcta ggtgctggcc caccaccaca catcacaggc tgatttcac aggcctcttc 960  
 ctaggggcct cgtgatctga ggggtgtgct ctacttcac tgtaagaaag aatcttgggtg 1020  
 gatttgtgtc tcaaatcaga taagagaagc ctgtttaaag agcagatgcc atcttctggc 1080  
 ttcctcaagg agccagttaa aaaaccagag cattcctttt tattgaaaaa taaaattaat 1140  
 ttgttatcag gttgtttcag ttgtattgga tgccctatct atctgctaaa gcaaaaagta 1200  
 ctaggctact aagtgcattt tcatcacaga aaagagttgc atttgtatta acaagaaatt 1260  
 tgtataccca cgcttcagct actatctaact catcacccga agatttaaga tacaccaaat 1320  
 ttcagtttgt ttgtaacatt gttcatcttt agtgcacttt gttttatata ataaagtatg 1380  
 cctgttatat taaataataa gaatatggca attagcgata tagcataccc aaacaaagat 1440  
 gttctcgata cagtctggca aagactatcc caagggtatt ttaatgaatt cagacatttt 1500  
 ttctctgtgga tatttctcca tcctaaaaaa agtggcaacc aaggaaaata ttttagatgca 1560

```

acttactaga gtgatgatgt gaaagaaatg gtgattctgg tatcatgggtg tttattttct 1620
ttcttataac tgcagagaaa atatcctgac taaaaaaaaat tcattttttt ggattccttt 1680
cttttacaaa ttgtgctgag gcaactatgg catagaaata aacatttgac attaaaaataa 1740
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 1784

```

<210> 44  
 <211> 82  
 <212> PRT  
 <213> Homo sapiens

```

<400> 44
Met Cys His Lys Ile Cys Thr Phe Phe Thr Phe Phe Pro Leu Tyr Ile
  1             5             10             15
Ser Tyr Gly Leu Gln Leu Val His Ser Glu Asn Asn Asn Asn Lys Ser
          20             25             30
Pro Phe Ile Phe Phe Asn Asn Cys Ile Ser Ala Gln Val Ile His Tyr
          35             40             45
Ser Leu Lys Pro Cys Leu Cys Asn Leu Thr Ser Asp Met Leu Ala Ile
          50             55             60
Lys Ala Cys Thr Cys Asn Asn Glu Lys Glu Lys Ala Phe Tyr Ile Thr
          65             70             75             80
Thr Gln

```

<210> 45  
 <211> 1034  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (598)

```

<400> 45
ggaagatggc ggccctctggg gcggagccgc aggtcctggg acaatacttg gtgttacgaa 60
aggatctatc acaagctccg ttctcctggc cggcggggcg actggtagcg caggcttgtc 120
acgcggccac cgcgcccttg cactcacc gcgaccacc gcacacagcc gcttacctcc 180
aagagctggg gcgcatgcgc aaagtgggcc tcgaggcccc agatgagacc accctaaagg 240
agctggccga gacctgcaa cagaagaaca ttgaccacat gctgtggctt gagcaaccag 300
agaatatcgc cacttgatt gctctccggc cctaccccaa ggaagaagtg ggccagtatt 360
tgaagaagtt ccgattgttc aagtaactgc tgctttgatg tgtttgaata cgcaggccac 420
ccattccaaa gcatcatgtg ttcttgcag tgtcagcttg ctcccgctct tcagttgtga 480
caatttcttg agggttaagc acatgttcat attaaagttg tcattaataa ctacttcctc 540
ttattaataa gttcaagtgg ggaaggtggg agagcagtat tgtctgggga tcattgcnca 600
aatagaagat ttggttagac tctcctgtgg ggctcaagga aactcccttc cagtcactcg 660
ggtttgaaac tttgcttttg aattccttct tattcacatc cagttatcat atttcattga 720
atccaagata acctcaactt caagatgcgg tagtatttta tgtattgtta aaaaatatgc 780
cggcaaatta aacacttgta tttcaataac aaagatgtta aaatttggcc agtgtggtgg 840
ctcacatctg ttaattccag ggttttggga agccaaggca ggaggatcgc ttgagcccat 900
gagttcaagg ttacagtcag ttctaatacag gccaccgcac tccagcgtgg gcaacagagt 960
gagacacggg ttctataaag attaataaca agttaaaaaa aaaaaaaaaa aaaaaaaaaa 1020
aaaaaaaaaa aaaa 1034

```

<210> 46

<211> 126  
 <212> PRT  
 <213> Homo sapiens

<400> 46  
 Met Ala Ala Ser Gly Ala Glu Pro Gln Val Leu Val Gln Tyr Leu Val  
           1                  5                  10                  15  
 Leu Arg Lys Asp Leu Ser Gln Ala Pro Phe Ser Trp Pro Ala Gly Ala  
                   20                  25                  30  
 Leu Val Ala Gln Ala Cys His Ala Ala Thr Ala Ala Leu His Thr His  
           35                  40                  45  
 Arg Asp His Pro His Thr Ala Ala Tyr Leu Gln Glu Leu Gly Arg Met  
           50                  55                  60  
 Arg Lys Val Val Leu Glu Ala Pro Asp Glu Thr Thr Leu Lys Glu Leu  
           65                  70                  75                  80  
 Ala Glu Thr Leu Gln Gln Lys Asn Ile Asp His Met Leu Trp Leu Glu  
                   85                  90                  95  
 Gln Pro Glu Asn Ile Ala Thr Cys Ile Ala Leu Arg Pro Tyr Pro Lys  
           100                  105                  110  
 Glu Glu Val Gly Gln Tyr Leu Lys Lys Phe Arg Leu Phe Lys  
           115                  120                  125

<210> 47  
 <211> 1626  
 <212> DNA  
 <213> Homo sapiens

<400> 47  
 caacttggtgt agctgaaggt ttgtttgtga cttattacag agcctgtgac ttaaaaaatcc 60  
 ttcccacaac cacaagctaa agtgggagaa gacaaactac ctcacctttt caaccaagag 120  
 ggaggagcaa aaatcagtgat acttttacag aagaacctgc cagcctgtga tgatcctacc 180  
 aaagagaaac ctcaatgagt tatggaattt cctttttggt gaattgagtg ctgtttttgc 240  
 tttttctcaga ttccaaatga gagtatacat ttttctttgt ttgatgtgct ggggtgagatc 300  
 tgataataaaa agaccatgcc ttgaattctc tcagctaagt gtaaaggatt ccttcagaga 360  
 tttattttatt ccgagaatag agaccattct gatgatgtat acaaggaaca acctaaactg 420  
 tgctgagcca ctgtttgaac aaaataactc acttaatggt aatttcaaca cacaaaagaa 480  
 aacagtcttg cttattcacg gatacagacc agtaggctcc atcccattat ggcttcagaa 540  
 cttcgttaagg attttgctga atgaagaaga tatgaatgta attgtagtag actggagccg 600  
 ggggtgctaca acttttattt ataataagagc agttaaaaac accagaaaag ttgctgtgag 660  
 tttgagtggtg cacattaaaa atcttttgaa gcatgggtgca tctcttgaca attttcattt 720  
 cataggtgtg agtttagggg ctcatatcag tggatttggt ggaaagatat ttcattggtca 780  
 acttggaaga ataacagggtc ttgaccctgc tgggccaagg ttctccagaa aaccaccata 840  
 tagcagatta gattacacgg atgcaaagtt tgtggatgtc atccattctg actccaatgg 900  
 aattcaattc attaaatgca accaccagag agcagttcac ttgttcatgg catctttaga 960  
 aacaaactgc aattttattt catttccttg tcgttcatac aaagattaca agactagctt 1020  
 atgtgtggag tgtgactgtt ttaaggaaaa atcatgtcct cggctgggtt atcaagccaa 1080  
 gctattttaa ggtgttttaa aagaaaggat ggaaggaaga cctcttagga cactgtgtt 1140  
 tttggataca agtgccattt attttgttct cagtataatt gttccagata aaactatgat 1200  
 ggatggctcg ttttcattta aattattaaa tcagcttgga atgattgaag agccaaggct 1260  
 ttatgaagaa agataacata tgtaaagag gcacccttac tctaaacaac tagtgacttt 1320  
 aaaagttcta agcgtatcag gagatggaga ccatcctggc taacatgggtg aaaccctgtc 1380  
 tctactaaaa attcagaaaa ttagctgggc atgggtggcag gtgcctgtag tcccagctac 1440

tcaggaggct gaggcaagag aattgcttga acccaggagg tggagggttgc agtgagctga 1500  
 gattgcaccg ctgccctcca gcctgggtga cagagcaaga ctccatttca aataaataaa 1560  
 taaataaata aataaataaa taaataaata aataaataaa gttaaagagt aaaaaaaaaa 1620  
 aaaaaa 1626

<210> 48

<211> 368

<212> PRT

<213> Homo sapiens

<400> 48

Met Ile Leu Pro Lys Arg Asn Leu Asn Glu Leu Trp Asn Phe Leu Phe  
 1 5 10 15  
 Gly Glu Leu Ser Ala Val Phe Ala Phe Leu Arg Phe Gln Met Arg Val  
 20 25 30  
 Tyr Ile Phe Leu Cys Leu Met Cys Trp Val Arg Ser Asp Asn Lys Arg  
 35 40 45  
 Pro Cys Leu Glu Phe Ser Gln Leu Ser Val Lys Asp Ser Phe Arg Asp  
 50 55 60  
 Leu Phe Ile Pro Arg Ile Glu Thr Ile Leu Met Met Tyr Thr Arg Asn  
 65 70 75 80  
 Asn Leu Asn Cys Ala Glu Pro Leu Phe Glu Gln Asn Asn Ser Leu Asn  
 85 90 95  
 Val Asn Phe Asn Thr Gln Lys Lys Thr Val Trp Leu Ile His Gly Tyr  
 100 105 110  
 Arg Pro Val Gly Ser Ile Pro Leu Trp Leu Gln Asn Phe Val Arg Ile  
 115 120 125  
 Leu Leu Asn Glu Glu Asp Met Asn Val Ile Val Val Asp Trp Ser Arg  
 130 135 140  
 Gly Ala Thr Thr Phe Ile Tyr Asn Arg Ala Val Lys Asn Thr Arg Lys  
 145 150 155 160  
 Val Ala Val Ser Leu Ser Val His Ile Lys Asn Leu Leu Lys His Gly  
 165 170 175  
 Ala Ser Leu Asp Asn Phe His Phe Ile Gly Val Ser Leu Gly Ala His  
 180 185 190  
 Ile Ser Gly Phe Val Gly Lys Ile Phe His Gly Gln Leu Gly Arg Ile  
 195 200 205  
 Thr Gly Leu Asp Pro Ala Gly Pro Arg Phe Ser Arg Lys Pro Pro Tyr  
 210 215 220  
 Ser Arg Leu Asp Tyr Thr Asp Ala Lys Phe Val Asp Val Ile His Ser  
 225 230 235 240  
 Asp Ser Asn Gly Ile Gln Phe Ile Lys Cys Asn His Gln Arg Ala Val  
 245 250 255  
 His Leu Phe Met Ala Ser Leu Glu Thr Asn Cys Asn Phe Ile Ser Phe



260	265	270
Pro Cys Arg Ser Tyr Lys Asp Tyr Lys Thr Ser Leu Cys Val Asp Cys		
275	280	285
Asp Cys Phe Lys Glu Lys Ser Cys Pro Arg Leu Gly Tyr Gln Ala Lys		
290	295	300
Leu Phe Lys Gly Val Leu Lys Glu Arg Met Glu Gly Arg Pro Leu Arg		
305	310	315
Thr Thr Val Phe Leu Asp Thr Ser Ala Tyr Tyr Phe Val Leu Ser Ile		
325	330	335
Ile Val Pro Asp Lys Thr Met Met Asp Gly Ser Phe Ser Phe Lys Leu		
340	345	350
Leu Asn Gln Leu Gly Met Ile Glu Glu Pro Arg Leu Tyr Glu Glu Arg		
355	360	365

&lt;210&gt; 49

&lt;211&gt; 1221

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

```

ggaaaagctg agaataatca cctctgataa agatcacaga agctgcccgg gaggtgtttg 60
attaaattca tgtattgaaa atattgttca gaccccatgt gacataactg gagccagtgc 120
agtgccatga agaactacga gattagcctg gatattaact tgtcttctag agaatagatt 180
tcatgttcca ttcttctgca atggttaatt cacacagaaa accaatgttt aacattcaca 240
gaggatttta ctgcttaaca gccatcttgc cccaaatatg catttgttct cagttctcag 300
tgccatctag ttatcacttc actgaggatc ctggggcttt ccagtagacc actaatgggg 360
aacgatttcc ttggcaggag ctaaggctcc ccagtgtggt cattcctctc cattatgacc 420
tctttgtcca cccaatctc acctctctgg actttgttgc atctgagaag atcgaagtct 480
tggtcagcaa tgctaccag tttatcatct tgcacagcaa agatcttgaa atcacgaatg 540
ccacccttca gtcagaggaa gattcaagat acatgaaacc aggaaaagaa ctgaaagttt 600
tgagttaccc tgctcatgaa caaattgcac tgctggttcc agagaaactt acgcctcacc 660
tgaaatacta tgtggctatg gacttccaag ccaagttagg tgatggcttt gaagggtttt 720
ataaaagcac atacagaact cttgggtggt aaacaagaat tcttgcacta acagattttg 780
agccaaccca ggcacgcatg gctttccctt gctttgatga accgttggtc aaagccaact 840
tttcaatcaa gatacgaaga gagagcagggc atattgcact atccaacatg ccaaagggtg 900
ccatctatgc atcccagac aaacggaatc aaacacatta tgctttgcag gcatcactga 960
agctacttga tttttatgaa aagtactttg atatctacta tccactctcc aaactgggta 1020
tgttcaaatt ccacattatt gtcttcattt ttgctcataa aacttgctta gatctcttcc 1080
ctctttctct ttgtatgtga tttaaatgag cactgaggaa ttcagttagc tcaggaaaaa 1140
ataatttggt cctcagagat gattcttgag tgtagaaaat aaaatattta tgacatgccc 1200
caaaaaaaaa aaaaaaaaaa a

```

&lt;210&gt; 50

&lt;211&gt; 305

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

Met Phe His Ser Ser Ala Met Val Asn Ser His Arg Lys Pro Met Phe
1 5 10 15
Asn Ile His Arg Gly Phe Tyr Cys Leu Thr Ala Ile Leu Pro Gln Ile
20 25 30

Cys Ile Cys Ser Gln Phe Ser Val Pro Ser Ser Tyr His Phe Thr Glu  
           35                                  40                                  45  
 Asp Pro Gly Ala Phe Pro Val Ala Thr Asn Gly Glu Arg Phe Pro Trp  
           50                                  55                                  60  
 Gln Glu Leu Arg Leu Pro Ser Val Val Ile Pro Leu His Tyr Asp Leu  
           65                                  70                                  75                                  80  
 Phe Val His Pro Asn Leu Thr Ser Leu Asp Phe Val Ala Ser Glu Lys  
                                   85                                  90                                  95  
 Ile Glu Val Leu Val Ser Asn Ala Thr Gln Phe Ile Ile Leu His Ser  
                                   100                                  105                                  110  
 Lys Asp Leu Glu Ile Thr Asn Ala Thr Leu Gln Ser Glu Glu Asp Ser  
           115                                  120                                  125  
 Arg Tyr Met Lys Pro Gly Lys Glu Leu Lys Val Leu Ser Tyr Pro Ala  
           130                                  135                                  140  
 His Glu Gln Ile Ala Leu Leu Val Pro Glu Lys Leu Thr Pro His Leu  
           145                                  150                                  155                                  160  
 Lys Tyr Tyr Val Ala Met Asp Phe Gln Ala Lys Leu Gly Asp Gly Phe  
                                   165                                  170                                  175  
 Glu Gly Phe Tyr Lys Ser Thr Tyr Arg Thr Leu Gly Gly Glu Thr Arg  
           180                                  185                                  190  
 Ile Leu Ala Val Thr Asp Phe Glu Pro Thr Gln Ala Arg Met Ala Phe  
           195                                  200                                  205  
 Pro Cys Phe Asp Glu Pro Leu Phe Lys Ala Asn Phe Ser Ile Lys Ile  
           210                                  215                                  220  
 Arg Arg Glu Ser Arg His Ile Ala Leu Ser Asn Met Pro Lys Val Ser  
           225                                  230                                  235                                  240  
 Ile Tyr Ala Ser Pro Asp Lys Arg Asn Gln Thr His Tyr Ala Leu Gln  
                                   245                                  250                                  255  
 Ala Ser Leu Lys Leu Leu Asp Phe Tyr Glu Lys Tyr Phe Asp Ile Tyr  
           260                                  265                                  270  
 Tyr Pro Leu Ser Lys Leu Gly Met Phe Lys Phe His Ile Ile Val Phe  
           275                                  280                                  285  
 Ile Phe Ala His Lys Thr Cys Leu Asp Leu Phe Pro Leu Ser Leu Cys  
           290                                  295                                  300  
 Met  
 305

&lt;210&gt; 51

&lt;211&gt; 951

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

```

gggtgggtgcg gagtctgceg ccgttcccgc ggctctctcc tcctccccgt tcccttcacc 60
cccaccccgcc acccctttcc ccattcccgc tccgtcaccc tcccgteccc cacactcagg 120
acaagaatgc cctgcccgga acaacccagc agcgccctaga tggctttggt cacgggtccag 180
cggtcaccta cccccagcac cacctccagc ccctgcgcct cggaggcaga cagtggggag 240
gaagaatgcc gggtcacagcc caggagcatc agcgagagct ttctaactgt caaagggtgct 300
gccctttttc taccacgggg aaatgggtca tccacaccaa gaatcagcca cagacgggaa 360
aagcatgcag gcgatctcca acagcatctc caagcaatgt tcattttact ccgcccagaa 420
gacaacatca ggctgggtgt aagactggaa agtacttacc agaatcgaac acgctatatg 480
gtagtgggtt caactaatgg tagacaagac actgaagaaa gcatcgctct aggaatggat 540
ttctcctcta atgacagcac ttgtaccatg ggcttagttt tgcctctctg gagcgacacg 600
ctaattcatt tggatggtga tgggtgggttc agtgtatcga cggataacag agttcacata 660
ttcaaacctg tatctgtgca ggcaatgtgg gtgacagggg attcaaggaa caaacactgt 720
gatgtactat tgggtggaaga atgaactgga gcagcctttc tggagagtga tttgccaata 780
tgccttatca ttttgcatga tctttgtcct agtaactcta tttctatgga tttactctaa 840
gtttgtaaac atggatgtgt gcaaagattt tagctctaag aatgtttgtc agtggttctaa 900
taatagcaaa aaataaaaaa caaatgattg aaaaataaaa aaaaaaaaaa a 951

```

&lt;210&gt; 52

&lt;211&gt; 194

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

```

Met Ala Leu Val Thr Val Gln Arg Ser Pro Thr Pro Ser Thr Thr Ser
  1                      5                      10                      15

Ser Pro Cys Ala Ser Glu Ala Asp Ser Gly Glu Glu Glu Cys Arg Ser
                20                      25                      30

Gln Pro Arg Ser Ile Ser Glu Ser Phe Leu Thr Val Lys Gly Ala Ala
        35                      40                      45

Leu Phe Leu Pro Arg Gly Asn Gly Ser Ser Thr Pro Arg Ile Ser His
  50                      55                      60

Arg Arg Asn Lys His Ala Gly Asp Leu Gln Gln His Leu Gln Ala Met
  65                      70                      75                      80

Phe Ile Leu Leu Arg Pro Glu Asp Asn Ile Arg Leu Ala Val Arg Leu
                85                      90                      95

Glu Ser Thr Tyr Gln Asn Arg Thr Arg Tyr Met Val Val Val Ser Thr
        100                      105                      110

Asn Gly Arg Gln Asp Thr Glu Glu Ser Ile Val Leu Gly Met Asp Phe
        115                      120                      125

Ser Ser Asn Asp Ser Thr Cys Thr Met Gly Leu Val Leu Pro Leu Trp
        130                      135                      140

Ser Asp Thr Leu Ile His Leu Asp Gly Asp Gly Gly Phe Ser Val Ser
        145                      150                      155                      160

Thr Asp Asn Arg Val His Ile Phe Lys Pro Val Ser Val Gln Ala Met
                165                      170                      175

Trp Val Asp Arg Asp Ser Arg Asn Lys His Cys Asp Val Leu Leu Val
        180                      185                      190

```

Glu Glu

<210> 53  
 <211> 1514  
 <212> DNA  
 <213> Homo sapiens

<400> 53  
 gcatgatatt tttacgggttc acccatattg catgtatcag gaatataatc ctttttatta 60  
 ttgagtagtg ttctattgta tgtatataacc acagtttatt tctcccttca tcctttgcta 120  
 gattttgggg ttttttcaca ttgcgctatt cagtataaac ctgctctcaa cattcatgtg 180  
 caagtctttg agtggacata tatttgcgtt tctcttgagt gaatgcacct tgttgggtca 240  
 cgtggcttaa cttaaaaaaa ttttaatcac tgtggtgcat atgtagtgat tattagtgat 300  
 tatctcataa ttttattttc ttgtttaatg atgttgagtg tatttcattt gtattttagt 360  
 ttgcaaagt ttgttcaa tcttcacctg ttttaaatga agacgtacga cttatttttg 420  
 tgttctgaac ataagttctt tgtcacataa aatgtgctat gaatggtgag ttttaaatac 480  
 tccaaatgaa tggctagaga attactattt gtagaaatat ttatatgtca aagggatgct 540  
 aacaattttac tttattgctc taaaatagaa aagttgccag aatgctgtgg agtttttagtg 600  
 gaaaacatga tagctggtgt tactgagtaa atttgagtgt taaatgtcaa tgtaagctaa 660  
 cggccaagat agggaccact gcagggtggt tacttgacgc tgtgactcaa ctggtccttc 720  
 actgccaaac atacctgggg ttggatcatt ggcctgacgt ttgcaaattg aggaacctta 780  
 gggcaaatca gtgaacttct gaactgcctt cgtcttcagt tatatgggga tttccccact 840  
 tttgagatcc ttgtaaggat tatatgagat gaagagatga gacaaggat ataaaagtcc 900  
 tagcacagag cgtgtcatat aatatggctt cacaagtacc ctcattctct tccagtcgt 960  
 tttttgtttt tgtttttgtt tttttgagac catctcactc tgttgcccag gctggagtgc 1020  
 ctcttcattt ttatttcttt attcagcaag tattgatcaa atgtgctttg taccaggtac 1080  
 tgagctcttc gttgggatat aatggtgatc aaggagattg tagattctgg cagggaaaac 1140  
 tgacatcaaa cacggcgacc cgacatagtg agaccctgtc tctactagaa gaactttaaa 1200  
 aatcacctag gtgtgggccc ggcacggtgg ctaacgcctg tgggtcccagc actttgggat 1260  
 gctgaggcgg gtggatcacg gggtcaggag atcgagacca tcctggataa cacggagaaa 1320  
 ccccgctctt actggaaata caaggaaatt ggccgggctg gggggcgggc atctgtggtc 1380  
 ccaattactc gggaggctgc agcaggagag tggcatgaac ccgggaggcg gatcttgcat 1440  
 tgagccgaga tcacgccact gcactccagc ctgggcgaca gaatgagact ccatctcaaa 1500  
 aaaaaaaaaa aaaa 1514

<210> 54  
 <211> 91  
 <212> PRT  
 <213> Homo sapiens

<400> 54  
 Met Ala Ser Gln Val Pro Ser Ser Pro Phe Gln Ser Phe Phe Val Phe  
 1 5 10 15  
 Val Phe Val Phe Leu Arg Pro Ser His Ser Val Ala Gln Ala Gly Val  
 20 25 30  
 Pro Leu His Phe Tyr Phe Phe Ile Gln Gln Val Leu Ile Lys Cys Ala  
 35 40 45  
 Leu Tyr Gln Val Leu Ser Ser Ser Leu Gly Tyr Asn Gly Asp Gln Gly  
 50 55 60  
 Asp Cys Arg Phe Trp Gln Gly Lys Leu Thr Ser Asn Thr Ala Thr Arg  
 65 70 75 80  
 His Ser Glu Thr Leu Ser Leu Leu Glu Glu Leu

85

90

<210> 55  
 <211> 1417  
 <212> DNA  
 <213> Homo sapiens

<400> 55  
 gtccaaatcc tattgtccac agtcagactt ctacaacctc ctctgaacaa atgcagcctc 60  
 caatgtttca ctctcaaagt accattgctg tgttacaggg ctcttcagtt cctcaagacc 120  
 agcagtcaac caacataatct ctttcccaga gtcccatgaa taatcttcag actaacacag 180  
 tagcccaaga agcatttttt gcagcaccga actcaatttc tccacttcag tcaacatcaa 240  
 acagtgaaca acaagctgct ttccaacagc aaagctccaat atcacacatc cagactccta 300  
 tgctttccca agaacaggca caacccccgc agcagggttt atttcagcct cagggtggccc 360  
 tgggctccct tccacctaata ccaatgcctc aaagccaaca aggaaccatg ttccagtcac 420  
 agcactcaat agttgccatg cagagtaact ctccatccca ggaacagcag cagcagcagc 480  
 aacagcagca gcaacagcag cagcaacaac aacagagcat tttattcagt aatcagaata 540  
 ccatggctac aatggcgtct ccaaagcaac caccaccaa catgatattc aacccaaatc 600  
 aaaatccaat ggctaatacag gagcaacaga accagtcaat ttttcaccaa caaagtaaca 660  
 tggccccaat gaatcaagag caacagccca tgcaatttca gagtcagtc acagtttcct 720  
 cacttcagaa cccaggtcct acccagtcgg aatcatcaca gacccccctg ttccatagct 780  
 ctcttcagat tcagttggta caagggtcac ctagtcttca agagcagcaa gtaactctct 840  
 tcttatctcc agcatccatg tctgccttgc agaccagtat aaatcaacaa gatatgcaac 900  
 agtctcctct ttattcccct cagaacaaca tgccctggaat tcaaggagcc acattttcgc 960  
 ctcaaccaca ggctacttta ttccacaaca cagcaggagg cacaatgaac caactgcaga 1020  
 attctcctgg ctcatctcag cagacatcag gaatgttctt atttggcatt caaaataact 1080  
 gtagtcagct tttaacctct ggaccagcta cattgcctga tcagttgatg gccataagtc 1140  
 agccaggcca accacaaaac gagggccagc cacctgtgac aacacttctt tctcagcaaa 1200  
 tgccagagaa ttctccactg gcacccctta taaacaccaa ccagaacatc gaaaagattg 1260  
 atttgcttgt ttcatgtcaa aaccaaggga acaacttgac tggctccttt taactggata 1320  
 taaattccac gaagaaaatc ctgattccaa gatgtcctga gatcttgtgg ttccatgaga 1380  
 attattactt taaaaacaaa acaaaaaaaaa aaaaaaa 1417

<210> 56  
 <211> 420  
 <212> PRT  
 <213> Homo sapiens

<400> 56  
 Met Gln Pro Pro Met Phe His Ser Gln Ser Thr Ile Ala Val Leu Gln  
 1 5 10 15  
 Gly Ser Ser Val Pro Gln Asp Gln Gln Ser Thr Asn Ile Phe Leu Ser  
 20 25 30  
 Gln Ser Pro Met Asn Asn Leu Gln Thr Asn Thr Val Ala Gln Glu Ala  
 35 40 45  
 Phe Phe Ala Ala Pro Asn Ser Ile Ser Pro Leu Gln Ser Thr Ser Asn  
 50 55 60  
 Ser Glu Gln Gln Ala Ala Phe Gln Gln Gln Ala Pro Ile Ser His Ile  
 65 70 75 80  
 Gln Thr Pro Met Leu Ser Gln Glu Gln Ala Gln Pro Pro Gln Gln Gly  
 85 90 95  
 Leu Phe Gln Pro Gln Val Ala Leu Gly Ser Leu Pro Pro Asn Pro Met  
 100 105 110

Pro Gln Ser Gln Gln Gly Thr Met Phe Gln Ser Gln His Ser Ile Val  
 115 120 125  
 Ala Met Gln Ser Asn Ser Pro Ser Gln Glu Gln Gln Gln Gln Gln  
 130 135 140  
 Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Ser Ile Leu Phe Ser  
 145 150 155 160  
 Asn Gln Asn Thr Met Ala Thr Met Ala Ser Pro Lys Gln Pro Pro Pro  
 165 170 175  
 Asn Met Ile Phe Asn Pro Asn Gln Asn Pro Met Ala Asn Gln Glu Gln  
 180 185 190  
 Gln Asn Gln Ser Ile Phe His Gln Gln Ser Asn Met Ala Pro Met Asn  
 195 200 205  
 Gln Glu Gln Gln Pro Met Gln Phe Gln Ser Gln Ser Thr Val Ser Ser  
 210 215 220  
 Leu Gln Asn Pro Gly Pro Thr Gln Ser Glu Ser Ser Gln Thr Pro Leu  
 225 230 235 240  
 Phe His Ser Ser Pro Gln Ile Gln Leu Val Gln Gly Ser Pro Ser Ser  
 245 250 255  
 Gln Glu Gln Gln Val Thr Leu Phe Leu Ser Pro Ala Ser Met Ser Ala  
 260 265 270  
 Leu Gln Thr Ser Ile Asn Gln Gln Asp Met Gln Gln Ser Pro Leu Tyr  
 275 280 285  
 Ser Pro Gln Asn Asn Met Pro Gly Ile Gln Gly Ala Thr Phe Ser Pro  
 290 295 300  
 Gln Pro Gln Ala Thr Leu Phe His Asn Thr Ala Gly Gly Thr Met Asn  
 305 310 315 320  
 Gln Leu Gln Asn Ser Pro Gly Ser Ser Gln Gln Thr Ser Gly Met Phe  
 325 330 335  
 Leu Phe Gly Ile Gln Asn Asn Cys Ser Gln Leu Leu Thr Ser Gly Pro  
 340 345 350  
 Ala Thr Leu Pro Asp Gln Leu Met Ala Ile Ser Gln Pro Gly Gln Pro  
 355 360 365  
 Gln Asn Glu Gly Gln Pro Pro Val Thr Thr Leu Leu Ser Gln Gln Met  
 370 375 380  
 Pro Glu Asn Ser Pro Leu Ala Ser Ser Ile Asn Thr Asn Gln Asn Ile  
 385 390 395 400  
 Glu Lys Ile Asp Leu Leu Val Ser Leu Gln Asn Gln Gly Asn Asn Leu  
 405 410 415  
 Thr Gly Ser Phe  
 420

<210> 57  
 <211> 2297  
 <212> DNA  
 <213> Homo sapiens

<400> 57  
 gaagtgaggg ttgaatgatc ccactttaac taaaaaatga ataagcgtac ttgaaatgat 60  
 tttttaaagt gtttggtagt ctatacttat gttctttctt tgtttccact atagacagta 120  
 ttcgtggcta ctttggggaa acaattgctc tgtactttgg atttttggag tatttcactt 180  
 ttgcattaat ccccatggct gtcattgggt taccttacta cttgtttgtg tgggaagact 240  
 atgacaagta cgtgatcttt gcctcgttca acctcatctg gtccacgggtg attctggaac 300  
 tgtggaagcg tggctgtgcc aacatgacct acaggtgggg gacactgctc atgaagagaa 360  
 agtttgagga gcccggcca ggatttcatt ggtcttggg tatcaattcc atcactggga 420  
 agggaggagcc tctgtacccc agctacaaga gacagttgcg catttacctg gtctccctgc 480  
 cattcgtgtg cctctgcctc tatttctcac tgtatgtcat gatgatttac ttcgacatgg 540  
 aggtttgggc cttgggtcta catgagaaca ggggtctga gtggaccagt gtctgttgt 600  
 atgtgccag catcatctat gccattgtga ttgagatcat gaatcgtctc tatcgatatg 660  
 ctgccgagtt tttaacttca tgggagaatc acagattgga atctgcctat cagaaccatc 720  
 taattctgaa agtttttagtg ttcaacttcc tcaattgctt tgccctacte ttctatatgt 780  
 cctttgtcct gaaagatatg aagcttttgc gccagagctt ggccactctc ctaattacct 840  
 cccagatcct caaccaaatt atggaatctt ttcttcctta ttggctccaa aggaagcatg 900  
 gtgtgcaggt gaagaggaag gtgcaggctt taaaggcaga cattgatgct acattatatg 960  
 aacaagtcac cctggaaaaa gaaatgggaa cttatttggg cacctttgat gattacttgg 1020  
 agttattcct gcagtttggg tatgtgagcc tttctcctg tgtttaccca ttagcagctg 1080  
 cctttgctgt gttaataaac ttcactgaag taaattcaga tgccttaaaa atgtgcaggg 1140  
 tcttcaaacg tccattctca gaaccttcag ccaatattgg tgtgtggcag atgatatttt 1200  
 gtttgacac aggtgtaaag agagggtga attgcaaggt catgaggaat cttttggggg 1260  
 aaatggaaat gtcctgtgtc ttgtttgtgg tgggtgtggg ttcacaggta aatacaccta 1320  
 tcaaaaggta atgaagtata taccttaaat gcatgcagtt tattgtatgt aaattataac 1380  
 tcaatatagg tgatttttaa aaaaacctga aagtttagtt acaaacatat tgcaagttca 1440  
 ggaagccagg cactgtcata tgctgctggt ggaagcatga gctgggtcaac ccatggagtg 1500  
 cagcttcgta tttatctatc aaaattacaa atgcatgtcc cctttgactc agccatttca 1560  
 ctttccagaa tttagcctaa ttccacattt gttaaatgat gtacttataa gatcaccaat 1620  
 tgtagcactg tttgtaatag caactaaatg cccaccaata agaaaatggg tacataaatt 1680  
 ctgatacgtc catgtaataa aatgcagaag cagtgtggca aagaatgagg gagctctttt 1740  
 agtattgaca cagaaagtcc ctcaagacac tttaaatgac taaagcaagg ggcctgacag 1800  
 tatgtagtat gctgaaaagg gagtaggaaa gagtgtatat attcaatatg cttattttgc 1860  
 atacaaactg tctggaagaa tacataagaa attgcaaata gtggttgtct tctagggaga 1920  
 atgggagctg cgaagtggaa gaaaagggga caagggaaaa agacactctt cactatgtac 1980  
 ttagaatttt ttatttttaa gccatgagaa tataatcaaa agtaaataaa tagaaatttt 2040  
 aaactaaagt aaagacagtc ttagatttct tatgaagaaa actgtggaaa aataatcaga 2100  
 ctacacaaag atttctaaat tgtaagatgg cagaagttct ccatgggaca aaatgtattc 2160  
 atattaagtt tggattcagt catcttttgt tttgcttctt ttaaaattaa tgtttctaat 2220  
 agtacttgtg tttcttgtaa ttaatgggta aattattaag tgggtgatcac catatacttt 2280  
 gtaaaaaaaa aaaaaaa 2297

<210> 58  
 <211> 378  
 <212> PRT  
 <213> Homo sapiens

<400> 58  
 Met Ala Val Ile Gly Leu Pro Tyr Tyr Leu Phe Val Trp Glu Asp Tyr  
 1 5 10 15  
 Asp Lys Tyr Val Ile Phe Ala Ser Phe Asn Leu Ile Trp Ser Thr Val  
 20 25 30

Ile Leu Glu Leu Trp Lys Arg Gly Cys Ala Asn Met Thr Tyr Arg Trp  
           35                          40                          45  
 Gly Thr Leu Leu Met Lys Arg Lys Phe Glu Glu Pro Arg Pro Gly Phe  
           50                          55                          60  
 His Gly Val Leu Gly Ile Asn Ser Ile Thr Gly Lys Glu Glu Pro Leu  
           65                          70                          75                          80  
 Tyr Pro Ser Tyr Lys Arg Gln Leu Arg Ile Tyr Leu Val Ser Leu Pro  
                           85                          90                          95  
 Phe Val Cys Leu Cys Leu Tyr Phe Ser Leu Tyr Val Met Met Ile Tyr  
                           100                          105                          110  
 Phe Asp Met Glu Val Trp Ala Leu Gly Leu His Glu Asn Ser Gly Ser  
                           115                          120                          125  
 Glu Trp Thr Ser Val Leu Leu Tyr Val Pro Ser Ile Ile Tyr Ala Ile  
           130                          135                          140  
 Val Ile Glu Ile Met Asn Arg Leu Tyr Arg Tyr Ala Ala Glu Phe Leu  
           145                          150                          155                          160  
 Thr Ser Trp Glu Asn His Arg Leu Glu Ser Ala Tyr Gln Asn His Leu  
                           165                          170                          175  
 Ile Leu Lys Val Leu Val Phe Asn Phe Leu Asn Cys Phe Ala Ser Leu  
                           180                          185                          190  
 Phe Tyr Ile Ala Phe Val Leu Lys Asp Met Lys Leu Leu Arg Gln Ser  
                           195                          200                          205  
 Leu Ala Thr Leu Leu Ile Thr Ser Gln Ile Leu Asn Gln Ile Met Glu  
           210                          215                          220  
 Ser Phe Leu Pro Tyr Trp Leu Gln Arg Lys His Gly Val Gln Val Lys  
           225                          230                          235                          240  
 Arg Lys Val Gln Ala Leu Lys Ala Asp Ile Asp Ala Thr Leu Tyr Glu  
                           245                          250                          255  
 Gln Val Ile Leu Glu Lys Glu Met Gly Thr Tyr Leu Gly Thr Phe Asp  
                           260                          265                          270  
 Asp Tyr Leu Glu Leu Phe Leu Gln Phe Gly Tyr Val Ser Leu Phe Ser  
                           275                          280                          285  
 Cys Val Tyr Pro Leu Ala Ala Ala Phe Ala Val Leu Asn Asn Phe Thr  
           290                          295                          300  
 Glu Val Asn Ser Asp Ala Leu Lys Met Cys Arg Val Phe Lys Arg Pro  
           305                          310                          315                          320  
 Phe Ser Glu Pro Ser Ala Asn Ile Gly Val Trp Gln Met Ile Phe Cys  
                           325                          330                          335  
 Leu Asp Thr Gly Val Lys Arg Gly Leu Asn Cys Lys Val Met Arg Asn  
           340                          345                          350



Leu Leu Gly Glu Met Glu Met Ser Cys Val Leu Phe Val Val Val Val  
 355 360 365

Val Ser Gln Val Asn Thr Pro Ile Lys Arg  
 370 375

<210> 59

<211> 4145

<212> DNA

<213> Homo sapiens

<400> 59

```

aggtctagaa ttcaagatga agtaaagaag gaaagagagg gtctggagaa tgacttgaaa 60
tctgtgaatt ttgacatgac aagcaagttt ttgacagccc tggctcaaga tgggtgtgata 120
aatgaagaag ctctttctgt tactgaacta gatcgagtct atggaggtct tacaactaaa 180
gtccaagaat ctctaaagaa acaggaggga ctcttataaa atattcaggt ctcacatcag 240
gaattttcaa aaatgaaaca atctaataat gaagctaact taagagaaga agttttgaag 300
aatttagcta ctgcatatga caactttgtt gaactgttag ctaatttgaa ggaaggcaca 360
aagttttaca atgagttgac tgaaatcctg gtcaggttcc agaacaaatg cagtgtatata 420
gtttttgcac ggaagacaga aagagatgaa ctcttaaagg acttgcaaca aagcattgcc 480
agagaacctt gtgctccttc aattcctaca cctgcgtatc agtcctcacc agcaggagga 540
catgcaccaa ctctcctaac tccagcgcca agaaccatgc cgcctactaa gccccagccc 600
ccagccaggc ctccaccacc tgtgcttcca gcaaatcgag ctcttctctg tactgtctcca 660
tctccagtgg gggctgggac tgctgcgcca gctccatcac aaacgcctgg ctcagctcct 720
cctccacagg cgcagggacc accctatccc acctatccag gatatcctgg gtattgcca 780
atgcccagtc ccatgggcta taatccttat gcgtatggcc agtataatat gccatatcca 840
ccagtgtatc accagagtcc tggacaggct ccataccggg gaccccagca gccttcatac 900
cccttccttc agccccaca gcagtcttac tatccacagc agtaatatgt ctgctcagca 960
gtcagctga ttcatatcag agggaaagaa ataccaaccc tgcaataagt gtactaaact 1020
ctacgctctg gttaatgtaa tgtactctcc tggactgaat gcagtgtata atttctgtct 1080
acagctagaa gctgtgcccc agttccacat ttgattacac atgtgagatt tgctgtgtgt 1140
gcagtataaa cactagggat aataggattt gaaattgcat tacagttcat aaaaattgaa 1200
aatgagaaat taaacctgca agtgaaacat ttgaaacgat tatacttttc tacataagac 1260
atggttgagg catcagatac ttacaaagat ggtttaagta tggatactag agaaaattaa 1320
gttttcttcc tctttggttt attgatttgg ttaatttcc attatgctat tttgcataat 1380
caaggcactg taaatcttat aattttaaaa taaattactt aagaacagtt gtcattgtta 1440
tgttttgtta ttgattctca ttactgtcta attttttttc tggattagt ctcattttgt 1500
atgtatataa gttaaacaga tactgttttt aagtgcataa atagtacaag ttattatcaa 1560
ggatgtttta cagggaatc aaaagaatat tatcatactt tatctttcgt atgctgatta 1620
gtaaacgatt tttagacatt attttagaaa gtccataaat gtggaagaaa caaacagttg 1680
ctaccaaaag ttcttcaaat aaacatacaa ataaatgtgt atatttaagt ttttattgtt 1740
agcttctcca gaaaattgat gcaaatcttg gtaataattc ttgcattttt tccccataac 1800
ctggttaaaa taaatagccc attggcaata cttcataatg taatggaatt gtttggggaa 1860
cacttactgt accctctcat cctttttcca cttactgtg ttaacttagt gacatttaat 1920
gccaatatg tatgaataga tctaagccat ttaatttttt ttccttaaaa gattggagta 1980
ttttataatt caaggagcat acaaaacaat ggttgggaac atatgccaat tatggaatag 2040
gctatgtatt taatattaat ctctgccatt aggatatcta ctcactgtat aaacctcagt 2100
aaaaatagtg aagacatgca tcatggaatg agaaaatgag aaaggaatga gttgtctaac 2160
atcacagtgg gatctgtttt ttgtgaggtt catttctgaa cacattaggc atatgagcag 2220
atttccagtg aatctattta tgtttatttt ctgagtttca acgctgacct tttcttgcat 2280
tattgtttca ttttaatgat agtgttactt gtcccactgt tgttttcatt gagtttggat 2340
ttatatttta atgttcgaat gaaagtatga ttgtaaaagg gagtgaattg gtttaaaaaat 2400
atatgtatat tttaaacttt gttgtgtgta ggaacatga aggcattgta attcaatata 2460
aatgaccttt caattcatgg aatattaaag ttggtttaaa gtccaatagt taaaccttag 2520
caaaaatagc tttttacttc atcagttgct aagattttaa actttggatt catcaaagt 2580
tgacatgggc ttgtttgact ttctgtaagt ggcatttaag ttccacattc ttattacttg 2640
aggtacttta tactaacata agacagttag agtttagagg attacaagtt gctagtttat 2700
aatgtcttac taatgcagaa acaaggaaaa aagcaaaatt ggcctgaata ttctcttggg 2760
gaaagagggc accaaagaaa agggtaagtg catctgaggg ccaaaagaga tgtataagcc 2820

```

```

ttttagccca ttccccatgc tgggcctgct cacagagcca caggaagatc attcagaaac 2880
taggaaagga ggccccacaca gctgatcctg ccacagcaca cctgactcac tcggctctgt 2940
tagtgtaacc ttttaaagtgt agcaacacaa accctttccc tcttgctcagt tcaactcatcc 3000
tttggtttct ttttaatcac ctgtgtctgg gcacagacaa tcacaataaa tgcagccctt 3060
tattactgtt aaggatcata ctgttggttt ggagttggaa gggactact ctgtgattca 3120
gggtgtgtgt acccatattt ataattaggc tttattatct tcctaaatca aggaaaggaa 3180
atcatcccca gaccatttat gctgagcttt ggaatactat tttaaactgg attgtactta 3240
aataatgaag ctctgcatag aggaactagt cagaagtggg gaaaacactg tctaattttt 3300
atcagtctgg tataaagtat tgatctaaga gaactctccc tgtgcccctt ggtctttatt 3360
ctcaattaag aaaaacagtc acatgtcacg acaaaccaat caatctttat gagatattcc 3420
tgtatccata ccccgcttg tttgcaattt ataaacctcc cttcaaaac taaggagttg 3480
cagaaaaaaa tggatttcac agagccttgt gtccctaaag ttctgtccca gtcagcagtc 3540
tttatagtc aaacagatta taaaaaatgt tttccatttg aactttacag tttgcaaaag 3600
tgcttttata ctttttctaa tttcagaaac aggataattt gtttaagtggg tttcagtttg 3660
ctaataggga ttttttgtgt tttgtttttt aattttcagc atctcttgaa gaatcttgct 3720
acagccaaat ggcattctac tttttaaaga cgtttgcaat tattagttga ttcacagtac 3780
agaacaaggt ataaaggaaa aaaccctgct aggtagtgtt ataattgcta gattaaaaat 3840
agactagaac aggttcattt taagatttac ttggaagagc aaagaaggaa aaattatatt 3900
tttaaagaaa gagaatattc aggccttatt tctgggtatga agtttatatt ttttaaaaaa 3960
atcctatatt atcacaccag agattttaga tcttttctg gttagaaaca ttgctggtag 4020
ttggattata tttttattgt attcatttat cttaggggga acattgtaaa gaaacaaaaa 4080
ggtccagatg aatgtatgct agaaaataaaa gttgaaagat tcttacttca aaaaaaaaaa 4140
aaaaa

```

&lt;210&gt; 60

&lt;211&gt; 289

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

```

Met Thr Ser Lys Phe Leu Thr Ala Leu Ala Gln Asp Gly Val Ile Asn
 1             5             10             15

Glu Glu Ala Leu Ser Val Thr Glu Leu Asp Arg Val Tyr Gly Gly Leu
      20             25             30

Thr Thr Lys Val Gln Glu Ser Leu Lys Lys Gln Glu Gly Leu Leu Lys
 35             40             45

Asn Ile Gln Val Ser His Gln Glu Phe Ser Lys Met Lys Gln Ser Asn
 50             55             60

Asn Glu Ala Asn Leu Arg Glu Glu Val Leu Lys Asn Leu Ala Thr Ala
 65             70             75             80

Tyr Asp Asn Phe Val Glu Leu Val Ala Asn Leu Lys Glu Gly Thr Lys
      85             90             95

Phe Tyr Asn Glu Leu Thr Glu Ile Leu Val Arg Phe Gln Asn Lys Cys
 100            105            110

Ser Asp Ile Val Phe Ala Arg Lys Thr Glu Arg Asp Glu Leu Leu Lys
 115            120            125

Asp Leu Gln Gln Ser Ile Ala Arg Glu Pro Ser Ala Pro Ser Ile Pro
 130            135            140

Thr Pro Ala Tyr Gln Ser Ser Pro Ala Gly Gly His Ala Pro Thr Pro
 145            150            155            160

```

Pro Thr Pro Ala Pro Arg Thr Met Pro Pro Thr Lys Pro Gln Pro Pro  
 165 170 175  
 Ala Arg Pro Pro Pro Pro Val Leu Pro Ala Asn Arg Ala Pro Ser Ala  
 180 185 190  
 Thr Ala Pro Ser Pro Val Gly Ala Gly Thr Ala Ala Pro Ala Pro Ser  
 195 200 205  
 Gln Thr Pro Gly Ser Ala Pro Pro Pro Gln Ala Gln Gly Pro Pro Tyr  
 210 215 220  
 Pro Thr Tyr Pro Gly Tyr Pro Gly Tyr Cys Gln Met Pro Met Pro Met  
 225 230 235 240  
 Gly Tyr Asn Pro Tyr Ala Tyr Gly Gln Tyr Asn Met Pro Tyr Pro Pro  
 245 250 255  
 Val Tyr His Gln Ser Pro Gly Gln Ala Pro Tyr Pro Gly Pro Gln Gln  
 260 265 270  
 Pro Ser Tyr Pro Phe Pro Gln Pro Pro Gln Gln Ser Tyr Tyr Pro Gln  
 275 280 285

Gln

&lt;210&gt; 61

&lt;211&gt; 1417

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 61

```

ggtgccccgac atggcgagtg tagtgctgcc gagcggatcc cagtgtgcgg cggcagcggc 60
ggcggcgggcg cctcccgggc tccggctccg gcttctgctg ttgctcttct ccgccgcggc 120
actgatcccc acaggtgatg ggcagaatct gtttacgaaa gacgtgacag tgatcgaggg 180
agaggttgcg accatcagtt gccaaagtcaa taagagtgcac gactctgtga ttcagctact 240
gaatcccaac aggcagacca tttatttcag ggacttcagg cctttgaagg acagcaggtt 300
tcagttgctg aatttttcta gcagtgaact caaagtatca ttgacaaacg tctcaatttc 360
tgatgaagga agatactttt gccagctcta taccgatccc ccacaggaaa gttacaccac 420
catcacagtc ctggtcccac cacgtaatct gatgatcgat atccagaaa acactgcggg 480
ggaaggtgag gagattgaag tcaactgcac tgctatggcc agcaagccag ccacgactat 540
caggtgggttc aaagggaaca cagagctaaa aggcaaactc gaggtggaag agtgggtcaga 600
catgtacact gtgaccagtc agctgatgct gaaggtgcac aaggaggacg atgggggtccc 660
agtgatctgc caggtggagc accctgcggg cactggaaac ctgcagaccc agcggtatct 720
agaagtacag tataagcctc aagtgcacat tcagatgact tatcctctac aaggcttaac 780
ccgggaaggg gacgcgcttg agttaacatg tgaagccatc gggaagcccc agcctgtgat 840
ggtaacttgg gtgagagtcg atgatgaaat gcctcaacac gccgtactgt ctgggcccac 900
cctgttcac aataaccta acaaaacaga taatggtaca taccgctgtg aagcttcaaa 960
catagtgggg aaagctcact cggattatat gctgtatgta tacgattccc gagcaggtga 1020
agaaggctcg atcagggcag tggatcatgc cgtgatcggt ggcgtcgtgg cgggtggtgg 1080
gttcgccatg ctgtgcttgc tcatcattct ggggcgctat tttgccagac ataaagggtac 1140
atacttcact catgaagcca aaggagccga tgacgcagca gacgcagaca cagctataat 1200
caatgcagaa ggaggacaga acaactccga agaaaagaaa gagtacttca tctagatcag 1260
cctttttggt tcaatgaggt gtccaactgg ccctatttag atgataaaga gacagtga 1320
ttggaacttg cgagaaattc gtgtgttttt ttatgaatgg gtggaaaggt gtgagactgg 1380
gaaggcttgg gatttgctgt gtaaaaaaaaa aaaaaaa 1417

```

&lt;210&gt; 62

<211> 414  
 <212> PRT  
 <213> Homo sapiens

<400> 62

```

Met Ala Ser Val Val Leu Pro Ser Gly Ser Gln Cys Ala Ala Ala Ala
 1              5              10              15

Ala Ala Ala Ala Pro Pro Gly Leu Arg Leu Arg Leu Leu Leu Leu Leu
 20              25              30

Phe Ser Ala Ala Ala Leu Ile Pro Thr Gly Asp Gly Gln Asn Leu Phe
 35              40              45

Thr Lys Asp Val Thr Val Ile Glu Gly Glu Val Ala Thr Ile Ser Cys
 50              55              60

Gln Val Asn Lys Ser Asp Asp Ser Val Ile Gln Leu Leu Asn Pro Asn
 65              70              75              80

Arg Gln Thr Ile Tyr Phe Arg Asp Phe Arg Pro Leu Lys Asp Ser Arg
 85              90              95

Phe Gln Leu Leu Asn Phe Ser Ser Ser Glu Leu Lys Val Ser Leu Thr
 100             105             110

Asn Val Ser Ile Ser Asp Glu Gly Arg Tyr Phe Cys Gln Leu Tyr Thr
 115             120             125

Asp Pro Pro Gln Glu Ser Tyr Thr Thr Ile Thr Val Leu Val Pro Pro
 130             135             140

Arg Asn Leu Met Ile Asp Ile Gln Lys Asp Thr Ala Val Glu Gly Glu
 145             150             155             160

Glu Ile Glu Val Asn Cys Thr Ala Met Ala Ser Lys Pro Ala Thr Thr
 165             170             175

Ile Arg Trp Phe Lys Gly Asn Thr Glu Leu Lys Gly Lys Ser Glu Val
 180             185             190

Glu Glu Trp Ser Asp Met Tyr Thr Val Thr Ser Gln Leu Met Leu Lys
 195             200             205

Val His Lys Glu Asp Asp Gly Val Pro Val Ile Cys Gln Val Glu His
 210             215             220

Pro Ala Val Thr Gly Asn Leu Gln Thr Gln Arg Tyr Leu Glu Val Gln
 225             230             235             240

Tyr Lys Pro Gln Val His Ile Gln Met Thr Tyr Pro Leu Gln Gly Leu
 245             250             255

Thr Arg Glu Gly Asp Ala Leu Glu Leu Thr Cys Glu Ala Ile Gly Lys
 260             265             270

Pro Gln Pro Val Met Val Thr Trp Val Arg Val Asp Asp Glu Met Pro
 275             280             285

Gln His Ala Val Leu Ser Gly Pro Asn Leu Phe Ile Asn Asn Leu Asn

```

```
<210> 63
<211> 1571
<212> DNA
<213> Homo sapiens
```

<210> 64  
<211> 417

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 64

```

Met Pro Leu Tyr Glu Gly Leu Gly Ser Gly Gly Glu Lys Thr Ala Val
 1              5              10              15

Val Ile Asp Leu Gly Glu Ala Phe Thr Lys Cys Gly Phe Ala Gly Glu
          20              25              30

Thr Gly Pro Arg Cys Ile Ile Pro Ser Val Ile Lys Arg Ala Gly Met
          35              40              45

Pro Lys Pro Val Arg Val Val Gln Tyr Asn Ile Asn Thr Glu Glu Leu
 50              55              60

Tyr Ser Tyr Leu Lys Glu Phe Ile His Ile Leu Tyr Phe Arg His Leu
 65              70              75              80

Leu Val Asn Pro Arg Asp Arg Arg Val Val Ile Ile Glu Ser Val Leu
          85              90              95

Cys Pro Ser His Phe Arg Glu Thr Leu Thr Arg Val Leu Phe Lys Tyr
          100              105              110

Phe Glu Val Pro Ser Val Leu Leu Ala Pro Ser His Leu Met Ala Leu
 115              120              125

Leu Thr Leu Gly Ile Asn Ser Ala Met Val Leu Asp Cys Gly Tyr Arg
 130              135              140

Glu Ser Leu Val Leu Pro Ile Tyr Glu Gly Ile Pro Val Leu Asn Cys
 145              150              155              160

Trp Gly Ala Leu Pro Leu Gly Gly Lys Ala Leu His Lys Glu Leu Glu
          165              170              175

Thr Gln Leu Leu Glu Gln Cys Thr Val Asp Thr Ser Val Ala Lys Glu
          180              185              190

Gln Ser Leu Pro Ser Val Met Gly Ser Val Pro Glu Gly Val Leu Glu
          195              200              205

Asp Ile Lys Ala Arg Thr Cys Phe Val Ser Asp Leu Lys Arg Gly Leu
 210              215              220

Lys Ile Gln Ala Ala Lys Phe Asn Ile Asp Gly Asn Asn Glu Arg Pro
 225              230              235              240

Ser Pro Pro Pro Asn Val Asp Tyr Pro Leu Asp Gly Glu Lys Ile Leu
          245              250              255

His Ile Leu Gly Ser Ile Arg Asp Ser Val Val Glu Ile Leu Phe Glu
          260              265              270

Gln Asp Asn Glu Glu Gln Ser Val Ala Thr Leu Ile Leu Asp Ser Leu
          275              280              285

Ile Gln Cys Pro Ile Asp Thr Arg Lys Gln Leu Ala Glu Asn Leu Val
 290              295              300

```

Val Ile Gly Gly Thr Ser Met Leu Pro Gly Phe Leu His Arg Leu Leu  
 305 310 315 320

Ala Glu Ile Arg Tyr Leu Val Glu Lys Pro Lys Tyr Lys Lys Ala Leu  
 325 330 335

Gly Thr Lys Thr Phe Arg Ile His Thr Pro Pro Ala Lys Ala Asn Cys  
 340 345 350

Val Ala Trp Leu Gly Gly Ala Ile Phe Gly Ala Leu Gln Asp Ile Leu  
 355 360 365

Gly Ser Arg Ser Val Ser Lys Glu Tyr Tyr Asn Gln Thr Gly Arg Ile  
 370 375 380

Pro Asp Trp Cys Ser Leu Asn Asn Pro Pro Leu Glu Met Met Phe Asp  
 385 390 395 400

Val Gly Lys Thr Gln Pro Pro Leu Met Lys Arg Ala Phe Ser Thr Glu  
 405 410 415

Lys

&lt;210&gt; 65

&lt;211&gt; 1752

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 65

```

ggccaatcag agggacggcc ccagaatggc atggtagatg gaacgcagct gagaggtctg 60
acaagatgta ccaggtccca ctaccactgg atcgggatgg gaccctggta cggtccgct 120
tcaccatggg ggccctggtc acgggtctgct gtccacttgt cgccttcctc ttctgcatcc 180
tctgggtccct gctcttcac ttcaaggaga caacggccac aactgtggg gtgcccatt 240
acctgcccctc ggtgagctca gccatcggcg gggaggtgcc ccagcgctac gtgtggcggt 300
tctgcatcgg cctgcaactcg gcgcctcgct tcttgggtggc cttcgctac tgggaacct 360
acctcagctg cacctccccg tgttctgct atcgcccgct ctgccgcctc aacttcggcc 420
tcaatgtcgt ggagaacctc gcgttgctag tgtcactta tgtctcctcc tccgaggact 480
tcaccatcca cgaaaatgct ttcatttgtt tcattgcctc atccctcggg cacatgctcc 540
tcacctgcat tctctggcgg ttgaccaaga agcacacagt aagtcaggag gatcgcaagt 600
cctacagctg gaaacagcgg ctcttcatca tcaacttcat ctcttcttc tcggcgctgg 660
ctgtctactt tcggcacaac atgtattgtg aggtctggagt gtacaccatc tttgccatcc 720
tggagtacac tgttgtctta accaatatgg cgttccacat gacggcctgg tgggacttcg 780
ggaacaagga gctgctcata acctctcagc ctgaggaaaa gcgattctga acccttcagt 840
cctgcttggg aggacgcagc ccaactgcca gaaacaagaa acacgatacc attctggcct 900
tccccacccc acatcctctc ttggccttac tgaagatggg ggaagggtaa gaaggaagg 960
tgtaggccaa ggctcacccc agtgctgctg gcttctcctc tccacccctc atatggcggt 1020
ggggtcctca aacatcacct ttacctgaga ggccccaaga agctgagctg gcagagagct 1080
ccaccatttg gtgctaaaaa aaaaaacgtc ctgaggttca tgaccaccat ccagtttctg 1140
gcctttacac agtcaccttt cactgaggtc aggagccctc gagcagtggc tgctccctga 1200
caaccacagc catttctctg cacgggggtc attcatagga ctaatgtatt tcatgatcta 1260
ctgtgcacat ccaggcctgt ggccacagtc ccctgctaaa gttgctcagg tgttctagtc 1320
ctgacttcac ctttttgatt tgggtgtgtg cctagggtat gtacccttcc ccatctgagc 1380
ctcgggtgtg ccatgtgtct ggcggggat ggggtgactg tatgatttcc aaggactcta 1440
ccagtcagtg gttctgatgt catcggttg aggtggtgt ctatacctaa aggatgacct 1500
gtccagaaa cagcaccagc acagcatgta ttttcttctc ttctgaaagt tctggcttgt 1560
agacccctcc cctcctttgc aaaggatagg gatagagggg tcagatgcag atctctactg 1620
taaaatgggc tccctgggtat ctctgtctt ccctactgct ccaaacccta aattttggtt 1680

```

gtacatttta tttgaaagga aaataaattt tttttttggg ccaaaaaaaaaa aaaaaaaaaa 1740  
 aaaaaaaaaa aa 1752

<210> 66  
 <211> 254  
 <212> PRT  
 <213> Homo sapiens

<400> 66  
 Met Tyr Gln Val Pro Leu Pro Leu Asp Arg Asp Gly Thr Leu Val Arg  
 1 5 10 15  
 Leu Arg Phe Thr Met Val Ala Leu Val Thr Val Cys Cys Pro Leu Val  
 20 25 30  
 Ala Phe Leu Phe Cys Ile Leu Trp Ser Leu Leu Phe His Phe Lys Glu  
 35 40 45  
 Thr Thr Ala Thr His Cys Gly Val Pro Asn Tyr Leu Pro Ser Val Ser  
 50 55 60  
 Ser Ala Ile Gly Gly Glu Val Pro Gln Arg Tyr Val Trp Arg Phe Cys  
 65 70 75 80  
 Ile Gly Leu His Ser Ala Pro Arg Phe Leu Val Ala Phe Ala Tyr Trp  
 85 90 95  
 Asn His Tyr Leu Ser Cys Thr Ser Pro Cys Ser Cys Tyr Arg Pro Leu  
 100 105 110  
 Cys Arg Leu Asn Phe Gly Leu Asn Val Val Glu Asn Leu Ala Leu Leu  
 115 120 125  
 Val Leu Thr Tyr Val Ser Ser Ser Glu Asp Phe Thr Ile His Glu Asn  
 130 135 140  
 Ala Phe Ile Val Phe Ile Ala Ser Ser Leu Gly His Met Leu Leu Thr  
 145 150 155 160  
 Cys Ile Leu Trp Arg Leu Thr Lys Lys His Thr Val Ser Gln Glu Asp  
 165 170 175  
 Arg Lys Ser Tyr Ser Trp Lys Gln Arg Leu Phe Ile Ile Asn Phe Ile  
 180 185 190  
 Ser Phe Phe Ser Ala Leu Ala Val Tyr Phe Arg His Asn Met Tyr Cys  
 195 200 205  
 Glu Ala Gly Val Tyr Thr Ile Phe Ala Ile Leu Glu Tyr Thr Val Val  
 210 215 220  
 Leu Thr Asn Met Ala Phe His Met Thr Ala Trp Trp Asp Phe Gly Asn  
 225 230 235 240  
 Lys Glu Leu Leu Ile Thr Ser Gln Pro Glu Glu Lys Arg Phe  
 245 250

<210> 67  
 <211> 781



&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 67

```

cactcctgca gacaaggcac tgattgcccc agaccatgta gttccagctc cagaagagtg 60
ctatgtgtat agtccattgg gctctgctta taaacttcaa agttacactg aaggatacgg 120
taaaaacacc agtttagtaa ccatttttat gatttggaat accatgatgg gaacatctat 180
actaagcatt ccttggggca taaaacaggc tggatttact actggaatgt gtgtcatcat 240
actgatgggc cttttaacac tttattgctg ctacagagta gtgaaatcac ggactatgat 300
gttttcattg gataccacta cctgggaata tccagatgtc tgcagacatt atttcggctc 360
ctttggggcag tggtcgagtc tcctcttctc cttgggtgtc ctcattggag caatgatagt 420
ttattgggtg cttatgtcaa attttctttt taatactgga aagtttattt ttagtaagta 480
tctatatcat atgcttttaa cacagtactt tcaaatacta ttaccactgt aatgttagtt 540
ctagccttaa attctaggac ttgggataaa taaaataaga agtaacatat ataatttttg 600
aaaatatatt ttattcagtt ggctttctgt ggttgtgctc tcaaataatag tgtatgctta 660
tttccaaaca ttaatctttg aaggaataat attcctccaa aatctttagt taaaataaaa 720
tatgtctata atccaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 780
a

```

781

&lt;210&gt; 68

&lt;211&gt; 127

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

```

Met Ile Trp Asn Thr Met Met Gly Thr Ser Ile Leu Ser Ile Pro Trp
  1                      5                      10                      15

Gly Ile Lys Gln Ala Gly Phe Thr Thr Gly Met Cys Val Ile Ile Leu
                20                      25                      30

Met Gly Leu Leu Thr Leu Tyr Cys Cys Tyr Arg Val Val Lys Ser Arg
  35                      40                      45

Thr Met Met Phe Ser Leu Asp Thr Thr Thr Trp Glu Tyr Pro Asp Val
  50                      55                      60

Cys Arg His Tyr Phe Gly Ser Phe Gly Gln Trp Ser Ser Leu Leu Phe
  65                      70                      75                      80

Ser Leu Val Ser Leu Ile Gly Ala Met Ile Val Tyr Trp Val Leu Met
                85                      90                      95

Ser Asn Phe Leu Phe Asn Thr Gly Lys Phe Ile Phe Ser Lys Tyr Leu
  100                      105                      110

Tyr His Met Leu Leu Thr Gln Tyr Phe Gln Ile Leu Leu Pro Leu
  115                      120                      125

```

&lt;210&gt; 69

&lt;211&gt; 649

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 69

```

gagcaactcc cttccccatc tctgctcacc atgtggacgc tgaaatcgct cctggtcctg 60
cttctgtgcc tcacctgcag ctatgccttt atgttctctt ctctgagaca gaaaactagc 120
gaaccccgagg ggaaggtgca atacggagag cactttcgga ttcggcgagaa tctaccagag 180
cacacccaag gctggccttg gagcaaatgg ctctggcttc tttttgttgt tgtgccgttt 240

```

```

gtgatactgc agtgtcaaag agacagtgcg aagaataagg agcagagtcc tcttggcctt 300
cgaggcggcc aacttcactc tccattaaag aaaaaaagaa atgcttcccc caacaaagac 360
tgtgcattca ataccttaat ggaactcgag gtggagctta tgaaatttgt gtccaaagtg 420
cggaatctta aacgtgccat ggcaacaggt agtggcagta acctcaggct tcgaaagtca 480
gagatgcctg cagatccata ccatgtcacg atctgtgaaa tatggggaga agaaagctct 540
agctgaatgg atttgtgtgt caggagagaa aaaagttagg tgttgacaaa ctgtatgcaa 600
actaataaaa ctattctgaa gaaaagaaaa aaaaaaaaaa aaaaaaaaaa 649

```

&lt;210&gt; 70

&lt;211&gt; 171

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 70

```

Met Trp Thr Leu Lys Ser Ser Leu Val Leu Leu Leu Cys Leu Thr Cys
  1                      5                      10                      15
Ser Tyr Ala Phe Met Phe Ser Ser Leu Arg Gln Lys Thr Ser Glu Pro
          20                      25                      30
Gln Gly Lys Val Gln Tyr Gly Glu His Phe Arg Ile Arg Gln Asn Leu
          35                      40                      45
Pro Glu His Thr Gln Gly Trp Leu Gly Ser Lys Trp Leu Trp Leu Leu
          50                      55                      60
Phe Val Val Val Pro Phe Val Ile Leu Gln Cys Gln Arg Asp Ser Glu
          65                      70                      75                      80
Lys Asn Lys Glu Gln Ser Pro Pro Gly Leu Arg Gly Gly Gln Leu His
          85                      90                      95
Ser Pro Leu Lys Lys Lys Arg Asn Ala Ser Pro Asn Lys Asp Cys Ala
          100                     105                     110
Phe Asn Thr Leu Met Glu Leu Glu Val Glu Leu Met Lys Phe Val Ser
          115                     120                     125
Lys Val Arg Asn Leu Lys Arg Ala Met Ala Thr Gly Ser Gly Ser Asn
          130                     135                     140
Leu Arg Leu Arg Lys Ser Glu Met Pro Ala Asp Pro Tyr His Val Thr
          145                     150                     155                     160
Ile Cys Glu Ile Trp Gly Glu Glu Ser Ser Ser
          165                     170

```

&lt;210&gt; 71

&lt;211&gt; 1456

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

```

cacggctgtc ttatctgcaa gtgcagagag gcctctgctt cagctggggc acccatcctg 60
tcgggcactt gtctcaccgt ggatggtcat catcataaaa atgaggagag ctggcacgat 120
gggtgcccgg aatgctactg tctcaatgga cgggaaatgt gtgccctgat cacctgcccg 180
gtgcctgcct gtggcaaccc caccattcac cctggacagt gctgcccata atgtgcagat 240
gactttgttg tgcagaagcc agagctcagt actccctcca ttgcccacgc ccctggagga 300
gaatactttg tggaaggaga aacgtggaac attgactcct gtactcagtg cacctgccac 360

```

```

agcgggacggg tgctgtgtga gacagaggtg tgcccaccgc tgctctgcca gaaccctca 420
cgcacccagg attcctgctg cccacagtgt acagatcaac ctttctggcc ttccttgtcc 480
cgcaataaca gcgtacctaa ttattgcaaa aatgatgaag gggatatatt cctggcagct 540
gagtcctgga agcctgacgt ttgtaccagc tgcattctgca ttgatagcgt aattagctgt 600
ttctctgagt cctgcccttc tgtatcctgt gaaagacctg tcttgagaaa aggccagtgt 660
tgtccctact gcatagaaga cacaattcca aagaaggtgg tgtgccactt cagtgggaag 720
gcctatgccg acgaggagcg gtgggacctt gacagctgca cccactgcta ctgcctgcag 780
ggccagaccc tctgctcgac cgtcagctgc cccctctgc cctgtgttga gcccatcaac 840
gtggaaggaa gttgctgccc aatgtgtcca gaaatgtatg tcccagaacc aaccaatata 900
cccattgaga agacaaacca tcgaggagag gttgacctgg aggttcccc gtggcccacg 960
cctagtga aa atgatatcgt ccatctccct agagatatgg gtcacctcca ggtagattac 1020
agagataaca ggctgcaccc aagtgaagat tcttcaactg actccattgc ctcagtgtgtg 1080
gttcccataa ttatatgcct ctctattata atagcattcc tattcatcaa tcagaagaaa 1140
cagtggatac cactgctttg ctggtatcga acaccaacta agccttcttc cttaaataat 1200
cagctagtat ctgtggactg caagaaagga accagagtcc aggtggacag tcccagaga 1260
atgctaagaa ttgcagaacc agatgcaaga ttcagtggct tctacagcat gcaaaaacag 1320
aaccatctac aggcagacaa tttctaccaa acagtgtgaa gaaaggcaac taggatgagg 1380
tttcaaaaaga cggaagacga ctaaattctgc tctaaaaagt aaactagaat ttgtgcactt 1440
aaaaaaaaa aaaaaa 1456

```

&lt;210&gt; 72

&lt;211&gt; 400

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 72

```

Met Cys Ala Leu Ile Thr Cys Pro Val Pro Ala Cys Gly Asn Pro Thr
  1                      5                      10                      15

```

```

Ile His Pro Gly Gln Cys Cys Pro Ser Cys Ala Asp Asp Phe Val Val
                20                      25                      30

```

```

Gln Lys Pro Glu Leu Ser Thr Pro Ser Ile Cys His Ala Pro Gly Gly
    35                      40                      45

```

```

Glu Tyr Phe Val Glu Gly Glu Thr Trp Asn Ile Asp Ser Cys Thr Gln
    50                      55                      60

```

```

Cys Thr Cys His Ser Gly Arg Val Leu Cys Glu Thr Glu Val Cys Pro
    65                      70                      75                      80

```

```

Pro Leu Leu Cys Gln Asn Pro Ser Arg Thr Gln Asp Ser Cys Cys Pro
                85                      90                      95

```

```

Gln Cys Thr Asp Gln Pro Phe Arg Pro Ser Leu Ser Arg Asn Asn Ser
    100                      105                      110

```

```

Val Pro Asn Tyr Cys Lys Asn Asp Glu Gly Asp Ile Phe Leu Ala Ala
    115                      120                      125

```

```

Glu Ser Trp Lys Pro Asp Val Cys Thr Ser Cys Ile Cys Ile Asp Ser
    130                      135                      140

```

```

Val Ile Ser Cys Phe Ser Glu Ser Cys Pro Ser Val Ser Cys Glu Arg
    145                      150                      155                      160

```

```

Pro Val Leu Arg Lys Gly Gln Cys Cys Pro Tyr Cys Ile Glu Asp Thr
                165                      170                      175

```

```

Ile Pro Lys Lys Val Val Cys His Phe Ser Gly Lys Ala Tyr Ala Asp

```

```
<210> 73
<211> 4723
<212> DNA
<213> Homo sapiens
```

65

gcctcttgaa	gccagggcag	gcagagcagg	gcaaaagcca	ggagcagga	cgtccgggag	780
cctggagcca	ttgccactag	gtgagctgtc	cacaggaccc	tgagtgggtc	ggggagttcg	840
gccttcattg	cctaggagcg	gcgcaggagt	gagcgagcg	gggcgcgcgg	agcggagccc	900
gcggatcttg	tgctgcgcca	ccgcgcccac	tcggcagctc	gggaggcggg	gaccggcccc	960
gaggctgcgc	cgctgcgggg	ccggccgact	cggaggagga	gagggaggag	gcgcgcgcgg	1020
cccgggctgg	agccgagcgc	agcagccacc	gccgcccgcc	cgccagaagt	ttgggttgaa	1080
ccggagctgc	cgggaggaaa	cttttttctt	ttttccccct	ccctccccgg	aggaggagga	1140
ggaggaggag	gggaagctgc	cgccggcgcc	aaggctcgtg	ggctcggggg	cggcgcggcc	1200
cgcagaaggg	gcgggggcct	cgccccgcga	ggggaggcgc	gccccggggg	ccccgagagg	1260
ggcggtgagg	accgcgggct	gctggtgcgg	cggcggcggc	ggcgcggtgt	ccccgcgcag	1320
gggaggggcg	ccgccccgct	cccgccccgg	ctgcgaggag	gaggcggcgg	cggcgcagga	1380
ggatgtactt	ggtggcgggg	gacagggggg	tgcccggtcg	cgggcacctc	ctggtctcgc	1440
tgctggggct	gctgctgctg	ctggcgcgct	ccggcacccg	ggcgctggtc	tgcttgcctt	1500
gtgacgagtc	caagtgcgag	gagcccagga	actgcccggg	gagcatcgtg	cagggcgctt	1560
gcggctgctg	ctacacgtgc	gccagccaga	ggaacgagag	ctgcggcgcc	accttcggga	1620
tttacggaac	ctgcgaccgg	gggctgcgtt	gtgtcatccg	ccccccgctc	aatggcgact	1680
ccctcaccga	gtacgaagcg	ggcgtttgcg	aagatgagaa	ctggactgat	gaccaactgc	1740
ttggttttaa	accatgcaat	gaaaacctta	ttgctggctg	caatataatc	aatgggaaat	1800
gtgaatgtaa	caccattcga	acctgcagca	atccctttga	gtttccaaat	caggatatgt	1860
gcctttcagc	tttaaagaga	attgaagaag	agaagccaga	ttgctccaag	gcccgcgtgt	1920
aagtccagtt	ctctccacgt	tgctctgaag	attctgttct	gatcgagggt	tatgctcctc	1980
ctggggagtg	ctgtccctta	cccagccgct	gcgtgtgcaa	ccccgcaggc	tgtctgcgca	2040
aagtctgcca	gccgggaaac	ctgaacatac	tagtgtcaaa	agcctcaggg	aagccgggag	2100
agtgtgtgta	cctctatgag	tgcaaaccag	ttttcggcgt	ggactgcagg	actgtggaat	2160
gccctctgtg	tcagcagacc	gcgtgtcccc	cggacagcta	tgaaactcaa	gtcagactaa	2220
ctgcagatgg	tgctgtactt	tgccaacaaa	gatcgagtg	tctctctggc	ttatgtgggt	2280
tccccgtgtg	tgagggtggga	tccactcccc	gcatagtctc	tcgtggcgat	gggacacctg	2340
gaaagtgtgt	tgatgtcttt	gaatgtgtta	atgatacaaa	gccagcctgc	gtattttaaca	2400
atgtggaata	ttatgatgga	gacatgtttc	gaatggacaa	ctgtcgggtc	tgctgatgcc	2460
aagggggcgt	tgccatctgc	ttcactgccc	agtgtgggtg	gataaaactgc	gagagggtact	2520
acgtgcccga	aggagagtg	tgcccagtg	gtgaagatcc	agtgtatcct	tttaataatc	2580
ccgctggctg	ctatgccaat	ggcctgatcc	ttgcccacgg	agaccgggtg	cggaagacg	2640
actgcacatt	gtcccagtg	gtcaacggg	acgccaactg	cgttgcgacc	gtctgcggac	2700
agacctgcac	aaacctgtg	aaagtgcctg	gggagtgttg	ccctgtgtgc	gaagaaccaa	2760
ccatcatcac	agttgatcca	cctgcatgtg	gggagttatc	aaactgcact	ctgacaggga	2820
aggactgcat	taatgggttc	aaacgcgatc	acaatgggtg	tcggacctgt	cagtgcataa	2880
acaccgagga	actatgttca	gaacgtaaac	aaggctgcac	cttgaactgt	cccttcgggt	2940
tccttactga	tgcccaaaac	tgtgagatct	gtgagtgcgc	cccaaggccc	aagaagtgca	3000
gaccataaat	ctgtgacaag	tattgtccac	ttggattgct	gaagaataag	cacggctgtg	3060
acatctgtcg	ctgtaagaaa	tgtccagagc	tctcatgcag	taagatctgc	cccttggggt	3120
tccagcagga	cagtcacggc	tgtcttatct	gcaagtgacg	agaggcctct	gcctcagctg	3180
ggccacccat	cctgtcgggc	acttgtctca	ccgtggatgg	tcacatcat	aaaaatgagg	3240
agagctggca	cgatgggtgc	cgggaatgct	actgtctcaa	tggaacggga	atgtgtgccc	3300
tgatcacctg	cccgggtgct	gcctgtggca	acccaccat	tcaccctgga	cagtgtgccc	3360
catcatgtgc	agatgacttt	gtggtgcaga	agccagagct	cagtactccc	tccatttgcc	3420
acgcccttg	aggagaatac	tttgtggaag	gagaaacgtg	gaacattgac	tcctgtactc	3480
agtgcacctg	ccacagcgga	cgggtgctgt	gtgagacaga	ggtgtgcccc	ccgtgtctct	3540
gccagaaccc	ctcacgcacc	caggattcct	gctgccaca	gtgtacagat	caaccttttc	3600
ggccttcctt	gtcccgaat	aacagcgta	ctaattactg	caaaaatgat	gaaggggata	3660
tattcctggc	agctgagtc	tggaagcctg	acgtttgtac	cagctgcata	tgcatgtata	3720
gcgtaattag	ctgtttctct	gagtcctgcc	cttctgtatc	ctgtgaaaga	cctgtcttga	3780
gaaaaggcca	gtgttgtccc	tactgcatag	aagacacaa	tccaaagaag	gtggtgtgcc	3840
acttcagtgg	gaaggcctat	gccgacgagg	agcgggtggg	ccttgacagc	tgacccact	3900
gctactgcct	gcagggccag	accctctgct	cgaccgtcag	ctgccccctt	ctgccctgtg	3960
ttgagcccat	caacgtggaa	ggaagtgtgt	gcccaatgtg	tccagaaatg	tatgtccccc	4020
aaccaaccaa	tataccatt	gagaagacaa	accatcgagg	agaggttgac	ctggaggttc	4080
ccctgtggcc	cacgcctagt	gaaaatgata	tcgtccatct	ccctagagat	atgggtcacc	4140
tccaggtaga	ttacagagat	aacaggctgc	acccaagtga	agattcttca	ctggactcca	4200
ttgcctcagt	tgtggttccc	ataattatat	gcctctctat	tataatagca	ttcctattca	4260
tcaatcagaa	gaaacagtgg	ataccactgc	tttctgtgta	tcgaacacca	actaagcctt	4320

```

cttccttaaa taatcagtta gtatctgtgg actgcaagaa aggaaccaga gtccaggtgg 4380
acagttccca gagaatgcta agaattgcag aaccagatgc aagattcagt ggcttctaca 4440
gcatgcaaaa acagaaccat ctacaggcag acaatttcta ccaaacagtg tgaagaaagg 4500
caactaggat gaggtttcaa aagacggaag acgactaaat ctgctctaaa aagtaaacta 4560
gaatttgtgc acttgcttag tggattgtat tggattgtga cttgatgtac agcgctaaga 4620
ccttactggg atgggctctg tctacagcaa tgtgcagaac aagcattccc cctcaaacct 4680
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa                      4723

```

&lt;210&gt; 74

&lt;211&gt; 1036

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 74

```

Met Tyr Leu Val Ala Gly Asp Arg Gly Leu Ala Gly Cys Gly His Leu
  1                      5                      10                      15

Leu Val Ser Leu Leu Gly Leu Leu Leu Leu Leu Ala Arg Ser Gly Thr
      20                      25                      30

Arg Ala Leu Val Cys Leu Pro Cys Asp Glu Ser Lys Cys Glu Glu Pro
      35                      40                      45

Arg Asn Cys Pro Gly Ser Ile Val Gln Gly Val Cys Gly Cys Cys Tyr
      50                      55                      60

Thr Cys Ala Ser Gln Arg Asn Glu Ser Cys Gly Gly Thr Phe Gly Ile
      65                      70                      75                      80

Tyr Gly Thr Cys Asp Arg Gly Leu Arg Cys Val Ile Arg Pro Pro Leu
      85                      90                      95

Asn Gly Asp Ser Leu Thr Glu Tyr Glu Ala Gly Val Cys Glu Asp Glu
      100                     105                     110

Asn Trp Thr Asp Asp Gln Leu Leu Gly Phe Lys Pro Cys Asn Glu Asn
      115                     120                     125

Leu Ile Ala Gly Cys Asn Ile Ile Asn Gly Lys Cys Glu Cys Asn Thr
      130                     135                     140

Ile Arg Thr Cys Ser Asn Pro Phe Glu Phe Pro Ser Gln Asp Met Cys
      145                     150                     155                     160

Leu Ser Ala Leu Lys Arg Ile Glu Glu Glu Lys Pro Asp Cys Ser Lys
      165                     170                     175

Ala Arg Cys Glu Val Gln Phe Ser Pro Arg Cys Pro Glu Asp Ser Val
      180                     185                     190

Leu Ile Glu Gly Tyr Ala Pro Pro Gly Glu Cys Cys Pro Leu Pro Ser
      195                     200                     205

Arg Cys Val Cys Asn Pro Ala Gly Cys Leu Arg Lys Val Cys Gln Pro
      210                     215                     220

Gly Asn Leu Asn Ile Leu Val Ser Lys Ala Ser Gly Lys Pro Gly Glu
      225                     230                     235                     240

Cys Cys Asp Leu Tyr Glu Cys Lys Pro Val Phe Gly Val Asp Cys Arg

```

245										250					255				
Thr	Val	Glu	Cys	Pro	Pro	Val	Gln	Gln	Thr	Ala	Cys	Pro	Pro	Asp	Ser				
			260						265					270					
Tyr	Glu	Thr	Gln	Val	Arg	Leu	Thr	Ala	Asp	Gly	Cys	Cys	Thr	Leu	Pro				
		275					280						285						
Thr	Arg	Cys	Glu	Cys	Leu	Ser	Gly	Leu	Cys	Gly	Phe	Pro	Val	Cys	Glu				
		290				295					300								
Val	Gly	Ser	Thr	Pro	Arg	Ile	Val	Ser	Arg	Gly	Asp	Gly	Thr	Pro	Gly				
305					310					315					320				
Lys	Cys	Cys	Asp	Val	Phe	Glu	Cys	Val	Asn	Asp	Thr	Lys	Pro	Ala	Cys				
				325					330					335					
Val	Phe	Asn	Asn	Val	Glu	Tyr	Tyr	Asp	Gly	Asp	Met	Phe	Arg	Met	Asp				
			340					345						350					
Asn	Cys	Arg	Phe	Cys	Arg	Cys	Gln	Gly	Gly	Val	Ala	Ile	Cys	Phe	Thr				
		355					360						365						
Ala	Gln	Cys	Gly	Glu	Ile	Asn	Cys	Glu	Arg	Tyr	Tyr	Val	Pro	Glu	Gly				
		370				375						380							
Glu	Cys	Cys	Pro	Val	Cys	Glu	Asp	Pro	Val	Tyr	Pro	Phe	Asn	Asn	Pro				
385					390					395					400				
Ala	Gly	Cys	Tyr	Ala	Asn	Gly	Leu	Ile	Leu	Ala	His	Gly	Asp	Arg	Trp				
				405					410					415					
Arg	Glu	Asp	Asp	Cys	Thr	Phe	Cys	Gln	Cys	Val	Asn	Gly	Glu	Arg	His				
			420					425					430						
Cys	Val	Ala	Thr	Val	Cys	Gly	Gln	Thr	Cys	Thr	Asn	Pro	Val	Lys	Val				
		435					440					445							
Pro	Gly	Glu	Cys	Cys	Pro	Val	Cys	Glu	Glu	Pro	Thr	Ile	Ile	Thr	Val				
		450				455						460							
Asp	Pro	Pro	Ala	Cys	Gly	Glu	Leu	Ser	Asn	Cys	Thr	Leu	Thr	Gly	Lys				
465					470					475					480				
Asp	Cys	Ile	Asn	Gly	Phe	Lys	Arg	Asp	His	Asn	Gly	Cys	Arg	Thr	Cys				
				485					490					495					
Gln	Cys	Ile	Asn	Thr	Glu	Glu	Leu	Cys	Ser	Glu	Arg	Lys	Gln	Gly	Cys				
			500					505					510						
Thr	Leu	Asn	Cys	Pro	Phe	Gly	Phe	Leu	Thr	Asp	Ala	Gln	Asn	Cys	Glu				
		515					520						525						
Ile	Cys	Glu	Cys	Arg	Pro	Arg	Pro	Lys	Lys	Cys	Arg	Pro	Ile	Ile	Cys				
		530				535					540								
Asp	Lys	Tyr	Cys	Pro	Leu	Gly	Leu	Leu	Lys	Asn	Lys	His	Gly	Cys	Asp				
545					550					555					560				
Ile	Cys	Arg	Cys	Lys	Lys	Cys	Pro	Glu	Leu	Ser	Cys	Ser	Lys	Ile	Cys				

	565		570		575
Pro Leu Gly Phe Gln Gln Asp Ser His Gly Cys Leu Ile Cys Lys Cys	580		585		590
Arg Glu Ala Ser Ala Ser Ala Gly Pro Pro Ile Leu Ser Gly Thr Cys	595		600		605
Leu Thr Val Asp Gly His His His Lys Asn Glu Glu Ser Trp His Asp	610		615		620
Gly Cys Arg Glu Cys Tyr Cys Leu Asn Gly Arg Glu Met Cys Ala Leu	625		630		635
Ile Thr Cys Pro Val Pro Ala Cys Gly Asn Pro Thr Ile His Pro Gly		645		650	655
Gln Cys Cys Pro Ser Cys Ala Asp Asp Phe Val Val Gln Lys Pro Glu		660		665	670
Leu Ser Thr Pro Ser Ile Cys His Ala Pro Gly Gly Glu Tyr Phe Val		675		680	685
Glu Gly Glu Thr Trp Asn Ile Asp Ser Cys Thr Gln Cys Thr Cys His		690		695	700
Ser Gly Arg Val Leu Cys Glu Thr Glu Val Cys Pro Pro Leu Leu Cys		705		710	715
Gln Asn Pro Ser Arg Thr Gln Asp Ser Cys Cys Pro Gln Cys Thr Asp		725		730	735
Gln Pro Phe Arg Pro Ser Leu Ser Arg Asn Asn Ser Val Pro Asn Tyr		740		745	750
Cys Lys Asn Asp Glu Gly Asp Ile Phe Leu Ala Ala Glu Ser Trp Lys		755		760	765
Pro Asp Val Cys Thr Ser Cys Ile Cys Ile Asp Ser Val Ile Ser Cys		770		775	780
Phe Ser Glu Ser Cys Pro Ser Val Ser Cys Glu Arg Pro Val Leu Arg		785		790	795
Lys Gly Gln Cys Cys Pro Tyr Cys Ile Glu Asp Thr Ile Pro Lys Lys		805		810	815
Val Val Cys His Phe Ser Gly Lys Ala Tyr Ala Asp Glu Glu Arg Trp		820		825	830
Asp Leu Asp Ser Cys Thr His Cys Tyr Cys Leu Gln Gly Gln Thr Leu		835		840	845
Cys Ser Thr Val Ser Cys Pro Pro Leu Pro Cys Val Glu Pro Ile Asn		850		855	860
Val Glu Gly Ser Cys Cys Pro Met Cys Pro Glu Met Tyr Val Pro Glu		865		870	875
					880
Pro Thr Asn Ile Pro Ile Glu Lys Thr Asn His Arg Gly Glu Val Asp					



	885		890		895
Leu Glu Val	Pro Leu Trp Pro Thr	Pro Ser Glu Asn Asp	Ile Val His		
	900	905	910		
Leu Pro Arg Asp Met Gly His	Leu Gln Val Asp Tyr Arg Asp Asn Arg				
	915	920	925		
Leu His Pro Ser Glu Asp Ser Ser	Leu Asp Ser Ile Ala Ser Val Val				
	930	935	940		
Val Pro Ile Ile Ile Cys Leu Ser	Ile Ile Ile Ala Phe Leu Phe Ile				
	945	950	955	960	
Asn Gln Lys Lys Gln Trp Ile Pro	Leu Leu Cys Trp Tyr Arg Thr Pro				
	965	970	975		
Thr Lys Pro Ser Ser Leu Asn Asn	Gln Leu Val Ser Val Asp Cys Lys				
	980	985	990		
Lys Gly Thr Arg Val Gln Val Asp	Ser Ser Gln Arg Met Leu Arg Ile				
	995	1000	1005		
Ala Glu Pro Asp Ala Arg Phe Ser	Gly Phe Tyr Ser Met Gln Lys Gln				
	1010	1015	1020		
Asn His Leu Gln Ala Asp Asn Phe	Tyr Gln Thr Val				
	1025	1030	1035		

&lt;210&gt; 75

&lt;211&gt; 3861

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 75

```

gtgcacgcgt ggcagacgga gaaggccagt gcccagcttg aaggttctgt caccttttgc 60
agtgggtccaa atgagaaaaa agtggaaaaat gggaggcatg aaatacatct ttctgttggt 120
gttcttttctt ttgctagaag gaggcaaaac agagcaagta aaacattcag agacatattg 180
catgttttcaa gacaagaagt acagagtggg tgagagatgg catccttacc tggaaacctta 240
tgggttggtt tactgcgtga actgcatctg ctacagagaat gggaatgtgc ttgcagccg 300
agtcagatgt ccaaagtgtt attgcctttc tcctgtgcat attcctcatc tgtgctgccc 360
tcgctgccc aactccttac cccagtgaa caataagtg accagcaagt cttgcgagta 420
caatgggaca acttaccac atggagagct gtctgtagct gaagggtct ttcagaatcg 480
gcaacccaat caatgcaccc agtcagctg ttccggaggga aacgtgtatt gtgggtctcaa 540
gacttgcccc aaattaacct gtgccttccc agtctctgtt ccagattcct gctgccgggt 600
atgcagagga gatggagaac tgtcatggga acattctgat ggtgatctt tccggcaacc 660
tgccaacaga gaagcaagac attcttacc cgcctctcac tatgatctc caccaagccg 720
acaggctgga ggtctgtccc gctttcctgg ggccagaagt caccggggag ctcttatgga 780
ttcccagcaa gcatcaggaa ccattgtgca aattgtcatc aataacaaac acaagcatgg 840
acaagtgtgt gtttccaatg gaaagacctt ttctcatggc gagtcttgcc acccaaacct 900
ccgggcattt ggcattgtgg agtgtgtgct atgtacttgt aatgtcacca agcaagagt 960
taagaaaatc cactgcccc aatcgataccc ctgcaagtat cctcaaaaaa tagacggaaa 1020
gtgctgcaag gtgtgtccag gtaaaaaagc aaaagaagaa cttccaggcc aaagctttga 1080
caataaaggc tacttctgcg gggaagaac gatgcctgtg tatgagtctg tattcatgga 1140
ggatggggag acaaccagaa aaatagcact ggagactgag agaccacctc aggtagaggt 1200
ccacgttttg actattcgaa agggcattct ccagcacttc catattgaga agatctccaa 1260
gaggatgttt gaggagcttc ctcaattcaa gctggtgacc agaacaacct tgagccagt 1320
gaagatcttc accgaaggag aagctcagat cagccagatg tgttcaagtc gtgtatgcag 1380
aacagagctt gaagatttag tcaaggtttt gtacctggag agatctgaaa agggccactg 1440

```

```

ttaggcaaga cagacagtat tggatagggg aaagcaagaa aactcaagct gcagctggac 1500
tgcaggctta ttttgcttaa gtcaacagtg ccctaaaact ccaaactcaa atgcagtcaa 1560
ttattcacgc catgcacagc ataatttgct cctttgtgtg gagggtgtg tcagcccttg 1620
aacatctcct ccaaagagac tagaagagtc ttaaattata tgtgggagga ggaggatag 1680
aacatcacaa cactgctcta gtttcttgga gaatcacatt tctttacagg ttaaagacaa 1740
acaagacccc aggggttttta tctagaaagt tattcaagtg aaagaaagag aagggaattg 1800
cttagtagga gttctgcagt atagaacaat tacttgtagt aaattatacc tttgaatttt 1860
agaatgtcat gtgttctttt aaaaaaatta gctcccatc ctcctctctc actccctccc 1920
tccctccttc tctctctctc tctctctccc tctctcacag acacacacac acacacacac 1980
acacacacgc acgtccacac tcacattaaa cttaaagcttt atttgaagca aagctagcca 2040
aaattctacg ttacttttcc cttgactgga tcccaagtag cttggaagtt tttgtgccc 2100
ggagagtaaa taactgtgaa caagaggctc tgcccttagg tctttgtggc tgtttaagtc 2160
accaacaata gagtcagggt aaagaataaa aacactttca tagcctcatt cttcacttta 2220
gaagtggtaa taatttttcc ctaatgatac cacttttctt tccccctgt acctatggga 2280
cttccagaaa gaagttaa at tgagtaaaat catcagaaac tgaatccatg taagaaaaaa 2340
taattgttga agaaagaagt tgatagaatt caaaaaggcc atctttttgc tttcacatca 2400
ataaaattta ccaagtaata gatcagtact cactaatatt tttgagacca tagttgtctg 2460
gtcagaaaaa ttatatataa ttagtaaat cttagaagctc tttaaaaggg aagttttcct 2520
tcttctccaa ttataggagt tgatttttac tttgcaaagt ggctcgttc tcatgagcat 2580
ctgcatgttg actcttcagt taagaaaatt gttgttcatt tagggaggtg gatattctga 2640
tgaagatctt tatcctaaac ctctctacta tcttgtctt attcatcaag cagatatttt 2700
agtcaagaat tccagagaag gctgctccta aaatgtctac ttgcagcca ataccagagc 2760
ataaactatc cattctgggg tctggcttta gaaatcatct ttgtgggaag acctaatct 2820
tcacagcaag gatctcaggc atgccttcta gatttgttcc ctctgagggg caggaatgaa 2880
ctgtagaaat gttttaagga cccagaaacc ccatatgtct cattccatga ctataggtga 2940
gagaattctt tcctaagagg gtttgatacc aataggggaa aatgtaaaat gttcagctct 3000
tatgacaacc tggcataaag gagtcaattc ttatgaaaga gacacaaggg ctttatggcc 3060
aggggtttctt gggacaagac tctcaccagc acatcacaca cgttctctct ggaagagaga 3120
agcagtacat cccggttgag aggtcacaaa gcattagtgt gtgtgtgtgt gtgtgtgtgt 3180
gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtatgt ggtaaagggg ggaaggtgtt 3240
atgcggctgc tcctccgct ccagaggtgg cagtgtattc ataagtggga gactagtaac 3300
tagatcctaa ggcaaaaggg tgtttctcct tttggatgat tcatcccaaa gccttcccac 3360
ccaggtgttc tctgaaagct tagccttaag agaacacgca gagagtttcc ctatagatac 3420
tctgcctcc aggtgtctgg acacaccttt gcaaaatgct gtgggaagca ggagctggg 3480
agctgtgtta agtcaaagta gaaaccctcc agtgtttggt gttgtgtaga gaataggaca 3540
tagggtaaaag aggccaagct gcctgtagtt agtagagaag aatggatgtg gttcttcttg 3600
tgtatttatt tttatcataa acacttgga caacaaagac cataagcatc atttagcagt 3660
tgtagccatt ttctagttaa ctcatgtaaa caagtaagag taacataaca gtattaccct 3720
ttcactgttc tcacaggaca tgtacctaat tatggtactt atttatgtag tcaactgtatt 3780
tctggatttt taaattaata aaaaagttaa ttttgaaaa tcaaaaaaaa aaaaaaaa 3840
aaaaaaaaa aaaaaaaaaa a 3861

```

&lt;210&gt; 76

&lt;211&gt; 457

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 76

```

Met Arg Lys Lys Trp Lys Met Gly Gly Met Lys Tyr Ile Phe Ser Leu
  1             5             10             15

```

```

Leu Phe Phe Leu Leu Leu Glu Gly Gly Lys Thr Glu Gln Val Lys His
      20             25             30

```

```

Ser Glu Thr Tyr Cys Met Phe Gln Asp Lys Lys Tyr Arg Val Gly Glu
  35             40             45

```

```

Arg Trp His Pro Tyr Leu Glu Pro Tyr Gly Leu Val Tyr Cys Val Asn
  50             55             60

```

Cys Ile Cys Ser Glu Asn Gly Asn Val Leu Cys Ser Arg Val Arg Cys  
 65 70 75 80  
 Pro Asn Val His Cys Leu Ser Pro Val His Ile Pro His Leu Cys Cys  
 85 90 95  
 Pro Arg Cys Pro Asp Ser Leu Pro Pro Val Asn Asn Lys Val Thr Ser  
 100 105 110  
 Lys Ser Cys Glu Tyr Asn Gly Thr Thr Tyr Gln His Gly Glu Leu Phe  
 115 120 125  
 Val Ala Glu Gly Leu Phe Gln Asn Arg Gln Pro Asn Gln Cys Thr Gln  
 130 135 140  
 Cys Ser Cys Ser Glu Gly Asn Val Tyr Cys Gly Leu Lys Thr Cys Pro  
 145 150 155 160  
 Lys Leu Thr Cys Ala Phe Pro Val Ser Val Pro Asp Ser Cys Cys Arg  
 165 170 175  
 Val Cys Arg Gly Asp Gly Glu Leu Ser Trp Glu His Ser Asp Gly Asp  
 180 185 190  
 Ile Phe Arg Gln Pro Ala Asn Arg Glu Ala Arg His Ser Tyr His Arg  
 195 200 205  
 Ser His Tyr Asp Pro Pro Pro Ser Arg Gln Ala Gly Gly Leu Ser Arg  
 210 215 220  
 Phe Pro Gly Ala Arg Ser His Arg Gly Ala Leu Met Asp Ser Gln Gln  
 225 230 235 240  
 Ala Ser Gly Thr Ile Val Gln Ile Val Ile Asn Asn Lys His Lys His  
 245 250 255  
 Gly Gln Val Cys Val Ser Asn Gly Lys Thr Tyr Ser His Gly Glu Ser  
 260 265 270  
 Trp His Pro Asn Leu Arg Ala Phe Gly Ile Val Glu Cys Val Leu Cys  
 275 280 285  
 Thr Cys Asn Val Thr Lys Gln Glu Cys Lys Lys Ile His Cys Pro Asn  
 290 295 300  
 Arg Tyr Pro Cys Lys Tyr Pro Gln Lys Ile Asp Gly Lys Cys Cys Lys  
 305 310 315 320  
 Val Cys Pro Gly Lys Lys Ala Lys Glu Glu Leu Pro Gly Gln Ser Phe  
 325 330 335  
 Asp Asn Lys Gly Tyr Phe Cys Gly Glu Glu Thr Met Pro Val Tyr Glu  
 340 345 350  
 Ser Val Phe Met Glu Asp Gly Glu Thr Thr Arg Lys Ile Ala Leu Glu  
 355 360 365  
 Thr Glu Arg Pro Pro Gln Val Glu Val His Val Trp Thr Ile Arg Lys  
 370 375 380

Gly Ile Leu Gln His Phe His Ile Glu Lys Ile Ser Lys Arg Met Phe  
 385 390 395 400

Glu Glu Leu Pro His Phe Lys Leu Val Thr Arg Thr Thr Leu Ser Gln  
 405 410 415

Trp Lys Ile Phe Thr Glu Gly Glu Ala Gln Ile Ser Gln Met Cys Ser  
 420 425 430

Ser Arg Val Cys Arg Thr Glu Leu Glu Asp Leu Val Lys Val Leu Tyr  
 435 440 445

Leu Glu Arg Ser Glu Lys Gly His Cys  
 450 455

<210> 77

<211> 2050

<212> DNA

<213> Homo sapiens

<400> 77

```

gtgctctgag aagccggact acgcggcagc ggctcttcaa agcggagccg ggagtttttg 60
ctacagtttt cgccaccatg agtcgcagct ataatgatga gctgcagttc ttggagaaga 120
tcaataaaaa ctgctggagg atcaagaagg gcttcgtgcc caacatgcag gttgaagggtg 180
ttttctatgt gaatgatgct ctggagaaat tgatgtttga ggaattaagg aatgcctgtc 240
gaggtggtgg tgttggtggc ttcctgccag ccatgaaaca gattggcaat gtggcagccc 300
tgcttggaaat tgttcacga tctattgggc ttcctgatgt ccattcagga tatgggttg 360
ctattgggaa catggcagcc tttgatatga atgaccctga agcagtagta tccccaggtg 420
gtgtcgggtt tgacatcaac tgtggtgtcc gcttgctaag aaccaattta gatgaaagtg 480
atgtccagcc tgtgaaggag caacttgccc aagctatgtt tgaccacatt cctgttgggg 540
tggggtcaaa aggtgtcatc ccaatgaatg ccaaagactt ggaggaggcc ttggagatgg 600
gggtggactg gtccttaaga gaagggtatg cctgggctga agacaaggag cactgcgagg 660
agtacggaag gatgctgcag gctgaccca ataaagtttc tgcaaggcg aagaaaagag 720
gccttctca gttggggacc ctgggagcag gcaaccatta tgcagaaatc caggttggtg 780
atgagatttt caatgagtat gctgctaaaa aaatgggcat cgaccataag ggacaggtgt 840
gtgtgatgat ccacagtga agcagaggct tgggccacca agtagccaca gatgcgctgg 900
tagctatgga gaaggccatg aagagagaca agattatagt caatgatcgg cagttggctt 960
gtgctcgaat cgcttccca gaggggtcaag actatctgaa gggaatggca gctgctggga 1020
actatgcctg ggtcaaccgc tcttccatga ccttcttaac ccgtcaggct ttcgccaagg 1080
tcttcaacac aaccctgat gacttggacc tacatgtgat ctatgatgtt tctcacaaca 1140
ttgccaaagt ggagcagcat gtggtggacg gaaaggaacg gacactgtta gtacacagga 1200
agggatccac ccgcgctttc cctcctcacc atccctcat tgctgttgat taccaactca 1260
ctggacagcc agtgctcatt ggtggcacca tgggaacctg tagttatgtt cttactggca 1320
ctgaacaggg catgactgag acctttggaa caacctgtca tggagcgggc cgtgcattgt 1380
cccagcaaaa atctcgacgt aatttagatt tccaggatgt cttagacaaa ttggcagata 1440
tgggaattgc gatccgtgtt gcctcaccca aactgggtat ggaagaggct cctgagtcct 1500
ataagaatgt gacagatgtg gtaaatacct gccatgatgc tggacagcag ggctgcctga 1560
ttaaactgag accaattgct gtgatcaaag gatagaacct tggacagcag ggctgcctga 1620
caccaccaac cctctctgaa gtggaagtgg actgacatgc tcttctgaca tcagactcaa 1680
ggcgggacaa gttgcaaagt gtgcagctgt aactgctcac gccaaaatgg ctgatgggga 1740
ggctgctgct ttcaggggccc cgtgcttgta aaataacctt ccaggaagag gcacattgcc 1800
cacctttgga aagggaggaa tatgccttct ccttggttgt tccacagagt ttaggaaaaa 1860
tctgttaggg atgggtagat gtcaaaactgc cttacgcagt catactgata ttagccatc 1920
agattgatct tcttcacacc aagctctggt tacattccga gaggtgtcat gaagaaagtt 1980
ctgttcaata agggaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040
aaaaaaaaaa

```

<210> 78

<211> 505

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 78

```

Met Ser Arg Ser Tyr Asn Asp Glu Leu Gln Phe Leu Glu Lys Ile Asn
 1              5              10              15

Lys Asn Cys Trp Arg Ile Lys Lys Gly Phe Val Pro Asn Met Gln Val
      20              25              30

Glu Gly Val Phe Tyr Val Asn Asp Ala Leu Glu Lys Leu Met Phe Glu
      35              40              45

Glu Leu Arg Asn Ala Cys Arg Gly Gly Gly Val Gly Gly Phe Leu Pro
 50              55              60

Ala Met Lys Gln Ile Gly Asn Val Ala Ala Leu Pro Gly Ile Val His
 65              70              75              80

Arg Ser Ile Gly Leu Pro Asp Val His Ser Gly Tyr Gly Phe Ala Ile
      85              90              95

Gly Asn Met Ala Ala Phe Asp Met Asn Asp Pro Glu Ala Val Val Ser
      100              105              110

Pro Gly Gly Val Gly Phe Asp Ile Asn Cys Gly Val Arg Leu Leu Arg
      115              120              125

Thr Asn Leu Asp Glu Ser Asp Val Gln Pro Val Lys Glu Gln Leu Ala
      130              135              140

Gln Ala Met Phe Asp His Ile Pro Val Gly Val Gly Ser Lys Gly Val
      145              150              155              160

Ile Pro Met Asn Ala Lys Asp Leu Glu Glu Ala Leu Glu Met Gly Val
      165              170              175

Asp Trp Ser Leu Arg Glu Gly Tyr Ala Trp Ala Glu Asp Lys Glu His
      180              185              190

Cys Glu Glu Tyr Gly Arg Met Leu Gln Ala Asp Pro Asn Lys Val Ser
      195              200              205

Ala Arg Ala Lys Lys Arg Gly Leu Pro Gln Leu Gly Thr Leu Gly Ala
      210              215              220

Gly Asn His Tyr Ala Glu Ile Gln Val Val Asp Glu Ile Phe Asn Glu
      225              230              235              240

Tyr Ala Ala Lys Lys Met Gly Ile Asp His Lys Gly Gln Val Cys Val
      245              250              255

Met Ile His Ser Gly Ser Arg Gly Leu Gly His Gln Val Ala Thr Asp
      260              265              270

Ala Leu Val Ala Met Glu Lys Ala Met Lys Arg Asp Lys Ile Ile Val
      275              280              285

Asn Asp Arg Gln Leu Ala Cys Ala Arg Ile Ala Ser Pro Glu Gly Gln
      290              295              300

```

Asp Tyr Leu Lys Gly Met Ala Ala Ala Gly Asn Tyr Ala Trp Val Asn  
 305 310 315 320  
 Arg Ser Ser Met Thr Phe Leu Thr Arg Gln Ala Phe Ala Lys Val Phe  
 325 330 335  
 Asn Thr Thr Pro Asp Asp Leu Asp Leu His Val Ile Tyr Asp Val Ser  
 340 345 350  
 His Asn Ile Ala Lys Val Glu Gln His Val Val Asp Gly Lys Glu Arg  
 355 360 365  
 Thr Leu Leu Val His Arg Lys Gly Ser Thr Arg Ala Phe Pro Pro His  
 370 375 380  
 His Pro Leu Ile Ala Val Asp Tyr Gln Leu Thr Gly Gln Pro Val Leu  
 385 390 395 400  
 Ile Gly Gly Thr Met Gly Thr Cys Ser Tyr Val Leu Thr Gly Thr Glu  
 405 410 415  
 Gln Gly Met Thr Glu Thr Phe Gly Thr Thr Cys His Gly Ala Gly Arg  
 420 425 430  
 Ala Leu Ser Arg Ala Lys Ser Arg Arg Asn Leu Asp Phe Gln Asp Val  
 435 440 445  
 Leu Asp Lys Leu Ala Asp Met Gly Ile Ala Ile Arg Val Ala Ser Pro  
 450 455 460  
 Lys Leu Val Met Glu Glu Ala Pro Glu Ser Tyr Lys Asn Val Thr Asp  
 465 470 475 480  
 Val Val Asn Thr Cys His Asp Ala Gly Ile Ser Lys Lys Ala Ile Lys  
 485 490 495  
 Leu Arg Pro Ile Ala Val Ile Lys Gly  
 500 505

&lt;210&gt; 79

&lt;211&gt; 1178

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 79

gccaaatgtc cgggtcaagat gtcacacagc tccagtgggt cagccagtct gagtcaggtt 60  
 tctccagggg aagaaacaga tcaaactgaa accgtgtcag ttcagtcttc ggtattgggg 120  
 aagggtgtaa aacatcgacc cccaccaatc aaacttccct caagctcagg aaatagttcc 180  
 tcaggtaact attttacacc acaacagaca agcagctttc tcaaattctc aactcctcct 240  
 ccttcttcta agccatcaag tattctctcg aaatcatctg tggatctcaa tcaagtttagc 300  
 atgcttttctc cagctgcctc atcacctgcc agtcatcac aaagaaccac ggccaccagc 360  
 gtcattggcaa actctgcttg acttaacttc atcaatgtag tgggctctgt ttgtggggcc 420  
 caggctttga tgagtgttc aaaccccatg ctgggctgta aactggtgc cataactcct 480  
 gcaggaataa acctgagcgg ccttctaccc tcaggagggt tgctaccaa tgactgccc 540  
 agtgcaatgc aggcagcttc tcaagcaggt gtccatttg gtttaaaaaa tacttcaagt 600  
 ctcaggccct taaatctact ccagcttcca ggtggttcac ttatttttaa cactctgcag 660  
 cagcagcaac agcagctctc ccagtttaca ccacaacaac ctcagcagcc cacaacttgt 720  
 agtcctcaac agccagggga gcagggttct gagcaagggt caaccagtca agaacaggcc 780

```

ttatctgctc agcaagctgc tgttattaac cttactggag taggaagttt tatgcagtca 840
caggcagctg cagttgcat tcttgcatc tcaaattggct atggcagcag cagcagcaca 900
aacagctcag ctacatcatc atcggcatc aggcatccag tcaaaaagta aaatgaagag 960
aggcatgcca accactccaa aattttgagt cttgcattac tttttgttcc ttttttaaaa 1020
acacaagagc actgaatcaa aagaattgag tttctacttt ttgttttttt taatgtgtca 1080
gtatttttaca ttgctagatg tacaaacttt atacagaagc acaaccttat cattttttaa 1140
taaaaacagg gaaatggttt aacaaaaaaa aaaaaaaa 1178

```

&lt;210&gt; 80

&lt;211&gt; 310

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

```

Met Ser His Ser Ser Gly Ser Ala Ser Leu Ser Gln Val Ser Pro
  1              5              10              15

Gly Lys Glu Thr Asp Gln Thr Glu Thr Val Ser Val Gln Ser Ser Val
      20              25              30

Leu Gly Lys Gly Val Lys His Arg Pro Pro Pro Ile Lys Leu Pro Ser
      35              40              45

Ser Ser Gly Asn Ser Ser Ser Gly Asn Tyr Phe Thr Pro Gln Gln Thr
      50              55              60

Ser Ser Phe Leu Lys Ser Pro Thr Pro Pro Pro Ser Ser Lys Pro Ser
      65              70              75              80

Ser Ile Pro Arg Lys Ser Ser Val Asp Leu Asn Gln Val Ser Met Leu
      85              90              95

Ser Pro Ala Ala Leu Ser Pro Ala Ser Ser Ser Gln Arg Thr Thr Ala
      100             105             110

Thr Gln Val Met Ala Asn Ser Ala Gly Leu Asn Phe Ile Asn Val Val
      115             120             125

Gly Ser Val Cys Gly Ala Gln Ala Leu Met Ser Gly Ser Asn Pro Met
      130             135             140

Leu Gly Cys Asn Thr Gly Ala Ile Thr Pro Ala Gly Ile Asn Leu Ser
      145             150             155             160

Gly Leu Leu Pro Ser Gly Gly Leu Leu Pro Asn Ala Leu Pro Ser Ala
      165             170             175

Met Gln Ala Ala Ser Gln Ala Gly Val Pro Phe Gly Leu Lys Asn Thr
      180             185             190

Ser Ser Leu Arg Pro Leu Asn Leu Leu Gln Leu Pro Gly Gly Ser Leu
      195             200             205

Ile Phe Asn Thr Leu Gln Gln Gln Gln Gln Leu Ser Gln Phe Thr
      210             215             220

Pro Gln Gln Pro Gln Gln Pro Thr Thr Cys Ser Pro Gln Gln Pro Gly
      225             230             235             240

Glu Gln Gly Ser Glu Gln Gly Ser Thr Ser Gln Glu Gln Ala Leu Ser

```

<210> 83  
<211> 832



&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 83

```

ccttgcatta ccttcttctg ccctatctgc tgctaggtgt aaacctgttt tttttcaccc 60
tgacttggtg aaccaatcct ggcattataa caaaagcaaa tgaattatta tttcttcatg 120
tttatgaatt tgaatgaagt atgtttccaa agaacgtgag gtgctctact tgtgatttaa 180
ggaaaccagc tcgatccaag cactgcagtg tgtgtaactg gtgtgtgcac cgtttcgacc 240
atcactgtgt ttgggtgaac aactgcatcg gggcctggaa catcaggtag ttcctcatct 300
acgtcttgac cttgacggcc tcggctgcca ccgtcgccat tgtgagcacc acttttctgg 360
tccacttggt ggtgatgtca gatttatacc aggagactta catcgatgac cttggacacc 420
ttccatgtta tggacacggc ctttcttatt cagtacctgt tcctgacttt tccacggatt 480
gtcttcatgc tgggctttgt cgtggttctg agcttctctc tgggtggcta cctgttgttt 540
gtcctgtatc tggcggccac caaccagact actaacgagt ggtacagagg tgactgggcc 600
tggtgccagc gttgtccctt tgtggcctgg cctccgtcag cagagcccca agtccaccgg 660
aacattcact cccatgggct tcggagcaac cttcaagaga tctttctacc tgcctttcca 720
tgtcatgaga ggaagaaaca agaatgacaa gtgtatgact gcctttgagc tgtagtcccc 780
gtttatttac acatgtggat cctcgtttct cctccgattg aattctagac ct 832

```

&lt;210&gt; 84

&lt;211&gt; 144

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 84

```

Met Phe Pro Lys Asn Val Arg Cys Ser Thr Cys Asp Leu Arg Lys Pro
  1              5              10              15

Ala Arg Ser Lys His Cys Ser Val Cys Asn Trp Cys Val His Arg Phe
      20              25              30

Asp His His Cys Val Trp Val Asn Asn Cys Ile Gly Ala Trp Asn Ile
      35              40              45

Arg Tyr Phe Leu Ile Tyr Val Leu Thr Leu Thr Ala Ser Ala Ala Thr
      50              55              60

Val Ala Ile Val Ser Thr Thr Phe Leu Val His Leu Val Val Met Ser
      65              70              75              80

Asp Leu Tyr Gln Glu Thr Tyr Ile Asp Asp Leu Gly His Leu Pro Cys
      85              90              95

Tyr Gly His Gly Leu Ser Tyr Ser Val Pro Val Pro Asp Phe Ser Thr
      100              105              110

Asp Cys Leu His Ala Gly Leu Cys Arg Gly Ser Glu Leu Pro Pro Gly
      115              120              125

Trp Leu Pro Val Val Cys Pro Val Ser Gly Gly His Gln Pro Asp Tyr
      130              135              140

```

&lt;210&gt; 85

&lt;211&gt; 3790

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 85

```

aacggcagtc tcaatctggc cccacacctt tcttgggctt gtaggaaggt ggacatgggc 60

```

tccccggagac	aagacaagtg	atatgttgaa	ctgttcgggtg	gctggaatca	actgctcctg	120
gagtgcaccta	aggccagtgt	ttatcagaac	ttagccagggt	ccagccaagc	aggcacagat	180
gctctgcttat	gaaatgccac	gcaggcagag	actgacaagc	ggtaggaact	gagctttccc	240
cttggaactgc	tgcttcctgc	tgtgttcagg	ggaggggggtc	actttctggtc	aactctgctg	300
ctgctgctgct	tgctgctgct	acttcagctt	cctctccact	caaggtaagc	aggctaaggg	360
agggcagggt	gctagggaaa	gctttgtacc	atgaacagga	tccgaaagtt	tttccgagga	420
agtgggctgag	tcttggcatt	tatctttgtg	gcttctgtca	tctggctcct	ctttgacatg	480
gcagctctcc	gcctctcatt	cagtgcagatc	aacactcggg	tcatcaagga	agacattgtg	540
aggaggggagc	ggataggatt	cagagttcag	ccagaccaag	gaaaaatttt	ttacagcagc	600
ataaaagaga	tgaaacctcc	cctaagggga	catgggaaag	gggcatgggg	caaagagaat	660
gttagaaaaa	ctgaggagag	tgtgctcaag	gttgagggtg	acttggaacca	aacctgaggg	720
gaaagaaaaa	tgagaatgc	cctgggaagg	ggcaagggtg	tgccgttgtg	gcctcctgca	780
catctgcaga	ccctccctgt	gactcctaac	aagcagaaga	cagacggggag	aggcaccaaa	840
cctgaagcct	cctctcacca	ggggacacca	aagcaaacga	cagctcagggt	ggctccaaag	900
acctcattca	tagcagcaaa	aggaactcag	gtagtcaaaa	tatcagtaca	catgggacgt	960
gtcagtttaa	aacaggagcc	ccggaagagt	catagtccca	gcagtgcacac	atcaaaacta	1020
gcagctgaaa	gggacttgaa	tgtgaccatc	agtcttagta	ctgatagacc	aaagcagcga	1080
tcacaggcag	tagcaaacga	gagggcacac	cctgccagca	cagcagtgcc	gaagtctggg	1140
gaagccatgg	ccttaaacaa	aactaagact	cagagcaaaag	aagtcaatgc	aaataaacac	1200
aaagccaata	cgagtcttcc	ttttcctaag	ttcactgtca	attcaaatcg	cttaagggaag	1260
caatctatta	atgagacacc	tttggaagt	ttgtcaaagg	atgatggagc	tagaggggct	1320
catgggaaga	aactcaattt	ctctgaaagc	catcttgtga	ttataaccaa	agaggaagag	1380
caaaaggcag	accccaaaga	ggctctaat	tctaaaacca	aaacaatatt	tcctaaagta	1440
ttgggtaaaa	gccaaagtaa	acacatttcc	aggaatagaa	gtgagatgtc	ttcctcttca	1500
cttgctccac	atagagtgcc	actgtcccaa	actaaccatg	ctttaactgg	agggctagag	1560
ccagcaaaaa	tcaacataac	tgccaaagcc	ccctctacag	aatacaacca	gagtcataata	1620
aaagcccttt	tacctgaaga	cagtggaaag	caccaggtgt	taagaattga	tgtgacactt	1680
tctccaagggt	accccaaagc	tccagggcag	tttgggcgctc	ctgtagttgt	cccccatgga	1740
aaggagaagg	aggcagaaag	aagatggaaa	gaaggaaact	tcaatgtcta	ccttagcgat	1800
ttgatcccgag	tggatagagc	cattgaagac	accagacctg	ctggatgtgc	agagcagcta	1860
gttcacaata	acctcccaac	caccagtgct	atcatgtgct	ttgtggatga	agtgtggtcc	1920
actctctga	gatctgttca	cagtgtcatc	aatcgctctc	ctccacacct	catcaaggag	1980
attctgctgg	tagatgactt	cagcaccaaa	cactatctaa	aagataattt	ggataaatac	2040
atgtcccagt	ttccaaaagt	tgggattctt	gcctcaaaag	agagacatgg	cttaataagg	2100
gccaggctgg	caggagcaca	gaatgcaaca	ggatgtgtgt	tgacattttt	agattctcat	2160
gtggaatgta	acgttggttg	gttggaacct	cttctggaaa	gagttttatt	aagtagaaag	2220
aaagtggcct	gtccagtaat	cgaagtcatc	aatgataagg	atatgagtta	catgacagtg	2280
gataactttc	aaagaggcat	ctttgtgtgg	cccatgaact	ttgggtggag	aacaattcct	2340
ccagatgtca	ttgcaaaaaa	cagaattaaa	gaaactgata	caataagggtg	ccctgtcatg	2400
gctgggtgac	tgttttctat	tgacaaaagt	tacttttttg	aacttgggaa	atacgaccct	2460
ggccttgatg	tttggggtgg	ggaaaatag	gactctcat	tcaagggtgtg	gatgtgtggt	2520
gggtgaaattg	agatcattcc	ctgctcccga	gtgggccata	tattcagaaa	tgacaattcca	2580
tattccttcc	ccaaagaccg	gatgaagaca	gtggagcgga	acttggtgcg	ggttgccgag	2640
gtctggctgg	atgagtataa	ggagctgttc	tatggccatg	gagaccacct	catcgaccaa	2700
gggctagatg	ttggcaacct	caccagcaa	agggagctgc	gaaagaaact	gaagtgcaaa	2760
agtttcaaat	ggtacttgga	gaatgtcttt	cctgacttaa	gggctcccat	tgtgagagct	2820
agtggatgct	ttattaatgt	ggctttgggt	aaatgcattt	ccattgaaaa	cactacagtc	2880
attctggaag	actgcgatgg	gagcaaaag	cttcaacaat	ttaattacac	ctgggttaaga	2940
cttattaat	gtggagaatg	gtgtatagcc	cccatccctg	ataaagggagc	cgtaaggctg	3000
cacccttgtg	ataacagaaa	caaagggtca	aaatggctgc	ataaatcaac	atcagctctt	3060
catccagaac	tggatgaatca	cattgttttt	gaaaacaatc	agcaattatt	atgcttgga	3120
ggaaattttt	ctcaaaaagat	cctgaaaagta	gctgcctgtg	acccagtgaa	gccatatcaa	3180
aagtggaaat	ttgaaaaata	ttatgaagcc	tgaagtgtaa	ctgatgtttt	tatatagtaa	3240
accattaaa	tactgtgaaa	ataacactga	acttggaaac	tatatctctc	agcggtagtt	3300
taaattttca	attttaataa	catttgaatg	gaagattttt	tataaatcac	aatatttggg	3360
atacaaaaag	atgactcagg	aaaacagtcc	aacattggac	tgaagtcctt	cttcggaact	3420
gggtggcctt	tgaattgcct	gctttccacc	ctatgctaga	cctcatcatg	caaatttccc	3480
tgtgaaagct	aacaggtaac	tggaaatgaa	gacagaagga	cttgagaaag	catgaggata	3540
ttcccaatga	ctatgtttgg	taataatcag	ctcttctggc	ccacaagtag	gaatgatcaa	3600
tgagaactta	acttagtctc	ttatttgggg	attttttcat	caaacaaaaa	tttcttgagt	3660

tcttatggct agaagacctc agatgcccac agctgtcacg tttgtgaaat ccctccagac 3720  
 tacatgcatg cttacctaac agtttgaaat agtattgatc tactgctggt aaaaaaaaaa 3780  
 aaaaaaaaaa 3790

<210> 86

<211> 940

<212> PRT

<213> Homo sapiens

<400> 86

Met Asn Arg Ile Arg Lys Phe Phe Arg Gly Ser Gly Arg Val Leu Ala  
 1 5 10 15

Phe Ile Phe Val Ala Ser Val Ile Trp Leu Leu Phe Asp Met Ala Ala  
 20 25 30

Leu Arg Leu Ser Phe Ser Glu Ile Asn Thr Arg Val Ile Lys Glu Asp  
 35 40 45

Ile Val Arg Arg Glu Arg Ile Gly Phe Arg Val Gln Pro Asp Gln Gly  
 50 55 60

Lys Ile Phe Tyr Ser Ser Ile Lys Glu Met Lys Pro Pro Leu Arg Gly  
 65 70 75 80

His Gly Lys Gly Ala Trp Gly Lys Glu Asn Val Arg Lys Thr Glu Glu  
 85 90 95

Ser Val Leu Lys Val Glu Val Asp Leu Asp Gln Thr Gln Arg Glu Arg  
 100 105 110

Lys Met Gln Asn Ala Leu Gly Arg Gly Lys Val Val Pro Leu Trp His  
 115 120 125

Pro Ala His Leu Gln Thr Leu Pro Val Thr Pro Asn Lys Gln Lys Thr  
 130 135 140

Asp Gly Arg Gly Thr Lys Pro Glu Ala Ser Ser His Gln Gly Thr Pro  
 145 150 155 160

Lys Gln Thr Thr Ala Gln Gly Ala Pro Lys Thr Ser Phe Ile Ala Ala  
 165 170 175

Lys Gly Thr Gln Val Val Lys Ile Ser Val His Met Gly Arg Val Ser  
 180 185 190

Leu Lys Gln Glu Pro Arg Lys Ser His Ser Pro Ser Ser Asp Thr Ser  
 195 200 205

Lys Leu Ala Ala Glu Arg Asp Leu Asn Val Thr Ile Ser Leu Ser Thr  
 210 215 220

Asp Arg Pro Lys Gln Arg Ser Gln Ala Val Ala Asn Glu Arg Ala His  
 225 230 235 240

Pro Ala Ser Thr Ala Val Pro Lys Ser Gly Glu Ala Met Ala Leu Asn  
 245 250 255

Lys Thr Lys Thr Gln Ser Lys Glu Val Asn Ala Asn Lys His Lys Ala  
 260 265 270

Asn Thr Ser Leu Pro Phe Pro Lys Phe Thr Val Asn Ser Asn Arg Leu  
 275 280 285  
 Arg Lys Gln Ser Ile Asn Glu Thr Pro Leu Gly Ser Leu Ser Lys Asp  
 290 295 300  
 Asp Gly Ala Arg Gly Ala His Gly Lys Lys Leu Asn Phe Ser Glu Ser  
 305 310 315 320  
 His Leu Val Ile Ile Thr Lys Glu Glu Glu Gln Lys Ala Asp Pro Lys  
 325 330 335  
 Glu Val Ser Asn Ser Lys Thr Lys Thr Ile Phe Pro Lys Val Leu Gly  
 340 345 350  
 Lys Ser Gln Ser Lys His Ile Ser Arg Asn Arg Ser Glu Met Ser Ser  
 355 360 365  
 Ser Ser Leu Ala Pro His Arg Val Pro Leu Ser Gln Thr Asn His Ala  
 370 375 380  
 Leu Thr Gly Gly Leu Glu Pro Ala Lys Ile Asn Ile Thr Ala Lys Ala  
 385 390 395 400  
 Pro Ser Thr Glu Tyr Asn Gln Ser His Ile Lys Ala Leu Leu Pro Glu  
 405 410 415  
 Asp Ser Gly Thr His Gln Val Leu Arg Ile Asp Val Thr Leu Ser Pro  
 420 425 430  
 Arg Asp Pro Lys Ala Pro Gly Gln Phe Gly Arg Pro Val Val Val Pro  
 435 440 445  
 His Gly Lys Glu Lys Glu Ala Glu Arg Arg Trp Lys Glu Gly Asn Phe  
 450 455 460  
 Asn Val Tyr Leu Ser Asp Leu Ile Pro Val Asp Arg Ala Ile Glu Asp  
 465 470 475 480  
 Thr Arg Pro Ala Gly Cys Ala Glu Gln Leu Val His Asn Asn Leu Pro  
 485 490 495  
 Thr Thr Ser Val Ile Met Cys Phe Val Asp Glu Val Trp Ser Thr Leu  
 500 505 510  
 Leu Arg Ser Val His Ser Val Ile Asn Arg Ser Pro Pro His Leu Ile  
 515 520 525  
 Lys Glu Ile Leu Leu Val Asp Asp Phe Ser Thr Lys Asp Tyr Leu Lys  
 530 535 540  
 Asp Asn Leu Asp Lys Tyr Met Ser Gln Phe Pro Lys Val Arg Ile Leu  
 545 550 555 560  
 Arg Leu Lys Glu Arg His Gly Leu Ile Arg Ala Arg Leu Ala Gly Ala  
 565 570 575  
 Gln Asn Ala Thr Gly Asp Val Leu Thr Phe Leu Asp Ser His Val Glu  
 580 585 590

Cys Asn Val Gly Trp Leu Glu Pro Leu Leu Glu Arg Val Tyr Leu Ser  
 595 600 605  
 Arg Lys Lys Val Ala Cys Pro Val Ile Glu Val Ile Asn Asp Lys Asp  
 610 615 620  
 Met Ser Tyr Met Thr Val Asp Asn Phe Gln Arg Gly Ile Phe Val Trp  
 625 630 635 640  
 Pro Met Asn Phe Gly Trp Arg Thr Ile Pro Pro Asp Val Ile Ala Lys  
 645 650 655  
 Asn Arg Ile Lys Glu Thr Asp Thr Ile Arg Cys Pro Val Met Ala Gly  
 660 665 670  
 Gly Leu Phe Ser Ile Asp Lys Ser Tyr Phe Phe Glu Leu Gly Thr Tyr  
 675 680 685  
 Asp Pro Gly Leu Asp Val Trp Gly Gly Glu Asn Met Glu Leu Ser Phe  
 690 695 700  
 Lys Val Trp Met Cys Gly Gly Glu Ile Glu Ile Ile Pro Cys Ser Arg  
 705 710 715 720  
 Val Gly His Ile Phe Arg Asn Asp Asn Pro Tyr Ser Phe Pro Lys Asp  
 725 730 735  
 Arg Met Lys Thr Val Glu Arg Asn Leu Val Arg Val Ala Glu Val Trp  
 740 745 750  
 Leu Asp Glu Tyr Lys Glu Leu Phe Tyr Gly His Gly Asp His Leu Ile  
 755 760 765  
 Asp Gln Gly Leu Asp Val Gly Asn Leu Thr Gln Gln Arg Glu Leu Arg  
 770 775 780  
 Lys Lys Leu Lys Cys Lys Ser Phe Lys Trp Tyr Leu Glu Asn Val Phe  
 785 790 795 800  
 Pro Asp Leu Arg Ala Pro Ile Val Arg Ala Ser Gly Val Leu Ile Asn  
 805 810 815  
 Val Ala Leu Gly Lys Cys Ile Ser Ile Glu Asn Thr Thr Val Ile Leu  
 820 825 830  
 Glu Asp Cys Asp Gly Ser Lys Glu Leu Gln Gln Phe Asn Tyr Thr Trp  
 835 840 845  
 Leu Arg Leu Ile Lys Cys Gly Glu Trp Cys Ile Ala Pro Ile Pro Asp  
 850 855 860  
 Lys Gly Ala Val Arg Leu His Pro Cys Asp Asn Arg Asn Lys Gly Leu  
 865 870 875 880  
 Lys Trp Leu His Lys Ser Thr Ser Val Phe His Pro Glu Leu Val Asn  
 885 890 895  
 His Ile Val Phe Glu Asn Asn Gln Gln Leu Leu Cys Leu Glu Gly Asn  
 900 905 910

Phe Ser Gln Lys Ile Leu Lys Val Ala Ala Cys Asp Pro Val Lys Pro  
 915 920 925

Tyr Gln Lys Trp Lys Phe Glu Lys Tyr Tyr Glu Ala  
 930 935 940

<210> 87  
 <211> 1200  
 <212> DNA  
 <213> Homo sapiens

<400> 87  
 ggcttctcgg agcggcgctg ggcgaccgga gcagggtcga gatgtcctac atccccggcc 60  
 agccggtcac cgccgtggtg caaagagttg aaattcacaa gctgcgtcaa ggtgagaact 120  
 taatccctggg ttccagcatt ggaggtggaa tcgaccagga tcttcccag aatcccttct 180  
 atgaagacaa gacggacaag ggtatttatg tcacacgggt gtctgaagga ggccctgctg 240  
 aaatcgctgg gctgcagatt ggagacaaga tcatgcaggt gaacggctgg gacatgacca 300  
 tggtcacaca cgaccaggcc cgcaagcggc tcaccaagcg ctggaggag gtggtgcgtc 360  
 tgctgggtgac gcggcagtcg ctgcagaagg cgtgcagcag tcatgctgtc ctagcagcca 420  
 ccaccatctg cgactcctgc ctgccgcctc tctgtacagt aacgccactt ccacactctg 480  
 tccccatctg gcttctgctg accagctttc tctcctggac accgaggatt ggaaataagg 540  
 gcctggagct gagtagtagc cagtctgctg tgaccacagg ctccaggtccg accctgctgc 600  
 ttggccacag cagtggctgg gcaagtggga accactatct cttgggagcc cccaaaagct 660  
 gggaaatgct ggaggaacca ggccctttccc gcttttgctt ggctgcaggg ttgggctccg 720  
 cccctgcccc ccagccttgg tgtgtccaca ccgcagtgtc tctgcccctc gggggactgg 780  
 acacacatcc tgccagaggc gctacgaagc tttgcccaga tgaagccagg tgggctccgc 840  
 gttcactccc actctcccga ggggtgctgg cctccccagg gtttgccttc ttacggattt 900  
 agacgaggtt cgaggctcac ctatcagggc agctctcagg attgtcattt tcctctttgc 960  
 ctgtgggttt aacttttgta tttttttaat cacaagtttg atacaaaatg tttttatcgt 1020  
 actctttgga gatgccatt ctacttttga atttagcttt tactaattcg catctggaag 1080  
 ctcagcaagt gcacaagcct tactttggtt accgtggaaa ccactgccgc ccctcccga 1140  
 tgtggtgtgc tgaataaaaa tgctggaatt caaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200

<210> 88  
 <211> 286  
 <212> PRT  
 <213> Homo sapiens

<400> 88  
 Met Ser Tyr Ile Pro Gly Gln Pro Val Thr Ala Val Val Gln Arg Val  
 1 5 10 15  
 Glu Ile His Lys Leu Arg Gln Gly Glu Asn Leu Ile Leu Gly Phe Ser  
 20 25 30  
 Ile Gly Gly Gly Ile Asp Gln Asp Pro Ser Gln Asn Pro Phe Tyr Glu  
 35 40 45  
 Asp Lys Thr Asp Lys Gly Ile Tyr Val Thr Arg Val Ser Glu Gly Gly  
 50 55 60  
 Pro Ala Glu Ile Ala Gly Leu Gln Ile Gly Asp Lys Ile Met Gln Val  
 65 70 75 80  
 Asn Gly Trp Asp Met Thr Met Val Thr His Asp Gln Ala Arg Lys Arg  
 85 90 95  
 Leu Thr Lys Arg Ser Glu Glu Val Val Arg Leu Leu Val Thr Arg Gln

100	105	110
Ser Leu Gln Lys Ala Cys Ser	Ser His Ala Val Leu	Ala Ala Thr Thr
115	120	125
Ile Cys Asp Ser Cys Leu Pro	Leu Cys Thr Val Thr	Pro Leu Pro
130	135	140
His Ser Val Pro Ile Trp Leu Leu	Leu Thr Ser Phe Leu Ser	Trp Thr
145	150	155
Pro Arg Ile Gly Asn Lys Gly Leu	Glu Leu Ser Ser Ser	Gln Ser Ala
165	170	175
Val Thr Thr Gly Ser Gly Pro Thr	Leu Leu Leu Gly His	Ser Ser Gly
180	185	190
Trp Ala Ser Gly Asn His Tyr Leu	Leu Gly Ala Pro Lys	Ser Trp Glu
195	200	205
Met Leu Glu Glu Pro Gly Leu Ser	Arg Phe Cys Leu Ala	Ala Gly Leu
210	215	220
Gly Ser Ala Pro Ala Pro Gln Pro	Trp Cys Val His Thr	Ala Val Leu
225	230	235
Leu Pro Leu Gly Gly Leu Asp Thr	His Pro Ala Arg Gly	Ala Thr Lys
245	250	255
Leu Cys Pro Asp Glu Ala Arg Trp	Ala Pro Arg Ser Leu	Pro Leu Ser
260	265	270
Arg Gly Val Leu Ala Ser Pro Gly	Phe Ala Phe Leu Arg	Ile
275	280	285

&lt;210&gt; 89

&lt;211&gt; 1023

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 89

```

ccaacatgga gactttgtac cgtgtcccgt tcttagtgct cgaatgtccc aacctgaagc 60
tgaagaagcc gccctgggtg cacatgccgt cggccatgac tgtgtatgct ctggtggtgg 120
tgtcttactt cctcatcacc ggaggaataa tttatgatgt tattgttgaa cctccaagtg 180
tcggttctat gactgatgaa catgggcatac agaggccagt agctttcttg gcctacagag 240
taaattggaca atatattatg gaaggacttg catccagctt cctattttaca atgggaggtt 300
taggtttcat aatcctggac cgatcgaatg caccaaatac cccaaaactc aatagattcc 360
ttcttctggt cattggattc gtctgtgtcc tattgagttt tttcatggct agagtattca 420
tgagaatgaa actgccgggc tatctgatgg gttagagtgc ctttgagaag aaatcagtgg 480
atactggatt tgctcctgtc aatgaagttt taaaggctgt accaatcctc taatatgaaa 540
tgtggaaaag aatgaagagc agcagtaaaa gaaatatcta gtgaaaaaac aggaagcgta 600
ttgaagcttg gactagaatt tcttcttggt attaaagaga caagtttatc acagaatttt 660
ttttcctgct ggcctattgc tataccaatg atggtgagtg gcattttctt tttagttttt 720
cattaaaata tattccatat ctacaactat aatatcaaat aaagtgatta ttttttacia 780
ccctcttaac attttttggg gatgacattt ctgattttca gaaattaaca taaaatccag 840
aagcaagatt ccgtaagctg agaactctgg acagttgatc agctttacct atggtgcttt 900
gcctttaact agagtgtgtg atggtagatt atttcagata tgtatgtaaa actgtttcct 960
gaacaataag atgtatgaac ggagcagaaa taaatacttt ttctaattaa aaaaaaaaaa 1020
aaa
1023

```

<210> 90  
 <211> 149  
 <212> PRT  
 <213> Homo sapiens

<400> 90  
 Met Glu Thr Leu Tyr Arg Val Pro Phe Leu Val Leu Glu Cys Pro Asn  
     1                    5                    10                    15  
 Leu Lys Leu Lys Lys Pro Pro Trp Leu His Met Pro Ser Ala Met Thr  
                     20                    25                    30  
 Val Tyr Ala Leu Val Val Val Ser Tyr Phe Leu Ile Thr Gly Gly Ile  
           35                    40                    45  
 Ile Tyr Asp Val Ile Val Glu Pro Pro Ser Val Gly Ser Met Thr Asp  
       50                    55                    60  
 Glu His Gly His Gln Arg Pro Val Ala Phe Leu Ala Tyr Arg Val Asn  
       65                    70                    75                    80  
 Gly Gln Tyr Ile Met Glu Gly Leu Ala Ser Ser Phe Leu Phe Thr Met  
                     85                    90                    95  
 Gly Gly Leu Gly Phe Ile Ile Leu Asp Arg Ser Asn Ala Pro Asn Ile  
           100                    105                    110  
 Pro Lys Leu Asn Arg Phe Leu Leu Leu Phe Ile Gly Phe Val Cys Val  
       115                    120                    125  
 Leu Leu Ser Phe Phe Met Ala Arg Val Phe Met Arg Met Lys Leu Pro  
       130                    135                    140  
 Gly Tyr Leu Met Gly  
 145

<210> 91  
 <211> 3901  
 <212> DNA  
 <213> Homo sapiens

<400> 91  
 gccatggagg gagtgagcgc gctgctggcc cgctgcccga cggccggcct ggccggcggc 60  
 ctgggggtca cggcgtgcgc cgcggccggc gtgttgctct accggatcgc gcggaggatg 120  
 aagccaacgc acacgatggt caactgctgg ttctgcaacc aggatacgt ggtgccctat 180  
 gggaaccgca actgctggga ctgtccccac tgcgagcagt acaacggctt ccaggagaac 240  
 ggcgactaca acaagccgat ccccgcccag tacttgagac acctgaacca cgtggtgagc 300  
 agcgcgcccga gcctgcgcga cccttcgcag ccgcagcagt gggtgagcag ccaagtctcg 360  
 ctgtgcaaga ggtgcaacca ccaccagacc accaagatca agcagctggc cgccttcgct 420  
 ccccgcgagg agggcaggta tgacgaggag gtcgaggtgt accggcatca cctggagcag 480  
 atgtacaagc tgtgcccggc gtgccaagcg gctgtggagt actacatcaa gcaccagaac 540  
 cgccagctgc gcgccctggt gctcagccac cagttcaagc gccgggaggc cgaccagacc 600  
 cacgcacaga acttctcctc cgccgtgaag tccccggtcc aggtcatcct gctccgtgac 660  
 ctgccttcc tggcctgcgc ctctctactg accaccgcgc tgtatggggc cagcggagac 720  
 ttcgccccag gcaccactgt gccctggcc ctgccacctg gtggcaatgg ctgagccaca 780  
 cctgacaatg gcaccacccc tggggccgag ggctggcggc agttgctggg cctactcccc 840  
 gagcacatgg cggagaagct gtgtgaggcc tgggcctttg ggagagcca ccagacgggc 900  
 gtcgtggcac tgggcctact cacctgcctg ctggcaatgc tgctggctgg ccgcatcagg 960



```

ctccggagga tcgatgcctt ctgcacctgc ctgtgggccc tgctgctggg gctgcacctg 1020
gctgaacagc acctgcaggc cgcctcgctt agctgggctaa acacgctcaa gttcagcacc 1080
acatctttgt gctgcctggg tggcttcacg gcggtgtgtg ccacaaggaa ggcaacgggc 1140
ccacggaggt tccggccccg aaggtcagag aagcagccat gactgcgggg ggaggacaca 1200
cggatgctca gggccaggct ttgccaggtc cgaagcgggc cctctctgt cctgcctctt 1260
ttcacctgct cacgcccctc cacccccacc ctacagcccc aggtcctggc ccagtcctctc 1320
cactgcctcg aagagtcagt ctgccctgcc ttttcctttc gggcaccacc agccatcccc 1380
gagtggcctg tagccactca ccactgctgc cacctctctg gccaatggcc ctttactggg 1440
cctgggactt ggaatgtggg cagcggccac acaggctctg gcccattggc tcctactggc 1500
agctccaggc acccccctct caccacggcg tttgctggct ctgacctgt tgggtgaggg 1560
tcctggctct gctgtcttcc cttctggcct ctgcacaggg gtggtgacag tggctacagg 1620
ctggggccct ggcgtgccct gaccgtgcag cagagtggag ctggggcagc agagagcccc 1680
agcctcacc ctagggagca cctgtgtct gtccccctgg tcctgcttat ggctggaccg 1740
gccctgcagg aggtgggtga gccgtgaagg aggcagagct gcagctctgg ctgctgcttg 1800
gcctctgctt ccaagacctt cccgagtcct cggaaatgga gagtgcagtt cttgggcccc 1860
gctgggcctt cgccatgagt ttggggagcg agaccccacc tgagacaggc agtaggagcc 1920
tgtgtgacc ttgggggaatc tgagcttttc caagggttaag gggcccaggg tatgacggc 1980
ttcagtgaac tcaggctggt gtcactcctt cctcctctga cctgtcacga gcctctgcag 2040
gtgcctgctc acctatggcc agcggcactc tgtctctcga ctcagggtgag ggggcagccc 2100
acagacctgc tcctcagtag cagggcctgg ccaggcccct gctgttctca gcctcagttt 2160
gccatctatg aaatgaggtg gacccctctc catagccctt ggggtgccagc tcagtgggtg 2220
tggggatcac atgaggtggc tcatgaggac aactcttgga agtcgagggg ctgccacgtg 2280
cagaggaagt tcccggcctg ggggctttat ccagggttcc cagtcgagag tggcccaggg 2340
ccgtccctca ccgggcatgt tccctctggc tgcccactcc ctacaggggc acatgtcctg 2400
ccactcgcca ctctgagcac gagttcacct tccagatgtg gccagggtgt gccagctcct 2460
ctctcctgtg cgttgggaacc ccgggggagg caagagcaga tcacagggtc atgagggtta 2520
cacccgtcac ctgggtctgc cgggatgggt tgggggggca ggtgccaggc ctactgctg 2580
tgaatctgcc acgcctgggg gtcttagagg ctgccccacc ccagtgattg ggtagcagct 2640
cacatcccac ccagcttcac aagtgaggaa cccagggtga tcggggagacc ctggggggct 2700
tctgtggcct ctgtgcccga tgacctgcgt ggcttcagac aaggccccag cgttactggg 2760
ctcagcttgt tgttctgtgt ggagcgtgag gtgagaaaac cctctgaaa agatgtgtgt 2820
ggggccacgc tcccactgg ttctgcagtg aggagttggg gcgggtgagc caagcgggc 2880
ccccatggtg tctacctgag gggcagggaa ccgcctgcct gtgcactcac gccaccccc 2940
agcccacaaa gagcccatct gagagaagga cgtggtggag ccaggacggg aaagcgtcct 3000
gtcggctggc catgctgttg cttgcgtctc gaatcttcgg ttctcgagga agtgttgaca 3060
gtgtgatgct aatgtctgct tttcttggcg ttgggtagaa gcaggacatc tgtgtgtatg 3120
tgcgatttta aattagatta ttataataa ccagagccag cctcgcgct ggccaggatc 3180
ctcctgccga gctgatgtcg ctctgcctc ctgcccgggt ccggaagcga catctcagga 3240
ggtagctctc agcagagtga ggattctgc cttctgtaga gttttgtgtg actttttaa 3300
ttattcatgt gtcccttaaa agtttacta cgtggagaaa attccagcac caagtgttgt 3360
ggcaacagct gagagagtgc aggcaccact gtgttgtggc ttgttgaccg ggaatgtgtc 3420
accctgcca gggaactctt ctctcgcgg gggacttggg atggccatca gaccttctag 3480
ggtctggtg gggtcacct aggtatgggt gaccgtccct gagacataag cgaggtagat 3540
tcagccatcc tcaccctcag acttgaggtc cccaccagg ccaagccggc ccccgtagc 3600
ccttgccctg gagcaaaccg ccaggacgca gcctccacgc cgcacctgcc acattcagcc 3660
ctgcccagga aggaacacat gaccctctc tctgtgactg ttgtgagtc tctgtctcat 3720
gtcgtagaat tgtggataat tgtctagtga cctctcatc actgtaacca tcgcgcctgg 3780
cctagatgtc gtgttttggg tgctgtgttt tcaataaatg cctctggggc cctgctttta 3840
caaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3900
a 3901

```

&lt;210&gt; 92

&lt;211&gt; 392

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 92

```

Met Glu Gly Val Ser Ala Leu Leu Ala Arg Cys Pro Thr Ala Gly Leu
  1                   5                   10                   15

```

Ala Gly Gly Leu Gly Val Thr Ala Cys Ala Ala Ala Gly Val Leu Leu  
                     20                    25                    30  
 Tyr Arg Ile Ala Arg Arg Met Lys Pro Thr His Thr Met Val Asn Cys  
                     35                    40                    45  
 Trp Phe Cys Asn Gln Asp Thr Leu Val Pro Tyr Gly Asn Arg Asn Cys  
                     50                    55                    60  
 Trp Asp Cys Pro His Cys Glu Gln Tyr Asn Gly Phe Gln Glu Asn Gly  
                     65                    70                    75                    80  
 Asp Tyr Asn Lys Pro Ile Pro Ala Gln Tyr Leu Glu His Leu Asn His  
                     85                    90                    95  
 Val Val Ser Ser Ala Pro Ser Leu Arg Asp Pro Ser Gln Pro Gln Gln  
                     100                    105                    110  
 Trp Val Ser Ser Gln Val Leu Leu Cys Lys Arg Cys Asn His His Gln  
                     115                    120                    125  
 Thr Thr Lys Ile Lys Gln Leu Ala Ala Phe Ala Pro Arg Glu Glu Gly  
                     130                    135                    140  
 Arg Tyr Asp Glu Glu Val Glu Val Tyr Arg His His Leu Glu Gln Met  
                     145                    150                    155                    160  
 Tyr Lys Leu Cys Arg Pro Cys Gln Ala Ala Val Glu Tyr Tyr Ile Lys  
                     165                    170                    175  
 His Gln Asn Arg Gln Leu Arg Ala Leu Leu Ser His Gln Phe Lys  
                     180                    185                    190  
 Arg Arg Glu Ala Asp Gln Thr His Ala Gln Asn Phe Ser Ser Ala Val  
                     195                    200                    205  
 Lys Ser Pro Val Gln Val Ile Leu Leu Arg Ala Leu Ala Phe Leu Ala  
                     210                    215                    220  
 Cys Ala Phe Leu Leu Thr Thr Ala Leu Tyr Gly Ala Ser Gly His Phe  
                     225                    230                    235                    240  
 Ala Pro Gly Thr Thr Val Pro Leu Ala Leu Pro Pro Gly Gly Asn Gly  
                     245                    250                    255  
 Ser Ala Thr Pro Asp Asn Gly Thr Thr Pro Gly Ala Glu Gly Trp Arg  
                     260                    265                    270  
 Gln Leu Leu Gly Leu Leu Pro Glu His Met Ala Glu Lys Leu Cys Glu  
                     275                    280                    285  
 Ala Trp Ala Phe Gly Gln Ser His Gln Thr Gly Val Val Ala Leu Gly  
                     290                    295                    300  
 Leu Leu Thr Cys Leu Leu Ala Met Leu Leu Ala Gly Arg Ile Arg Leu  
                     305                    310                    315                    320  
 Arg Arg Ile Asp Ala Phe Cys Thr Cys Leu Trp Ala Leu Leu Leu Gly  
                     325                    330                    335

Leu His Leu Ala Glu Gln His Leu Gln Ala Ala Ser Pro Ser Trp Leu  
 340 345 350

Asn Thr Leu Lys Phe Ser Thr Thr Ser Leu Cys Cys Leu Val Gly Phe  
 355 360 365

Thr Ala Ala Val Ala Thr Arg Lys Ala Thr Gly Pro Arg Arg Phe Arg  
 370 375 380

Pro Arg Arg Ser Glu Lys Gln Pro  
 385 390

<210> 93

<211> 2203

<212> DNA

<213> Homo sapiens

<400> 93

```

cagcgggtggg aggcggcgac cagccgggttg aggcggcgag cttggcctca ccacaatgtg 60
gcacgaggct cggaagcatg agcggaaagt tgcaggcatg atgggtcgact acaagaagag 120
ggcggagcgg agacgggagt attatgaaaa gatcaagaag gacccagccc agttcctgca 180
ggtacatggc cgagcttgca aggtgcacct ggattctgca gtcgccctgg ccgctgagag 240
ccctgttaat atgatgcctt ggcaggggga caccaacaac atgattgacc gattcgatgt 300
ccgtgcccac ctggaccaca tccccgacta cccccccct ctgctcacca ccattctccc 360
agaacaggag tcggacgaac ggaagtgtaa ctacgagcgc tacagaggcc tggtcagaa 420
cgactttgcc ggcattctcag aggagcagtg cctgtaccag atctacattg atgagttgta 480
cggaggcctc cagagaccca gcgaagatga gaagaagaag ctggcagaga agaaggcttc 540
catcggttat acctacgagg acagcacggt ggccgaggtg gagaaggcgg cagaaaagcc 600
agaggaggag gagtacgagg ccgaggaggga gagcaactcg gacgaagatg aggtcatccc 660
cgacatcgac gtggagggtg acgtggatga attgaaccag gagcagggtg cagatctcaa 720
caaacaggcc acgacttatg gcatggccga cgggtgactt gtcaggatgc tccggaaaga 780
caaggaggag gcagaggcca tcaagcatgc caaggctctt gaggaggaga aggccatgta 840
ctcgggacgc cgctctcgac gccagcggag agagtctcgg gagaagcggc tgaggggtcg 900
caagatcagc caccacagct atgcccgccg agacagcccc acctatgacc cctataagcg 960
gtcacccctc gagtccagct cagagtcccg ctcccgctcc cgctcccgca ccccgggccg 1020
cgaggagaag atcacgttca tcaccagttt tgggggcagc gatgaggagg cagccgcagc 1080
cgctgctgcc gcagcagcat caggagtcac cacagggaag ccccccgcac ctcccagcc 1140
tggcgggccc gccccgggac gtaatgccag cgcccgccgc cgctcctect cctcctectc 1200
ctcctcttct gcctcgagga cctccagctc ccgctccagc tctcgctcca gctcccgctc 1260
tcgcccgtgt gggggctact accgttccgg ccgccacgcc cgctcccggt ccgctcctcg 1320
gtcccgcctc cgctcccgct cccggcgcta ttcccggctc cgtagccgtg gccggcggca 1380
ctcagggtgg ggctcccag agggacaccg gtactcccgc tcgcccgcgc gccgtggtgg 1440
ttacggggcc cggcgagaa gcaggagccg ctcccactca ggggaccgct acaggcgggg 1500
cggccggggc ctcaggcacc acagcagtag ccgcagccgc agcagctggt ccctcagccc 1560
gtcccgcagt cgcagcctga ctgcagccg cagccatagc cccagcccca gccagagccg 1620
cagccgcagc cgcagccgca gccagagccc ctgcctatca cccgcaagag agaagctgac 1680
caggccggcc gcgtcccctg ctgtgggcga gaagctgaaa aagaccgaac ctgccgctgg 1740
taaagagaca ggagctgcca aacccaagct gacgcctcag gagaagctga aactgaggat 1800
gcagaaggcg ctgaacaggc agttcaaggc ggataagaag gcggcacaag aaaagatgat 1860
ccagcaggag catgagcggc aggagcggga agacagctt cgagccatgg cccgcaagat 1920
ccgcatgaag gagcgggaac gccgagagaa ggagagagaa gagtgggaac gccagtacag 1980
ccggcagagc cgctcaccct cccccgata cagtcgagaa tacagctctt ctccaaggcg 2040
ctcaaggtcc cgatcccga gccccatta ccgacattag gcagaagagt ggggggtggg 2100
gaggacaagg ggggtggtaa ggggtcgaag ctgtgatgct gctggtttta tctctagtga 2160
aataaagtca aaagtatttt aattcccgtc aaaaaaaaaa aaa 2203

```

<210> 94

<211> 674

<212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 94

```

Met Trp His Glu Ala Arg Lys His Glu Arg Lys Leu Arg Gly Met Met
 1             5             10             15

Val Asp Tyr Lys Lys Arg Ala Glu Arg Arg Arg Glu Tyr Tyr Glu Lys
          20             25             30

Ile Lys Lys Asp Pro Ala Gln Phe Leu Gln Val His Gly Arg Ala Cys
          35             40             45

Lys Val His Leu Asp Ser Ala Val Ala Leu Ala Ala Glu Ser Pro Val
          50             55             60

Asn Met Met Pro Trp Gln Gly Asp Thr Asn Asn Met Ile Asp Arg Phe
 65             70             75             80

Asp Val Arg Ala His Leu Asp His Ile Pro Asp Tyr Thr Pro Pro Leu
          85             90             95

Leu Thr Thr Ile Ser Pro Glu Gln Glu Ser Asp Glu Arg Lys Cys Asn
          100             105             110

Tyr Glu Arg Tyr Arg Gly Leu Val Gln Asn Asp Phe Ala Gly Ile Ser
          115             120             125

Glu Glu Gln Cys Leu Tyr Gln Ile Tyr Ile Asp Glu Leu Tyr Gly Gly
          130             135             140

Leu Gln Arg Pro Ser Glu Asp Glu Lys Lys Lys Leu Ala Glu Lys Lys
          145             150             155             160

Ala Ser Ile Gly Tyr Thr Tyr Glu Asp Ser Thr Val Ala Glu Val Glu
          165             170             175

Lys Ala Ala Glu Lys Pro Glu Glu Glu Glu Ser Ala Ala Glu Glu Glu
          180             185             190

Ser Asn Ser Asp Glu Asp Glu Val Ile Pro Asp Ile Asp Val Glu Val
          195             200             205

Asp Val Asp Glu Leu Asn Gln Glu Gln Val Ala Asp Leu Asn Lys Gln
          210             215             220

Ala Thr Thr Tyr Gly Met Ala Asp Gly Asp Phe Val Arg Met Leu Arg
          225             230             235             240

Lys Asp Lys Glu Glu Ala Glu Ala Ile Lys His Ala Lys Ala Leu Glu
          245             250             255

Glu Glu Lys Ala Met Tyr Ser Gly Arg Arg Ser Arg Arg Gln Arg Arg
          260             265             270

Glu Phe Arg Glu Lys Arg Leu Arg Gly Arg Lys Ile Ser Pro Pro Ser
          275             280             285

Tyr Ala Arg Arg Asp Ser Pro Thr Tyr Asp Pro Tyr Lys Arg Ser Pro
          290             295             300

```

Ser Glu Ser Ser Ser Glu Ser Arg Ser Arg Ser Arg Ser Pro Thr Pro  
 305 310 315 320  
 Gly Arg Glu Glu Lys Ile Thr Phe Ile Thr Ser Phe Gly Gly Ser Asp  
 325 330 335  
 Glu Glu Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ser Gly Val Thr  
 340 345 350  
 Thr Gly Lys Pro Pro Ala Pro Pro Gln Pro Gly Gly Pro Ala Pro Gly  
 355 360 365  
 Arg Asn Ala Ser Ala Arg Arg Arg Ser Ser Ser Ser Ser Ser Ser  
 370 375 380  
 Ser Ala Ser Arg Thr Ser Ser Ser Arg Ser Ser Ser Arg Ser Ser Ser  
 385 390 395 400  
 Arg Ser Arg Arg Gly Gly Gly Tyr Tyr Arg Ser Gly Arg His Ala Arg  
 405 410 415  
 Ser Arg Ser Arg Ser Trp Ser Arg Ser Arg Ser Arg Ser Arg Arg Tyr  
 420 425 430  
 Ser Arg Ser Arg Ser Arg Gly Arg Arg His Ser Gly Gly Gly Ser Arg  
 435 440 445  
 Asp Gly His Arg Tyr Ser Arg Ser Pro Ala Arg Arg Gly Gly Tyr Gly  
 450 455 460  
 Pro Arg Arg Arg Ser Arg Ser Arg Ser His Ser Gly Asp Arg Tyr Arg  
 465 470 475 480  
 Arg Gly Gly Arg Gly Leu Arg His His Ser Ser Ser Arg Ser Arg Ser  
 485 490 495  
 Ser Trp Ser Leu Ser Pro Ser Arg Ser Arg Ser Leu Thr Arg Ser Arg  
 500 505 510  
 Ser His Ser Pro Ser Pro Ser Gln Ser Arg Ser Arg Ser Arg Ser Arg  
 515 520 525  
 Ser Gln Ser Pro Ser Pro Ser Pro Ala Arg Glu Lys Leu Thr Arg Pro  
 530 535 540  
 Ala Ala Ser Pro Ala Val Gly Glu Lys Leu Lys Lys Thr Glu Pro Ala  
 545 550 555 560  
 Ala Gly Lys Glu Thr Gly Ala Ala Lys Pro Lys Leu Thr Pro Gln Glu  
 565 570 575  
 Lys Leu Lys Leu Arg Met Gln Lys Ala Leu Asn Arg Gln Phe Lys Ala  
 580 585 590  
 Asp Lys Lys Ala Ala Gln Glu Lys Met Ile Gln Gln Glu His Glu Arg  
 595 600 605  
 Gln Glu Arg Glu Asp Glu Leu Arg Ala Met Ala Arg Lys Ile Arg Met  
 610 615 620

Lys Glu Arg Glu Arg Arg Glu Lys Glu Arg Glu Glu Trp Glu Arg Gln  
 625 630 635 640

Tyr Ser Arg Gln Ser Arg Ser Pro Ser Pro Arg Tyr Ser Arg Glu Tyr  
 645 650 655

Ser Ser Ser Arg Arg Arg Ser Arg Ser Arg Ser Arg Ser Pro His Tyr  
 660 665 670

Arg His

<210> 95

<211> 1014

<212> DNA

<213> Homo sapiens

<400> 95

```

gggcgcgcgc gcctcctcct ccatggctgt ttaccgcgct gcattgtggg agtttgacct 60
ccgcgcgcgc caaccgcgc ctcagcttgc gccgcgcgc ccgcgcgcgc cgccgcgcac 120
gccatgggag ccgtgactga cgacgaagtt atacggaagc gtctcctcat tgatggagat 180
ggtgctggag atgatcggag aattaatctg ctagtgaaga gtttcattaa atggtgcaac 240
tctgggtccc aggaagaggg atatatgccag taccaacgta tgctgagcac gctgtctcaa 300
tgtgaatttt caatgggcaa aactttacta gtatatgata tgaatctcag agaaatggaa 360
aattatgaaa aaatttacia ggaaatagaa tgtagcatag ctggagcaca tgaaaaaatt 420
gctgagtgcg aaaagcaaat tcttcaagca aaacgaatac gaaaaaatcg ccaagaatat 480
gatgcttttg caaaagtgat tcagcaccat ccagacaggc atgagacatt aaaggaacta 540
gaggctctgg gaaaagaatt agagcatctt tcacacatta aagaaagtgt tgaagataag 600
ctggaattga gacggaaaca gtttcatgtt cttcttagta ccatccatga acttcagcaa 660
acattggaaa atgatgaaa actctcagag gtagaagaag ctcaggaagc aagcatggaa 720
acagatccta agccatagac aggctaattg cccaccactc ccaggaatat tgaaatagct 780
acatgaccat aatgtgttta aaatgtggta tgctcttgag atattttaaag ttttggcagt 840
aaaatactct gtttttaagt atgaatgtat ttcattcata tttcctctca caaagggaaa 900
tgacttcagt atagatttgt ttttattaaa atgcattttt tattcttaag tggtaggaag 960
caacatccaa aaatgcttaa taaaatgctt ttaagctgca aaaaaaaaaa aaaa 1014

```

<210> 96

<211> 204

<212> PRT

<213> Homo sapiens

<400> 96

Met Gly Ala Val Thr Asp Asp Glu Val Ile Arg Lys Arg Leu Leu Ile  
 1 5 10 15

Asp Gly Asp Gly Ala Gly Asp Asp Arg Arg Ile Asn Leu Leu Val Lys  
 20 25 30

Ser Phe Ile Lys Trp Cys Asn Ser Gly Ser Gln Glu Glu Gly Tyr Ser  
 35 40 45

Gln Tyr Gln Arg Met Leu Ser Thr Leu Ser Gln Cys Glu Phe Ser Met  
 50 55 60

Gly Lys Thr Leu Leu Val Tyr Asp Met Asn Leu Arg Glu Met Glu Asn  
 65 70 75 80

Tyr Glu Lys Ile Tyr Lys Glu Ile Glu Cys Ser Ile Ala Gly Ala His  
 85 90 95

Glu Lys Ile Ala Glu Cys Lys Lys Gln Ile Leu Gln Ala Lys Arg Ile  
 100 105 110  
 Arg Lys Asn Arg Gln Glu Tyr Asp Ala Leu Ala Lys Val Ile Gln His  
 115 120 125  
 His Pro Asp Arg His Glu Thr Leu Lys Glu Leu Glu Ala Leu Gly Lys  
 130 135 140  
 Glu Leu Glu His Leu Ser His Ile Lys Glu Ser Val Glu Asp Lys Leu  
 145 150 155 160  
 Glu Leu Arg Arg Lys Gln Phe His Val Leu Leu Ser Thr Ile His Glu  
 165 170 175  
 Leu Gln Gln Thr Leu Glu Asn Asp Glu Lys Leu Ser Glu Val Glu Glu  
 180 185 190  
 Ala Gln Glu Ala Ser Met Glu Thr Asp Pro Lys Pro  
 195 200

<210> 97  
 <211> 955  
 <212> DNA  
 <213> Homo sapiens

<400> 97  
 aatcctcaac aaaatagtag caaaacaaat ccaatagtag atcacgaaga taatacagta 60  
 tgatcaaatg ggatttattt caaggatgca tagatgattc aacattcacg atcaataaaa 120  
 tttattctgt taatttttca taaggatgt ttttaaataa gaatgggtatt gtatattaga 180  
 cataaaaatga ctgttttagt tagcattctt agagctagct tcataatcca attaatatac 240  
 ttgcaacttg agtgcaggtg ttttaatttt tataactgta tcctgtatgc tattcaaatg 300  
 agctaattgt agttattctt ataccattg gtattgggtt ccatagtata cataagtttt 360  
 atttttgttt ttctgttag accttcaaat atttactttc catagtttct ctggcataaa 420  
 agctcccagt ttctatcttc aacagttcag gtcttgggat atctatcgtt ttatttggtt 480  
 ttataacttt tatttagaag agttacatcc tttttagctt atttaatgat aaaaagttca 540  
 cttttccac ttttgatttt gaatgaattg ctgcccctaa catggatcta tcttggtttt 600  
 acagagagaa gagaagagga agatattgaa gagaagaaat cgattaagaa aaaaattaaa 660  
 gaacttaagt ttttagattc taaaattgcc cagaaccttt gtaagtatca tattccaata 720  
 ccattcaagg acagtggaaa tttttcttta aatgatttca ttttcttta gaccgattat 780  
 tcattatttg ctattttcat tttgttatta tatgcatgat aaattcacag ataactctcc 840  
 tttaggtaaa ttatgggatt aaatgcttca aaagataagt gcatattaga aaatacaaat 900  
 aagaagaggt tttaaaatga aattctacct ttcataactg aaaaaaaaaa aaaaa 955

<210> 98  
 <211> 97  
 <212> PRT  
 <213> Homo sapiens

<400> 98  
 Met Ile Lys Ser Ser Leu Phe Pro Leu Leu Tyr Leu Asn Glu Leu Leu  
 1 5 10 15  
 Pro Leu Thr Trp Ile Tyr Leu Gly Phe Thr Glu Arg Arg Glu Glu Glu  
 20 25 30  
 Asp Ile Glu Glu Lys Lys Ser Ile Lys Lys Lys Ile Lys Glu Leu Lys  
 35 40 45

Phe Leu Asp Ser Lys Ile Ala Gln Asn Leu Cys Lys Tyr His Ile Pro  
 50 55 60

Ile Pro Phe Lys Asp Ser Gly Asn Ile Ser Leu Asn Asp Phe Ile Phe  
 65 70 75 80

Phe Lys Thr Asp Tyr Ser Leu Phe Ala Ile Phe Ile Leu Leu Leu Tyr  
 85 90 95

Ala

<210> 99

<211> 1375

<212> DNA

<213> Homo sapiens

<400> 99

```

gtcttctttt agggagcagg agtgcactctg gtaattgagg gtggatgttg tgtgtgctgg 60
ggaggggtcc ttctgttttg tgctaccctt gtctactctg cccctggatg gtgcggggtg 120
ctttctccac cccacactc cctgctcagc tcctcgtgct gccctgcatg cccaggcctt 180
tgagccaagc tgcttttttg ggcagggagt agcagcagggt gggaggggtt acccatcagc 240
ccttgcaagt cccccactca ggctcttgga aggtccagggt atgggctctg atgagagggt 300
aaaagatgct cagggaaaca caggcctcag ctgcctagag gaccctcccc ctgecttgca 360
gtgggctcgg gtagagcagt atcaggagct agggttgtct gctgcccaca ctctgtcttt 420
ttgggatatc taactgctaa ggagggagtt gacatcccc ttctggctca tgtgtctgac 480
accaacaaca tgggtctctgt ccctctctct ttgactctcc ctttgtcctc cccatagagc 540
tggggtgggg tggatcccta tacctggggc aggcagcccc aaagtggggg agggggatgg 600
cagagactgt aaaggcgcca ctggactctg gcaaggcctt tattacctt actccctcc 660
ctctccatc accagctca aggcctgagg ggtgcagggg ctccctggcag ctactgggtg 720
aggtttcctg gcacagactc acccttctt ctggcaccac ctctttccct tttgaagaga 780
cagcaacagc cgtagcaaaa gcagctgctg ctctgtctat gaggggtgat atatttttta 840
cccaaagctc tggaattgta catttatctt ttaaaactca aagagggaaa gagccttgta 900
tcatatgtga acattgtatc ataggtaatg ttgtacagac ccttttatac agtgatctgt 960
cttgttctctg cagcaaaaat cctctatgga cataggaggt gctgtgtccc atgccctctt 1020
gccctgacag tgtcccatgg gccccttct gctccctgcc ccctccctgc tactgctgat 1080
gcactctctc ctccctgcag cccctggctt cccagccttc ctctgaccc cttccaacag 1140
ccttggaact ccagctgcca ccacctctg ggtcggacac tgggaccac tggcccagtc 1200
ttggctgctg cttacccta gccttgatgc ctgcccagg acccccagcc cctcccgtt 1260
gccctgcagc ttaacagag tgaaccatgt gtattgtaca ggcgcgggtg tcattgcaga 1320
aaccgctggg tggagaagaa gccgataaag tctatgaatc aaaaaaaaaa aaaaa 1375

```

<210> 100

<211> 132

<212> PRT

<213> Homo sapiens

<400> 100

Met Ala Glu Thr Val Lys Ala Pro Leu Asp Ser Gly Lys Ala Phe Ile  
 1 5 10 15

Thr Phe Thr Pro Leu Pro Leu Pro Ser Pro Ala Ser Arg Pro Glu Gly  
 20 25 30

Cys Arg Gly Ser Trp Gln Leu Leu Gly Glu Val Ser Trp His Arg Leu  
 35 40 45

Thr Leu Leu Ser Gly Thr Thr Ser Phe Pro Phe Glu Glu Thr Ala Thr



50

55

60

Ala Val Ala Lys Ala Ala Ala Pro Ala Met Arg Val Tyr Ile Phe  
65 70 75 80

Phe Thr Gln Ser Ser Gly Ile Val His Leu Phe Phe Lys Thr Gln Arg  
85 90 95

Gly Lys Glu Pro Cys Ile Ile Cys Glu His Cys Ile Ile Gly Asn Val  
100 105 110

Val Gln Thr Leu Leu Tyr Ser Asp Leu Ser Cys Ser Cys Ser Lys Asn  
115 120 125

Pro Leu Trp Thr  
130

&lt;210&gt; 101

&lt;211&gt; 1213

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 101

```

ggcttcaggt tgaagtcctt ggttcttcca gttcctcacg ggtaggtag gggctcctgc 60
atcaccttca gaatccagtt ccaacccccca ctctccttag gccttggtgct ctgctctgcc 120
ctgccaggtt gcccttggtc atgtgagtag catgggaggg tgggtggggac ggcagtggtg 180
atgaaggggg tgcaccacag gcctcatgaa gcagttccca catgggagtg tggctggggc 240
gtggccacca cagagcacat ggctgtgtct aggcgcaagc acttttagcag tatctgttta 300
catgcgcaag gatcaagccg actacctgtg ctgtctactg ggacagcagt ctccgagcta 360
ctccgtacct cctctgcca ggtcgtggag ttaggccccca gtccctactt gtcactggtt 420
cccactgtgc tctaaactgt gcagcacctg ggagctctgg cctggggctg gaggccctgg 480
taggagctgc agttggaggc cgttctgtgc ccagcagcgg tgagcggctc ccatggggcc 540
tgtgtctgca gggagccagg gctgcgccac atgtgtctgt aaactggcac ccacctggcg 600
tgctgtctgc gccacttgct tctgcagca cctcctaccc tgctccgtgt cctccctctc 660
cccgcgcctg gctcaggagt gctggaaaag ctacgcctc ggcctgggag cctggcctct 720
tgatatacct cgagcttccc ctgtgtctcc cagccccagg accactggcc ccttggcctg 780
aggggctggg ggccccacga cctgcagcgt cgagtcgggg agagagcccg gagcggcggtg 840
ccatctcggc tcggccttgc tgagagcctc cgccttgggt ttctccctgt ctggattcag 900
tggctcacgt tgggtgtaca cagctagaat agatatattt agagagagag atatttttaa 960
gacaaagccc acaattagct gtctttaaac accgcagaa cccctcccag aagaagagcg 1020
atccctcgga cggtcggggc gggcaccctc agccgggctc tttgcagaag cagcaccgct 1080
gactgtgggc ccggccctca gatgtgtaca tatacggcta tttcctattt tactgttctt 1140
cagatttagt acttgtaaat aaacacacac attaaggaga gattaaacat ttttgctaaa 1200
aaaaaaaaaa aaa 1213

```

&lt;210&gt; 102

&lt;211&gt; 100

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 102

```

Met Lys Gly Val His His Arg Pro His Glu Ala Val Pro Thr Trp Ala
  1           5           10           15

Cys Gly Trp Gly Val Ala Thr Thr Glu His Met Ala Val Ser Arg Arg
          20           25           30

Lys His Phe Ser Ser Ile Cys Leu His Ala Gln Gly Ser Ser Arg Leu
    35           40           45

```

Pro Val Leu Ser Thr Gly Thr Ala Val Ser Glu Leu Leu Arg Thr Ser  
 50 55 60

Leu Cys Gln Val Val Glu Leu Gly Pro Ser Pro Tyr Leu Ser Leu Val  
 65 70 75 80

Pro Thr Val Leu Leu Thr Val Gln His Leu Gly Ala Leu Ala Trp Gly  
 85 90 95

Trp Arg Pro Trp  
 100

<210> 103

<211> 1036

<212> DNA

<213> Homo sapiens

<400> 103

```
cctcaaatgc tttcttttctt cagatgcttt ttctgtgtaca tgatactagt agacactttt 60
ctcttttatat ttactgatag tgaaaatcat acgcaataaaa atattgatgt ttgaaggcag 120
tgggtcaccaa ttggttaaaaa aactatgaaa tgtaaaactga attggttatat ctctatcctt 180
tttgcttttc tctgtgtttt taatgtatgg aataaatctc ataaatagaa agaaaaataa 240
tctagaaatt tttcaaagct agtactcttt ctctttataa atgtacacaa ttttaatctt 300
tttacaattt tatttaactg tacctactgt acttattgta gattcaatga cgcagttaag 360
tcatcaccca aggatttatg aatttgagat tactgacctg ttttcttcat attgcattca 420
catcaatatt tgtgaatttg ttgttcagct tttcattcaa acaaaaaata ttccctcaag 480
aaagctccat ttttatcata aacatttcaa cataaccaac attagaacaa gtctgccatg 540
ttaaaaaataa tttaaagact tatctctgaa aacggtatcc agaaacgcag gtgttcccag 600
taatgtagct tcaaaaaataa aatgtgtctat ttatatgaca tgaaattcat aacttttgga 660
agggtatatt tatgacagca taaaaaataa attctgtgct ataaagaaga tccaacaaat 720
taaccatata agcacagaaa atagagaaac acagttattg aatctactct tgtcattaac 780
attttcaaaa aacaaaatgc atattgtaat atttggtaca tgacacttgc atgttgatat 840
gcctatatac ttacaaagta ttcaatgtgt acttagcggc gcttaaaata tgtcatgtac 900
aactcttata aacattttta cagggttccc atttgcactt catctttcag taaagtcttg 960
tcagaaaaaaa attgtctgat aaatatggaa aaataaaatt tgaatttttag ttaaaaaaaa 1020
aaaaaaaaaa aaaaaa 1036
```

<210> 104

<211> 87

<212> PRT

<213> Homo sapiens

<400> 104

Met Tyr Thr Ile Leu Ile Phe Leu Gln Ile Tyr Leu Thr Val Pro Thr  
 1 5 10 15

Val Leu Ile Val Asp Ser Met Thr Gln Leu Ser His His Pro Arg Ile  
 20 25 30

Tyr Glu Phe Glu Ile Thr Asp Leu Phe Ser Ser Tyr Cys Ile His Ile  
 35 40 45

Asn Ile Cys Glu Phe Val Val Gln Leu Phe Ile Gln Thr Lys Asn Ile  
 50 55 60

Pro Ser Arg Lys Leu His Phe Tyr His Lys His Phe Asn Ile Thr Asn  
 65 70 75 80

Ile Arg Thr Ser Leu Pro Cys

85

&lt;210&gt; 105

&lt;211&gt; 2349

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 105

```

tttttttttt tttttggatt cttggtaaaa ttttatccaa aaaacaggat acatatatat 60
ttagagaagg aaatatgaaa tcaagagttt tggcagcccc tgcttttttt ttttttttag 120
ctccctaaag actgtagcag gataaaagga tcaactggctc cgagtctctt tgagataaca 180
agtgatgaaa taaaaaagaa agcccatacc ctcaaataag gtcaggtaac cccattgccc 240
accctcccta caaggtaaaa aatgagtact tttagtaaca gttcagaatt catctttatc 300
tcctacctgc ctcatcgggtg gaagtgtaaa gtcatgattt ttttttagaca ttgatacttg 360
tgtctataga caaataaact catattagac ggccaaagag gcctaccact gctgcaccag 420
cagtatacct cactgactgc ctccactctg cccctgcgcc cagatgctcc tgttgaaaag 480
tcacccgagg agacagctac ccaggtcccc agtctggaga gtctgacttt aaagctagag 540
cacgaggtgg tggccaggag ccgaccaacc ccacaagact atgagatgag agtatccccc 600
tctgatacta cccctctggt ttcccgaggt gttccaccag tcaaactgga ggatgaggat 660
gattcggact ctgagctgga cttgagcaag ctgtcaccat cttcttcttc ttctcatcc 720
tcattccagct ccagctccag cactgatgag agtgaggatg agaaggaaga gaagctaact 780
gaccagtccc gctcaaagct ctatgatgaa gagagtctcc tgctccctcac tatgtcccaa 840
gatggattcc caaatgaaga tggagaacaa atgaccctcg agcttctgct actgcaggaa 900
agacaaagag cctctgagtg gcccaaggat cgtgtcctga taaaccgtat tgacctcgctc 960
tgccaggctg tactctcagg gaagtggcct tctagccgta ggagccagga aatggttaaca 1020
ggaggaatth tggggccagg caaccacttg ctagacagtc cctcattgac tcctggagaa 1080
tatgggtgact ctccagtccc cacaccacga agtagtagtg cagcttccat ggcagaggag 1140
gaagcatctg cagtcagcac agcggcagcc cagttcacca aacttcgccg aggcattggat 1200
gaaaaggagt ttacagttca aatcaaagat gaggaaggat tgaagttaac attccagaag 1260
cacaagttga tggcgaatgg agtaatggga gatggacatc cactgtttca taagaagaag 1320
gggaacagaa agaagctagt agagctggag gtggagtga tggagagacc taatcacctt 1380
gatgtggacc tggagacccg gatccctgtc atcaataagg tggatggtag tttgctggtg 1440
ggtgaggatg cccctcgccg ggctgaactg gagatgtggt tacagggtca tccagagttt 1500
gctgttgatc cccgatttct agcgtatatg gaggatcgca gaaaacagaa gtggcaaaaga 1560
tgtaaaaaaa ataataaggc agaattgaac tgtttgggaa tggaaaccagt acagacagct 1620
aactctagaa atgggaaaaa ggtcatcac actgaaacgg tgttcaaccg ggttttgcca 1680
gggcctattg caccagagag cagcaagaag cgggcccgtg ggatgagacc agacctttct 1740
aagatgatgg ccctcatgca ggggtgaagc actgggtctc tatctctgca taacacgttc 1800
caacacagca gtagtggcct acagtctgtg tcatcttgg gtcacagcag tgccacttct 1860
gcattcttgc cttttatgcc atttgtgatg ggtggtgcac catcatcccc tcatgtagac 1920
tccagcacca tgcttcatca ccaccaccac caccaccacc cccaccatca ccaccatcac 1980
catccaggct tgagagcccc tggctacccc tcttcaccag tgactaccgc ctctggtact 2040
accttgcggt tgccaccact gcaacctgag gaggatgacg atgaggatga agaagatgat 2100
gatgacttat ctcagggcta tgatagctca gaaagggact tctcactcat tgatgatcct 2160
atgatgccag ctaactcaga ctccagtga gatgctgatg actgaagccc cagcatgggc 2220
cccattgctt gggcggtgc tgatttttca ttactctgg cccttgagct atggaaacgt 2280
gggaggggca ggggagatgt ggggaagtcc aggactccag gaggtgaaaa ggaacaaaaa 2340
aaaaaaaaa                                     2349

```

&lt;210&gt; 106

&lt;211&gt; 539

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 106

```

Met Arg Val Ser Pro Ser Asp Thr Thr Pro Leu Val Ser Arg Ser Val
  1                      5                      10                      15

```

Pro Pro Val Lys Leu Glu Asp Glu Asp Asp Ser Asp Ser Glu Leu Asp  
                   20                                  25                                  30  
 Leu Ser Lys Leu Ser Pro Ser Ser Ser Ser Ser Ser Ser Ser Ser  
                   35                                  40                                  45  
 Ser Ser Ser Ser Thr Asp Glu Ser Glu Asp Glu Lys Glu Glu Lys Leu  
                   50                                  55                                  60  
 Thr Asp Gln Ser Arg Ser Lys Leu Tyr Asp Glu Glu Ser Leu Leu Ser  
                   65                                  70                                  75                                  80  
 Leu Thr Met Ser Gln Asp Gly Phe Pro Asn Glu Asp Gly Glu Gln Met  
                                   85                                  90                                  95  
 Thr Pro Glu Leu Leu Leu Gln Glu Arg Gln Arg Ala Ser Glu Trp  
                   100                                  105                                  110  
 Pro Lys Asp Arg Val Leu Ile Asn Arg Ile Asp Leu Val Cys Gln Ala  
                   115                                  120                                  125  
 Val Leu Ser Gly Lys Trp Pro Ser Ser Arg Arg Ser Gln Glu Met Val  
                   130                                  135                                  140  
 Thr Gly Gly Ile Leu Gly Pro Gly Asn His Leu Leu Asp Ser Pro Ser  
                   145                                  150                                  155                                  160  
 Leu Thr Pro Gly Glu Tyr Gly Asp Ser Pro Val Pro Thr Pro Arg Ser  
                                   165                                  170                                  175  
 Ser Ser Ala Ala Ser Met Ala Glu Glu Glu Ala Ser Ala Val Ser Thr  
                   180                                  185                                  190  
 Ala Ala Ala Gln Phe Thr Lys Leu Arg Arg Gly Met Asp Glu Lys Glu  
                   195                                  200                                  205  
 Phe Thr Val Gln Ile Lys Asp Glu Glu Gly Leu Lys Leu Thr Phe Gln  
                   210                                  215                                  220  
 Lys His Lys Leu Met Ala Asn Gly Val Met Gly Asp Gly His Pro Leu  
                   225                                  230                                  235                                  240  
 Phe His Lys Lys Lys Gly Asn Arg Lys Lys Leu Val Glu Leu Glu Val  
                                   245                                  250                                  255  
 Glu Cys Met Glu Glu Pro Asn His Leu Asp Val Asp Leu Glu Thr Arg  
                   260                                  265                                  270  
 Ile Pro Val Ile Asn Lys Val Asp Gly Thr Leu Leu Val Gly Glu Asp  
                   275                                  280                                  285  
 Ala Pro Arg Arg Ala Glu Leu Glu Met Trp Leu Gln Gly His Pro Glu  
                   290                                  295                                  300  
 Phe Ala Val Asp Pro Arg Phe Leu Ala Tyr Met Glu Asp Arg Arg Lys  
                   305                                  310                                  315                                  320  
 Gln Lys Trp Gln Arg Cys Lys Lys Asn Asn Lys Ala Glu Leu Asn Cys  
                                   325                                  330                                  335

Leu Gly Met Glu Pro Val Gln Thr Ala Asn Ser Arg Asn Gly Lys Lys  
 340 345 350  
 Gly His His Thr Glu Thr Val Phe Asn Arg Val Leu Pro Gly Pro Ile  
 355 360 365  
 Ala Pro Glu Ser Ser Lys Lys Arg Ala Arg Arg Met Arg Pro Asp Leu  
 370 375 380  
 Ser Lys Met Met Ala Leu Met Gln Gly Gly Ser Thr Gly Ser Leu Ser  
 385 390 395 400  
 Leu His Asn Thr Phe Gln His Ser Ser Ser Gly Leu Gln Ser Val Ser  
 405 410 415  
 Ser Leu Gly His Ser Ser Ala Thr Ser Ala Ser Leu Pro Phe Met Pro  
 420 425 430  
 Phe Val Met Gly Gly Ala Pro Ser Ser Pro His Val Asp Ser Ser Thr  
 435 440 445  
 Met Leu His His His His His His Pro His Pro His His His His His  
 450 455 460  
 His His Pro Gly Leu Arg Ala Pro Gly Tyr Pro Ser Ser Pro Val Thr  
 465 470 475 480  
 Thr Ala Ser Gly Thr Thr Leu Arg Leu Pro Pro Leu Gln Pro Glu Glu  
 485 490 495  
 Asp Asp Asp Glu Asp Glu Glu Asp Asp Asp Asp Leu Ser Gln Gly Tyr  
 500 505 510  
 Asp Ser Ser Glu Arg Asp Phe Ser Leu Ile Asp Asp Pro Met Met Pro  
 515 520 525  
 Ala Asn Ser Asp Ser Ser Glu Asp Ala Asp Asp  
 530 535

&lt;210&gt; 107

&lt;211&gt; 3004

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 107

ggggcatgag catctcaggg ctgccagaat ggcttttggct gaggatcatag caccagcgtg 60  
 tgtcatgtct tggctgcgtt tctggggccc atggcccctc cttacgtggc aactattgtc 120  
 ttactagtc aaggaggctc agcctctggt gtgggtcaag gaccgctcc agctgacctc 180  
 taacccctg gggccacctg agcctggtc ttcccgtcc tcccatctcc catgggaatc 240  
 tccccatgca cctgctcccc cagcagcccc gggggacttt gattacctgg ggccctctgc 300  
 ttcttcgcag atgtcagccc tgcctcagga accaactgaa aatttggtc cattcctgaa 360  
 ggaattggat tcagctggag agctgcccct ggggccagag ccgttcttgg ctgcacatca 420  
 ggacttaaat gacaagcgga ctccagaaga aaggctccca gaggtggttc cgcttctcaa 480  
 ccgggatcag aaccaggccc tagttcagct tctcgccctc aagtgggttc aaactacaga 540  
 tctagatcgg gctgcaggtc atcaggcaga tgaaatactt gttccactag acagtaaggt 600  
 ttcaagacca accaaatttg ttgtttcgcc caagaacctg aagaaagatc tagctgaacg 660  
 ttggagcctt cctgagattg ttgggattcc acaccaatta tccaaacctc agcgtcagaa 720  
 acagactttg ccagatgatt atttgagtat ggacacactg tatcccggca gcctacctcc 780  
 agaactccgg gtgaacgcag atgagcctcc agggcctcct gagcaagttg gactttctca 840

```

attccatcta gagcccaaaa gtcaaaatcc agagaccctt gaagacatcc agtcctcttc 900
actccaggaa gaagccccag cgcagcttct acagctccct caggaggtag aaccttcaac 960
ccagcaggag gccccagctc tgctccaga gtcctctatg gagagtctag ctcaaactcc 1020
actgaatcat gaagtgcacag ttcaacctcc aggtgaggat caagctcatt ataatttgcc 1080
caagttttaca gtcaaacctg cagatgtgga ggttaccatg acttcagagc ctaaaaatga 1140
gacagaatct acccaagccc agcaggaggc cccaattcag cctcccaggg aggcggaacc 1200
ttctttctaca gccctgagga ctacagatcc tctccagaa caccctgagg tgacacttcc 1260
accttcagac aagggtcagg ctacagattc acacctgact gaagccacag ttcaacctct 1320
ggacctggag cttagcataa ctacagagcc tactacagag gttaaaccgt ctccaaccac 1380
ggaggaaacc tcagctcagc ctccagacct ggggcttgcc ataactccag aaccactac 1440
agagattgga cattccacag ccctggagaa gactagagct cctcatccag accagggtca 1500
gactctgcat cgaagcctga ctgaagtcac aggtccacct acaaagttag aatcttcgca 1560
ggattcattg gtgcagtctg aaactgcacc agaggaacag aaggcctcca caagcaccac 1620
catatgtgag ctctgcacct gcggagatga gactctgtca tgtgttggtc tcagcccaaa 1680
gcagaggctc cgccaagtgc ctgtgccaga gcccgacacc tacaatggca tcttcaccac 1740
cttaaatctc caaggaaact atatttcata ccttgatgga aatgtatgga aagcatacag 1800
ttggaccgag aaactaattc tcagtgaata ttatttgact gaattacctt aggattcatt 1860
tgaaggcctg ctatacctcc agtatttaga ttatcctgc aataaaatac gatataattga 1920
aagacaaaca tttgaatcac taccattttt gcagtatata aatctgggct gcaattttaa 1980
tacaaaactg agccttgga ctttcaggc ctggcacgga atgcagtttt tacacaactt 2040
aattctcaat cgcaatcctc tgactactgt cgaagatcca tatctctttg aactgccggc 2100
attaaaatat ctagacatgg gaacaacaca catcacactt acaacactta agaacattct 2160
cacgatgact gttgaactgg aaaaactgat cttacctagc catatggcct gctgcctctg 2220
ccaattttaa aatagcattg aggtgtctg caagacagtc aagctgcatt gcaacactgc 2280
atgtctgact aacagcatac attgtcctga agaagcatct gtagggaatc cagaaggagc 2340
gttcatgaag atgttacaag cccggaagca gcacatgagc actcagctga ctattgagtc 2400
ggaggcgccc tcagacagca gtggcatcaa cttgtcaggc tttgggggtg atcagcttga 2460
aattcagcta accgagcagc tacggtccct catccccaac gaggatgtga gaaagttcat 2520
gtctcatgtt atccggacct tgaaaatgga atgttcagaa acacatgtgc aaggggagctg 2580
tgccaagctc atgtcgcgaa caggcctcct gatgaagctt ctcagcgagc agcaggaagc 2640
aaaggcattg aatgtagaat gggatacgga ccaacaaaaa acaaattata ttaatgagaa 2700
catggaacag aatgaacaga aagagcagaa gtcaagttag ctcatgaaag aagttccagg 2760
agatgactat aagaacaaac tcatcttcgc aatatctgtg actgtaatac taataatttt 2820
gattataatt ttttgtctta tagagggtgaa ttcacataaa agggcatcag aaaaatacaa 2880
agacaaccca tcaatatcag gagcctgagc atgagttaaa gcatgtggat ggcctggagc 2940
tatgttttta aaattgttat taaatattgg ttttttactt aaaaaaaaaa aaaaaaaaaa 3000
aaaa

```

&lt;210&gt; 108

&lt;211&gt; 959

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 108

```

Met Ala Phe Ala Glu Cys Ile Ala Pro Ala Cys Val Met Ser Trp Leu
  1             5             10             15

```

```

Arg Phe Trp Gly Pro Trp Pro Leu Leu Thr Trp Gln Leu Leu Ser Leu
      20             25             30

```

```

Leu Val Lys Glu Ala Gln Pro Leu Val Trp Val Lys Asp Pro Leu Gln
      35             40             45

```

```

Leu Thr Ser Asn Pro Leu Gly Pro Pro Glu Pro Trp Ser Ser Arg Ser
      50             55             60

```

```

Ser His Leu Pro Trp Glu Ser Pro His Ala Pro Ala Pro Pro Ala Ala
      65             70             75             80

```

```

Pro Gly Asp Phe Asp Tyr Leu Gly Pro Ser Ala Ser Ser Gln Met Ser

```

85										90					95				
Ala	Leu	Pro	Gln	Glu	Pro	Thr	Glu	Asn	Leu	Ala	Pro	Phe	Leu	Lys	Glu				
			100					105					110						
Leu	Asp	Ser	Ala	Gly	Glu	Leu	Pro	Leu	Gly	Pro	Glu	Pro	Phe	Leu	Ala				
		115					120					125							
Ala	His	Gln	Asp	Leu	Asn	Asp	Lys	Arg	Thr	Pro	Glu	Glu	Arg	Leu	Pro				
		130				135					140								
Glu	Val	Val	Pro	Leu	Leu	Asn	Arg	Asp	Gln	Asn	Gln	Ala	Leu	Val	Gln				
145				150				155						160					
Leu	Pro	Arg	Leu	Lys	Trp	Val	Gln	Thr	Thr	Asp	Leu	Asp	Arg	Ala	Ala				
			165					170					175						
Gly	His	Gln	Ala	Asp	Glu	Ile	Leu	Val	Pro	Leu	Asp	Ser	Lys	Val	Ser				
		180					185						190						
Arg	Pro	Thr	Lys	Phe	Val	Val	Ser	Pro	Lys	Asn	Leu	Lys	Lys	Asp	Leu				
		195					200					205							
Ala	Glu	Arg	Trp	Ser	Leu	Pro	Glu	Ile	Val	Gly	Ile	Pro	His	Gln	Leu				
	210					215					220								
Ser	Lys	Pro	Gln	Arg	Gln	Lys	Gln	Thr	Leu	Pro	Asp	Asp	Tyr	Leu	Ser				
225				230						235				240					
Met	Asp	Thr	Leu	Tyr	Pro	Gly	Ser	Leu	Pro	Pro	Glu	Leu	Arg	Val	Asn				
			245					250					255						
Ala	Asp	Glu	Pro	Pro	Gly	Pro	Pro	Glu	Gln	Val	Gly	Leu	Ser	Gln	Phe				
		260					265						270						
His	Leu	Glu	Pro	Lys	Ser	Gln	Asn	Pro	Glu	Thr	Leu	Glu	Asp	Ile	Gln				
	275						280					285							
Ser	Ser	Ser	Leu	Gln	Glu	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Gln	Leu	Pro				
	290				295						300								
Gln	Glu	Val	Glu	Pro	Ser	Thr	Gln	Gln	Glu	Ala	Pro	Ala	Leu	Pro	Pro				
305				310					315					320					
Glu	Ser	Ser	Met	Glu	Ser	Leu	Ala	Gln	Thr	Pro	Leu	Asn	His	Glu	Val				
			325					330					335						
Thr	Val	Gln	Pro	Pro	Gly	Glu	Asp	Gln	Ala	His	Tyr	Asn	Leu	Pro	Lys				
		340					345						350						
Phe	Thr	Val	Lys	Pro	Ala	Asp	Val	Glu	Val	Thr	Met	Thr	Ser	Glu	Pro				
		355				360						365							
Lys	Asn	Glu	Thr	Glu	Ser	Thr	Gln	Ala	Gln	Gln	Glu	Ala	Pro	Ile	Gln				
	370					375					380								
Pro	Pro	Glu	Glu	Ala	Glu	Pro	Ser	Ser	Thr	Ala	Leu	Arg	Thr	Thr	Asp				
385				390					395					400					
Pro	Pro	Pro	Glu	His	Pro	Glu	Val	Thr	Leu	Pro	Pro	Ser	Asp	Lys	Gly				

	405		410		415
Gln Ala Gln His Ser His Leu Thr	Glu Ala Thr Val Gln Pro Leu Asp				
420	425	430			
Leu Glu Leu Ser Ile Thr Thr	Glu Pro Thr Thr Glu Val Lys Pro Ser				
435	440	445			
Pro Thr Thr Glu Glu Thr Ser Ala Gln Pro Pro Asp	Pro Gly Leu Ala				
450	455	460			
Ile Thr Pro Glu Pro Thr Thr	Glu Ile Gly His Ser Thr Ala Leu Glu				
465	470	475	480		
Lys Thr Arg Ala Pro His Pro Asp	Gln Val Gln Thr Leu His Arg Ser				
485	490	495			
Leu Thr Glu Val Thr Gly Pro Pro	Thr Lys Leu Glu Ser Ser Gln Asp				
500	505	510			
Ser Leu Val Gln Ser Glu Thr Ala	Pro Glu Glu Gln Lys Ala Ser Thr				
515	520	525			
Ser Thr Asn Ile Cys Glu Leu Cys	Thr Cys Gly Asp Glu Thr Leu Ser				
530	535	540			
Cys Val Gly Leu Ser Pro Lys Gln Arg	Leu Arg Gln Val Pro Val Pro				
545	550	555	560		
Glu Pro Asp Thr Tyr Asn Gly Ile	Phe Thr Thr Leu Asn Phe Gln Gly				
565	570	575			
Asn Tyr Ile Ser Tyr Leu Asp Gly	Asn Val Trp Lys Ala Tyr Ser Trp				
580	585	590			
Thr Glu Lys Leu Ile Leu Ser	Glu Asn Tyr Leu Thr Glu Leu Pro Lys				
595	600	605			
Asp Ser Phe Glu Gly Leu Leu Tyr	Leu Gln Tyr Leu Asp Leu Ser Cys				
610	615	620			
Asn Lys Ile Arg Tyr Ile Glu Arg	Gln Thr Phe Glu Ser Leu Pro Phe				
625	630	635	640		
Leu Gln Tyr Ile Asn Leu Gly Cys	Asn Leu Ile Thr Lys Leu Ser Leu				
645	650	655			
Gly Thr Phe Gln Ala Trp His Gly	Met Gln Phe Leu His Asn Leu Ile				
660	665	670			
Leu Asn Arg Asn Pro Leu Thr	Thr Val Glu Asp Pro Tyr Leu Phe Glu				
675	680	685			
Leu Pro Ala Leu Lys Tyr Leu Asp	Met Gly Thr Thr His Ile Thr Leu				
690	695	700			
Thr Thr Leu Lys Asn Ile Leu Thr	Met Thr Val Glu Leu Glu Lys Leu				
705	710	715	720		
Ile Leu Pro Ser His Met Ala Cys	Cys Leu Cys Gln Phe Lys Asn Ser				



725										730					735				
Ile	Glu	Ala	Val	Cys	Lys	Thr	Val	Lys	Leu	His	Cys	Asn	Thr	Ala	Cys				
			740						745				750						
Leu	Thr	Asn	Ser	Ile	His	Cys	Pro	Glu	Glu	Ala	Ser	Val	Gly	Asn	Pro				
		755					760					765							
Glu	Gly	Ala	Phe	Met	Lys	Met	Leu	Gln	Ala	Arg	Lys	Gln	His	Met	Ser				
		770				775					780								
Thr	Gln	Leu	Thr	Ile	Glu	Ser	Glu	Ala	Pro	Ser	Asp	Ser	Ser	Gly	Ile				
	785				790					795					800				
Asn	Leu	Ser	Gly	Phe	Gly	Gly	Asp	Gln	Leu	Glu	Ile	Gln	Leu	Thr	Glu				
				805					810					815					
Gln	Leu	Arg	Ser	Leu	Ile	Pro	Asn	Glu	Asp	Val	Arg	Lys	Phe	Met	Ser				
			820					825					830						
His	Val	Ile	Arg	Thr	Leu	Lys	Met	Glu	Cys	Ser	Glu	Thr	His	Val	Gln				
		835					840					845							
Gly	Ser	Cys	Ala	Lys	Leu	Met	Ser	Arg	Thr	Gly	Leu	Leu	Met	Lys	Leu				
		850				855					860								
Leu	Ser	Glu	Gln	Gln	Glu	Ala	Lys	Ala	Leu	Asn	Val	Glu	Trp	Asp	Thr				
	865				870					875					880				
Asp	Gln	Gln	Lys	Thr	Asn	Tyr	Ile	Asn	Glu	Asn	Met	Glu	Gln	Asn	Glu				
			885					890					895						
Gln	Lys	Glu	Gln	Lys	Ser	Ser	Glu	Leu	Met	Lys	Glu	Val	Pro	Gly	Asp				
			900					905					910						
Asp	Tyr	Lys	Asn	Lys	Leu	Ile	Phe	Ala	Ile	Ser	Val	Thr	Val	Ile	Leu				
		915					920					925							
Ile	Ile	Leu	Ile	Ile	Ile	Phe	Cys	Leu	Ile	Glu	Val	Asn	Ser	His	Lys				
	930					935					940								
Arg	Ala	Ser	Glu	Lys	Tyr	Lys	Asp	Asn	Pro	Ser	Ile	Ser	Gly	Ala					
	945					950					955								

&lt;210&gt; 109

&lt;211&gt; 1331

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 109

```

gtttcttttct tttccatgat atcattatat ggacagttta ggttggtctc atggattata 60
accatttgga tatttggttc actaacaatt ttcttactgg ccagagttct tgggtggagaa 120
gttgcatatg gccaaagtcct tggagttata ggatattcat tacttctctc cattgtaata 180
gccccgtgtac ttttggtggt tggatcattt gaagtgggtg ctacacttat aaaactgttt 240
ggtgtgtttt gggctgccta cagtgtctgt tcattgttag tgggtgaaga attcaagacc 300
aaaaagcctc ttctgattta tccaatcttt ttattataca tttatTTTTT gtcgttatat 360
actggtgtgt gatccaagtt atacatgaat agaaaaagat ggtgttaaatt ttgtgtgtag 420
gctgggaatt cttgctgaag gaattggaga aaacctgttg ctgcaaaatt ttacatgttc 480
cagatggaaa gggaagtcta agcgcttttt aaaacaattt ttttttgtat ttaattaagc 540

```

```

aattgcagtt atctgggatt tttgggtcag aattttaaat tctgtttgat tctccatatt 600
ccagtgaata aaatacaaaa gcattgtgtt tttaagattg tgcgatatt cacctaaaaa 660
cttgtgccaa aagcacctgg attggtaatt atatttact taaagggtaa atttgacaat 720
atcttgataa tcaaaagtgc aatttttttc ttcaaaatgt tttctccagc atcacagatc 780
ctgcagatat atattttatat ttatacatat atattttatga aataattctt actcacaaaa 840
tatatttctg ataaacatta agatattaaa tctgatgcac aaacttttta atttggccat 900
taatcttttt tatttaaaaa tttaaatttg tttttaaaat tgtatatagt ttttaaaatc 960
tcacacatgc ttcgatactt ccttggttaag aattcttaat aactactaaa actgattttt 1020
aatagttgct gatatatatt tgggttggtt ggggtatactt ttcaaaacca tttttgaatg 1080
tccaaacatc tgattttaaag tttctgttta tctttctgac caaaggagca agagggtataa 1140
tggatatggc attcattaaa atctttacta tgtacaaaaa cagtaatatt tacagcatca 1200
gtaaatattt ttaagtggta cttctaaatc ataaaagttg gggaaagaga cctttaaaaa 1260
cttgtggtgt tgaacaatgt tatatgaagt agaaaaaata aaatacttcc cagttgaaaa 1320
aaaaaaaaa a

```

1331

&lt;210&gt; 110

&lt;211&gt; 118

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

```

Met Ile Ser Leu Tyr Gly Gln Phe Arg Val Val Ser Trp Ile Ile Thr
  1             5             10             15
Ile Trp Ile Phe Gly Ser Leu Thr Ile Phe Leu Leu Ala Arg Val Leu
          20             25             30
Gly Gly Glu Val Ala Tyr Gly Gln Val Leu Gly Val Ile Gly Tyr Ser
          35             40             45
Leu Leu Pro Leu Ile Val Ile Ala Pro Val Leu Leu Val Val Gly Ser
          50             55             60
Phe Glu Val Val Ser Thr Leu Ile Lys Leu Phe Gly Val Phe Trp Ala
          65             70             75             80
Ala Tyr Ser Ala Ala Ser Leu Leu Val Gly Glu Glu Phe Lys Thr Lys
          85             90             95
Lys Pro Leu Leu Ile Tyr Pro Ile Phe Leu Leu Tyr Ile Tyr Phe Leu
          100            105            110
Ser Leu Tyr Thr Gly Val
          115

```

&lt;210&gt; 111

&lt;211&gt; 2610

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 111

```

aattgaccat ctctgtgtaa ttctgattgt gtaccttgag acatacagag gatttctatt 60
tctctttttt gcctaagaat gtataattag tgtcatttta ttggggagaa attttatttt 120
tttgtcattt ctctgaagtt gacatttgat gagtgtattc tgaattctac cactcctctg 180
gggaaaaaaaa tctcaatgga agattgtaca gtttaagtac ttgtttttcc ctcttgaggt 240
atgctatttt taaggtcatt aatttaaaat agaaatattt ctttaaacca ttgggggtaaa 300
attttttaaa atatggtgct ttgatataat ttgagaaaa gtgcctgaat cctaaaacta 360
actgtctata aagttaaagtc cgtataacaa atgattatat atataacaaa gtaaaagtaa 420
atagtgtaga atattaaagt tctcttccag ggcaatgggt attgtggggg attaaacaga 480

```

```

agccccccttg ctatgatcaa ctgcaaagga aagaaactaa atatggttca gtgataaaca 540
aaaaatagcaa agactgatca ggaaaatgag aaagagctat gcaaatacag atgatacaca 600
ggccagagaa ttttaagtgc taatagcttt actcctgcat attccatcag tagaatgagc 660
tttttctttt agtctaagtg acactttgaa ggtcttttgca ttcattacgc ttttatatac 720
tgcttttctt ttctcttctt tttatttcat ccaacacact aagaaatgaa gaagtttatt 780
gatatatgtc ccaataaaaat acaaactctgt tgcaacacaa tgcataatc ttattagcat 840
tcttggttagt tctaaaattc aattcatcca tagacactta ttcccagcct tcaaattgga 900
agctctcttt atgcaagaga aaaagggtatt taaaagtgc caaatttttag aatgaggaat 960
tggttatgct gaatttctgt tccaaaatca ctagggtaaaa ttcccatatc taataatgta 1020
tttttaggaac aagccaaggt ttgtcattaa gacatagtaa cattaaacta atatttaaaa 1080
atagatttag ctatatttta tcatgagaaa aggatttctg tggccaggac aaaaccacc 1140
ttgatcctaa agagacatta gcagtgccca tgtcagagcc ttctcacatt cttggaataa 1200
agtaaaccac tccaaggaag gtgagggcac caaatggatc attcttcaaa atgaaaggca 1260
gtctcttact ttctcttgag acatttcttc cctatttggg gattttatga ctgctgttct 1320
ggagagccca catgttttgc ttccatgcag gccaaatcct tctcttgggt ttgggcaaga 1380
gaagtttaac cacctaacag cattttgatt agacaagaat atcatctgat cttatccagg 1440
gacaagaaag tggcatgacc aaaggttctc cttatgtctc ttcttttaaa tgatttcta 1500
attttcagaa gggctctgga tagactagtt tctgataagg agattttgct gtggtgtctg 1560
ttcttccagg ttaaaggcac acaaagcctt ctgtggtcca cttgtgcttg tcttctgaat 1620
tactgaaac atggcagaat aaggtttaag gaaaatgaat ttgacttta catagttaaa 1680
tgagtaggag ttacagcaaa aaaaaaaaaa aagacctaaa acttttattt aatagtattc 1740
ttcacctcag gaaattcaga ctttagttg ttccaggtcag aaaatgtgga tatatcctgt 1800
aacactgtct tcacctcact caagttaaatt gtaacagcaa atactttcag ctttactttc 1860
aaaagggttc cagaactttc cacatttcat cactttcact tctacccttc tggcaaaaat 1920
taccaagatt tttagtagtt tactgttccc tggtttctct gatttgatcc ttgctaccat 1980
tctgtttagt cctcaaagaa aaaaaatcaa cattttaaaa cgtttcaatt cttactaatg 2040
gttctcatct cagaagaaaa aacaacgaaa tatcttatgt taatctaaaa aaccttcagt 2100
gacctacttg atctcatttt ctaccatttt cctcctcttt ttctgaaata catcaacaca 2160
gagcactttt cctctccttt aatgcacaaa gatggcagga cttttgaatg ttacatttat 2220
ttatcttctt ctagagtgc ttcccttata caccatgtg acttggtcct ccttctcttc 2280
tagtctttgt ttatatatat attattatca cagagggcta ggaaagaaaa caccactgc 2340
tgcgcccccac actcatccac ctgccctgta ccacttactt tgttttggtt ttctctgtag 2400
aattcatgac tttttgaaat ataatttttt taatgtgtac atactttatg ctttctctca 2460
ttcatatgta aagtctggaa gacacagact ggtttttttg ttactgttg atgcttcagt 2520
ccctaaaata tgcatagcat gaattgccac tttttaaatt aataaatctg gaacattgtt 2580
aaaattcaaa aaaaaaaaaa aaaaaaaaaa
2610

```

&lt;210&gt; 112

&lt;211&gt; 116

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 112

```

Met Ala Gly Leu Leu Asn Val Thr Phe Ile Tyr Leu Leu Leu Glu Cys
 1             5             10             15

Leu Ser Leu Tyr Thr His Val Thr Cys Ser Ser Leu Pro Ser Ser Leu
      20             25             30

Cys Leu Tyr Ile Tyr Tyr Tyr His Arg Gly Leu Gly Lys Lys Thr Pro
 35             40             45

Thr Ala Ala Pro His Thr His Pro Pro Ala Leu Tyr His Leu Leu Cys
 50             55             60

Phe Val Phe Leu Cys Arg Ile His Asp Phe Leu Lys Tyr Asn Phe Phe
 65             70             75             80

Asn Val Tyr Ile Leu Tyr Ala Phe Ser His Ser Tyr Val Lys Ser Gly
      85             90             95

```

Arg His Arg Leu Val Phe Leu Phe Thr Val Asp Ala Ser Val Pro Lys  
 100 105 110

Ile Cys Ile Ala  
 115

<210> 113  
 <211> 2759  
 <212> DNA  
 <213> Homo sapiens

<400> 113  
 tttttttttt tttgaaagac acacgttatt ttattaatat agccatctct cccactgcc 60  
 ccagtgggtga aggtgtttgc attgcaacat ggagggggcac caaatgctct gcgggcccta 120  
 gcccgtgcc acaggctagg cctgcctgca gccaagaagg ctgctcaaac tctagatgcc 180  
 atttgagggc atgaggacct gagcccagag gtggcagtggt cctaccagc gaagtcaaca 240  
 gatcgtgctc caggtcccag ctctgggctg ggccaggact aaatcctggc tcccctttct 300  
 tgggtactaag gggattagtg cttggtttgc tgtaggggggt cagagtaggg agggttccag 360  
 gaagggttcc agagtgggct cacaggggac ctctccctt ggctcttgg agtccaggtc 420  
 gtcgagggcg caaagctgca cgccatcctg ggcaagctgg gcccgagcg tgggcgcggt 480  
 gaggacgcgc agtcatgca gccgtccca agagcaagag aaagcgtcgg ggcttccacc 540  
 gcagccgcgc gtgggaggca cactggggtg gccggggtgc gccatcagct cggctgtcag 600  
 ggtgtggccc gctagggtac cttccaggac ccgcgcagg gccccggaca cgcggtgagc 660  
 ggacatgtgc cggccgcaag tgctcaggcc cacgaaggcg tctgtccacc gcaggccgtg 720  
 gcgggagaag gggccacggc ggcccggcg tgcgctcca cggcgaggc gaaggcacgc 780  
 gcgggggcct ccagccaagt gcagccacc acaccgcgt ccagcggcag tcgcgtaaag 840  
 cgcaccccat aggcctgcag cgcctcggcg aacacctggc acacgcctgg gagcacgtgc 900  
 acgtgctggg gccgctccgc gtgctgggg gccctgccca gcagctcccg gaagcagctt 960  
 agttgggcct cgagctcctc ccgcacctag agggcgagcg agagacacct tgagcgaccg 1020  
 ggagtagctg ccgaggatac cctcctcgtg cttccgtgtg atggctccag tggttgaaatg 1080  
 cggaagtcac ccaccgccag ctcttaacgg cctcacagta cctccgggc ggagctctgg 1140  
 gggctcctgc gagcatcctc ctgtagctgc ggctccgcac ctgaggcaaa tccacgtctc 1200  
 cggccgccac cgcctcccgg aatcccatct tgccaaggaa gaagccttcc gggccgagca 1260  
 gcgatgaggc gccacggcgg gccggaccca cggggcgccc ctcgacagc ttggcgtgga 1320  
 ggcccggtgg gatgctgtgc ctgcgggcca gctccgccgc gctctccgtg gccgcaccgt 1380  
 tgaccagcag ggacacgctg gtcacggccc cggccagaaa ggctccacg ataccctcat 1440  
 cgcgtcgcgg gcagtaacca aagtctccg cggtgaccac caggcgcccg cgggagccgg 1500  
 cagatccggg tcttgtggct aagagtgcg tggctactcg aatcaaaaca gaggaggggg 1560  
 aggaagccgg cggccagaaa cggcagtggc agcagcgtcc ggagcagccg cagcctctctg 1620  
 gaagctccag gcggtctttc tgccgagcct cggctccggc ccccatcctc cccgccccat 1680  
 cggttgttgt ctgggcggat ttaaacagtc aagtaaaatc aagctgggta atcatggcag 1740  
 aaggtggatt tgatccctgt gaatgtgttt gctctcatga acatgcaatg agaagactga 1800  
 tcaatctggt acggcagctc cagtcctact gcacagacac agagtgtctt caggaattac 1860  
 cgggaccctc tgggtgataat ggcacagtg ttacaatgat cttggtagcc tggatgggta 1920  
 ttgcattgat cttgttctta ctgagacctc ctaatctaag aggatccagc ctacctggaa 1980  
 agccaaccag tctcataat ggacaagatc caccagctcc tcctgtggac taactttgtg 2040  
 atatgggaag tgaaaatagt taacaccttg caccagcaaa cgaacgaaga tgaccagagt 2100  
 actcttaacc ccattagaac tggttttctt tttgtatctg caatatggga tggattgttt 2160  
 tcatgagct tctagaaatt tcaactgcaa gtttattttt gcttccgtgt ttactgccat 2220  
 tctatcttac agtatatttg agtgaatgat tatattttta aaaagttaca tggggctttt 2280  
 ttggttgtcc taaacttaca aacattccac tcattctgtt tgtaactgtg attataattt 2340  
 ttgtgataat ttctggcctg attgaaggaa atttgagagg tctgcattta tatattttta 2400  
 atagatttga taggttttta aattgctttt tttcataagg tatttataaa gttatttggg 2460  
 gttgtctggg attgtgtgaa agaaaattag aaccacgctg tattttacatt taccttggta 2520  
 gtttatttgt ggtggcagc tttctgtagt tttggggact gtggtagctc ttggattgtt 2580  
 ttgcaaatga cagctgaaat ctgtgtcatg gattaaactg gcttatgtgg ctagaatagg 2640  
 aagagaaaaa aaatgaaatg gttgtttact aattttatac tcccattaaa aatctcta 2700  
 gtaagaaaaa ccttaataa acatgattga tcaatatgaa aaaaaaaaaa aaaaaaaaaa 2759

<210> 114  
 <211> 99  
 <212> PRT  
 <213> Homo sapiens

<400> 114  
 Met Ala Glu Gly Gly Phe Asp Pro Cys Glu Cys Val Cys Ser His Glu  
           1                  5                  10                  15  
 His Ala Met Arg Arg Leu Ile Asn Leu Leu Arg Gln Ser Gln Ser Tyr  
                   20                  25                  30  
 Cys Thr Asp Thr Glu Cys Leu Gln Glu Leu Pro Gly Pro Ser Gly Asp  
                   35                  40                  45  
 Asn Gly Ile Ser Val Thr Met Ile Leu Val Ala Trp Met Val Ile Ala  
           50                  55                  60  
 Leu Ile Leu Phe Leu Leu Arg Pro Pro Asn Leu Arg Gly Ser Ser Leu  
           65                  70                  75                  80  
 Pro Gly Lys Pro Thr Ser Pro His Asn Gly Gln Asp Pro Pro Ala Pro  
                   85                  90                  95  
 Pro Val Asp

<210> 115  
 <211> 1404  
 <212> DNA  
 <213> Homo sapiens

<400> 115  
 aatcgggacg ggacgaatta ttggttgggg gaaacccacg aggggacgcg gccgaggagg 60  
 gtcgctgtcc acccgggggc gtgggagtga ggtaccagat tcagcccat tggcccccac 120  
 gcctctgttc tcggaatccg ggtgctgcgg attgaggtcc cggttcctaa cgggtgggac 180  
 ggtgtcctcg ggatgagatt tggcgcttcc tcggggcttt ggtgggacg gtgtcctcag 240  
 gatgagattt agggtttcc cggggcttcc gggatcttca cctaataacc ggtattattt 300  
 tatgagagga gtggtcttgg ctgtcagaac tggatccctg gggatgattt tgggaattag 360  
 tggagtgatc tctgaagacc tagggctatg atctggagct gctgtggctg aaatttgggg 420  
 cctctgaagt ggcattggaga ttgaggtcca gagagcctga gatcttgagg gctgacattt 480  
 ggagagatgg ggtcgagggt tgtctttggg ccttgactgc tttgggcctt tctcactctc 540  
 attcccggga tgctttgcca gaatctctgc tggattggcc gtaacctgt ccccgagcgg 600  
 gctcacagg tctgaaggcc acgcatgagg caaaggtaaa gttctgagcc acccggtgcc 660  
 tccttccag gactgcaaga tggaggaagg cgggaacctt ggaggcctga ttaagatggt 720  
 ccatctactg gtcttgtcag gtgcctgggg catgcaaatg tgggtgacct tcgtctcagg 780  
 tagggaccct cagcttggat gtcattgggt cctgggggtg ggatggaaat aagaggggaa 840  
 ccgggaagtg ccctaacacc cctgtggtcc ccataccctg caggcttcc gcttttccga 900  
 agccttcccc gacatacctt cggactagtg cagagcaaac tcttccccct ctacttccac 960  
 atctccatgg gctgtgcctt catcaacctc tgcattcttg cttcacagca tgcttgggct 1020  
 cagctcacat tctgggaggg cagccagctt tactgtctgt tcctgagcct tacgtgggcc 1080  
 actgtcaacg cccgctgggt ggaaccccg accacagctg ccatgtgggc cctgcaaac 1140  
 gtggagaagg agcgaggcct ggggtggggag gtaccaggca gccaccaggg tcccgatccc 1200  
 taccgccagc tgcgagagaa ggaccccaag tacagtgtc tccgccagaa tttcttccgc 1260  
 taccatgggc tgtcctctct ttgcaatctg ggctgcgtcc tgagcaatgg gctctgtctc 1320  
 gctggccttg ccctggaaat aaggagcctc tagcatgggc cctgcatgct aataaatgct 1380  
 tcttcagaaa aaaaaaaaaa aaaa 1404

<210> 116  
 <211> 184  
 <212> PRT  
 <213> Homo sapiens

<400> 116

```

Met Ser Trp Val Pro Gly Val Gly Met Glu Ile Arg Gly Glu Pro Gly
 1              5              10              15

Ser Ala Leu Thr Pro Leu Trp Ser Pro Tyr Pro Ala Gly Phe Leu Leu
      20              25              30

Phe Arg Ser Leu Pro Arg His Thr Phe Gly Leu Val Gln Ser Lys Leu
      35              40              45

Phe Pro Phe Tyr Phe His Ile Ser Met Gly Cys Ala Phe Ile Asn Leu
      50              55              60

Cys Ile Leu Ala Ser Gln His Ala Trp Ala Gln Leu Thr Phe Trp Glu
      65              70              75              80

Ala Ser Gln Leu Tyr Leu Leu Phe Leu Ser Leu Thr Leu Ala Thr Val
      85              90              95

Asn Ala Arg Trp Leu Glu Pro Arg Thr Thr Ala Ala Met Trp Ala Leu
      100             105             110

Gln Thr Val Glu Lys Glu Arg Gly Leu Gly Gly Glu Val Pro Gly Ser
      115             120             125

His Gln Gly Pro Asp Pro Tyr Arg Gln Leu Arg Glu Lys Asp Pro Lys
      130             135             140

Tyr Ser Ala Leu Arg Gln Asn Phe Phe Arg Tyr His Gly Leu Ser Ser
      145             150             155             160

Leu Cys Asn Leu Gly Cys Val Leu Ser Asn Gly Leu Cys Leu Ala Gly
      165             170             175

Leu Ala Leu Glu Ile Arg Ser Leu
      180

```

<210> 117  
 <211> 1801  
 <212> DNA  
 <213> Homo sapiens

<400> 117

```

tgaagaagggt gtttactttt tttgaaatta ccttgagaca tttcaaactg tgcagaagat 60
atatgcacaa aagcaaagt cttgcagttt gctatagcca cttatacatc atctggctct 120
tgaatagctt taattcagct gttgaatctc acttgaattt gagcaaaacc ttcattcttta 180
tatgtatctg gacaaattac ttcaattgct tgacagtaat gaccaatcaa tttatttaaa 240
atagtatcat ttagtaggac agtggttttc tctggtttga gcaacgaatt caaccagtcc 300
tctgggttga tcatcatcat catcatcatt tggttatcag ttcttgagtt atttttacca 360
gggagtttta tacctttaga caactatttt gaattatctc aggaatgtca tatatctctg 420
cctctttaga gtcagtcact ggcactttgt ctgtttggtg acatcatggt tccctgactg 480
ttcttcatct ttgtagttat acattgatat ctgtgcattg aatatgtagg tatttataaa 540
cagtctttgc aatctggctt tgtctgtgat tgtccttgta tagtaggtct gtccagaaat 600
tgtaagcata ctgtcttttt tggctcttaa gcccgtaac gctacagccc gtgtagtgcc 660

```

```

aaatggtgcc ctaagcccag gttccctgca gtccactctg tgatttggtt gttgactgct 720
gtgagcccca ccccatctct ttgtttctaa tttacaccta gcaggctaac cctgctggca 780
cctgcagtgcc ttccaggggg aaaggaccat agtgtgcccc tgtgaagagt ctcagaatgg 840
tgcggaaggt gaatgcccac ctcccgctct cttttcccac tgtagaaact gattccagag 900
aaattctcca agtgcggtgc tatgtgggct tgcgggagag gtgttatgat caaacagaac 960
cattctcttt accctctggt catggttttt cttggctctg tgggtccagt agttgtcaca 1020
gcttactccc caatttctgg gatattcagg gcaataatct tgccactggg tatttgctag 1080
ttgaaattat gtggtaggga gagaagccag taagcttcac ttctccggtt ggctgatgtc 1140
actctcgaat tctgtacttt catacggatt gtaataggga gggctctaaa ggaagcttca 1200
gttttggatt tttagagctt cttctaagta caattgttga atcaagagg aaagatggta 1260
tgttataact ggaaatacgc tgagattaaa aaggagataa attagcccca ttgggacagt 1320
gctattggga atgtgaattg ataccacttt cctggagtct agtttagtat ttattttata 1380
tgtaatgcct gaaaagatgt gtgtctcctt tgactcaata cttataggaa tttacacaaa 1440
ggttataatc aaagattata gaactgttat actggaaaaa cacgaaatat tctaattatg 1500
aatatttggt ggattgggta aataaactat ggtttgatat gctctaaaaa taatgttaca 1560
gaaaaaagtg tactgatatg gcaaaatgta tgacttatag ttaaaaaagc aggttagatg 1620
ttgatagata cagtatgata gaaaaagatc aggaaggat atgctgacat ttaaatctgg 1680
atatttatga gtgttttttt ttttcaatc tttgtacat catgtatttt ctagaaattg 1740
tattactatc tttgtaataa agtaaattat ttttaaggga ctaaaaaaaa aaaaaaaaaa 1800
a                                                                 1801

```

&lt;210&gt; 118

&lt;211&gt; 86

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 118

```

Met Val Arg Lys Val Asn Ala His Leu Pro Leu Ser Phe Pro Thr Val
  1              5              10              15

Glu Thr Asp Ser Arg Glu Ile Leu Gln Val Arg Cys Tyr Val Gly Leu
          20              25              30

Arg Glu Arg Cys Tyr Asp Gln Thr Glu Pro Phe Ser Leu Pro Ser Val
          35              40              45

His Gly Phe Ser Trp Leu Cys Gly Pro Val Ser Cys His Ser Phe Thr
          50              55              60

Pro Asn Phe Trp Asp Ile Gln Gly Asn Asn Leu Ala Thr Gly Tyr Leu
          65              70              75              80

Leu Val Glu Ile Met Trp
                  85

```

&lt;210&gt; 119

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (2)

&lt;223&gt; biotinylated phosphoramidite residue

&lt;400&gt; 119

ancgggagcc tcttgaccat ctcctcttc

29

<210> 120

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 120

cncacagaaa attcaataag accctcgct

29

<210> 121

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 121

cncagctctt cgtagggaag ttctgactt

29

<210> 122

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 122

antcctgcac accagccagt aacgccacc

29

<210> 123

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature



<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 123  
gnggctggaa agatgtgtgg ggatcaaga 29

<210> 124  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 124  
anatgggtct aagccacaca acagggtga 29

<210> 125  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 125  
ancggcaggg aacttacagg gacagagct 29

<210> 126  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 126  
nggttttcgg tgatgatggt gtagaggat 29

<210> 127  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 127

cnagaacaca tagggatgcg agagcaagc

29

<210> 128

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 128

anactgaaaa ctgagtatgt gcgagtgtgta

29

<210> 129

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 129

gntatccatt tcctcttctt catctgagt

29

<210> 130

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 130

tncttgaggc aatgggtgaa gtccggcg

29

<210> 131

<211> 29

<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 131  
tntctgtctg tgccttctc tatccgaac

29

<210> 132  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 132  
gngcatctca ctggatgtca tcatcatca

29

<210> 133  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 133  
tngtccatgt gaagggcatg ggccagttg

29

<210> 134  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 134

gngcactgta ttgagctgat tgctgaagc

29

<210> 135

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 135

cncagaagca gaagaatgac aggcaacac

29

<210> 136

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 136

anacattctg agtagttgca tgatttccc

29

<210> 137

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 137

gnccagaaaag ttgaggacat gctgggcag

29

<210> 138

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 138  
angggaacaa gacaactgga gaaggggtca 29

<210> 139  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 139  
tngtctccca ggtagacaga gggcttcag 29

<210> 140  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 140  
anccatctac atgtgcattg acaagctta 29

<210> 141  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 141  
tngtgataga tccttttcgta acaccaagt 29

<210> 142  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 142

angaccagat ctcacccagc acatcaaac

29

<210> 143

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 143

tntttggggc aagatggctg ttaagcagt

29

<210> 144

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 144

tnggttggtc cgggcagggc attcttgtc

29

<210> 145

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 145

tnacacgctc tgtgctagga cttttatat

29

<210> 146

<211> 29

<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 146  
cnggatgtgt gatattggag cttgctggt

29

<210> 147  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 147  
ancatcaaag gtgcccaaat aagttccca

29

<210> 148  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 148  
anatcactgc atttggtctg gaacctgac

29

<210> 149  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 149

gntgacttca atctcctcac cttccaccg

29

<210> 150

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 150

gnagtgccac ctatgactac caaattctc

29

<210> 151

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 151

gngggatgag gcaatgaaca caatgaaag

29

<210> 152

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 152

cnaaactggt gtttttaccg tatccttca

29

<210> 153

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature



<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 153  
tngattctgc cgaatccgaa agtgccttc 29

<210> 154  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 154  
angttgatgg gctcaacaca gggcagagg 29

<210> 155  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 155  
anggatgcca tctctcacc actctgtac 29

<210> 156  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 156  
anaccaccac ctcgacaggc attccttaa 29

<210> 157  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 157  
cncctgaggg tagaaggccg ctcaggttt 29

<210> 158  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 158  
tngagttagc agagcaagaa gcaaggagg 29

<210> 159  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 159  
tttgcatgta tagttctctg cagtagcat 29

<210> 160  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 160  
anctgccatg tcaaagagga gccagatga 29

<210> 161  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 161  
tnaggcttgt gcacttgctg agcttccag 29

<210> 162  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 162  
cnttcgagca ctaagaacgg gacacgta 29

<210> 163  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 163  
angagaagtt ctgtgcgtgg gtctggtcg 29

<210> 164  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 164  
angtcgttct gcaccaggcc tctgtagcg 29

<210> 165  
<211> 29  
<212> DNA

<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 165  
gntcatctcc agcaccatct ccatcaatg 29

<210> 166  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<400> 166  
actgtctttg aatggtattg 20

<210> 167  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 167  
angttcacat atgatacaag gctctttcc 29

<210> 168  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 168  
cnagtagaca gcacaggtag tcggcttga 29

<210> 169  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 169  
gngaatgcaa tatgaagaaa acaggtcag 29

<210> 170  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 170  
anatcaaggt gattaggctc ttccatgca 29

<210> 171  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 171  
tntacattca atgcctttgc ttctgctg 29

<210> 172  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 172  
antccatgag accaccctaa actgtccat 29

<210> 173  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 173  
gngaggaaaa gtgctctgtg ttgatgtat 29

<210> 174  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 174  
antccacctt ctgccatgat taccagct 29

<210> 175  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 175  
anccaagatg cagaggttga tgaaggcac 29

<210> 176  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 176  
tntaatgcct gaaaagatgt gtgtctcct

29

<210> 177  
<211> 388  
<212> PRT  
<213> Homo sapiens

<400> 177  
Met His Leu Tyr Lys Thr Asn Lys Met Thr Ser Leu Lys Glu Asp Val  
1 5 10 15  
Arg Arg Ser Ala Met Leu Cys Ile Leu Thr Val Pro Ala Ala Met Thr  
20 25 30  
Ser His Asp Leu Met Lys Phe Val Ala Pro Phe Asn Glu Val Ile Glu  
35 40 45  
Gln Met Lys Ile Ile Arg Asp Ser Thr Pro Asn Gln Tyr Met Val Leu  
50 55 60  
Ile Lys Phe Arg Ala Gln Ala Asp Ala Asp Ser Phe Tyr Met Thr Cys  
65 70 75 80  
Asn Gly Arg Gln Phe Asn Ser Ile Glu Asp Asp Val Cys Gln Leu Val  
85 90 95  
Tyr Val Glu Arg Ala Glu Val Leu Lys Ser Glu Asp Gly Ala Ser Leu  
100 105 110  
Pro Val Met Asp Leu Thr Glu Leu Pro Lys Cys Thr Val Cys Leu Glu  
115 120 125  
Arg Met Asp Glu Ser Val Asn Gly Ile Leu Thr Thr Leu Cys Asn His  
130 135 140  
Ser Phe His Ser Gln Cys Leu Gln Arg Trp Asp Asp Thr Thr Cys Pro  
145 150 155 160  
Val Cys Arg Tyr Cys Gln Thr Pro Glu Pro Val Glu Glu Asn Lys Cys  
165 170 175  
Phe Glu Cys Gly Val Gln Glu Asn Leu Trp Ile Cys Leu Ile Cys Gly  
180 185 190  
His Ile Gly Cys Gly Arg Tyr Val Ser Arg His Ala Tyr Lys His Phe  
195 200 205  
Glu Glu Thr Gln His Thr Tyr Ala Met Gln Leu Thr Asn His Arg Val  
210 215 220  
Trp Asp Tyr Ala Gly Asp Asn Tyr Val His Arg Leu Val Ala Ser Lys  
225 230 235 240  
Thr Asp Gly Lys Ile Val Gln Tyr Glu Cys Glu Gly Asp Thr Cys Gln  
245 250 255  
Glu Glu Lys Ile Asp Ala Leu Gln Leu Glu Tyr Ser Tyr Leu Leu Thr  
260 265 270

Ser Gln Leu Glu Ser Gln Arg Ile Tyr Trp Glu Asn Lys Ile Val Arg  
 275 280 285

Ile Glu Lys Asp Thr Ala Glu Glu Ile Asn Asn Met Lys Thr Lys Phe  
 290 295 300

Lys Glu Thr Ile Glu Lys Cys Asp Asn Leu Glu His Lys Leu Asn Asp  
 305 310 315 320

Leu Leu Lys Glu Lys Gln Ser Val Glu Arg Lys Cys Thr Gln Leu Asn  
 325 330 335

Thr Lys Val Ala Lys Leu Thr Asn Glu Leu Lys Glu Glu Gln Glu Met  
 340 345 350

Asn Lys Cys Leu Arg Ala Asn Gln Val Leu Leu Gln Asn Lys Leu Lys  
 355 360 365

Glu Glu Glu Arg Val Leu Lys Glu Thr Cys Asp Gln Lys Asp Leu Gln  
 370 375 380

Ile Thr Glu Ile  
 385

&lt;210&gt; 178

&lt;211&gt; 171

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 178

Met Met Met Gln Cys Val Ser Arg Met Leu Ala His Pro Leu His Val  
 1 5 10 15

Ile Ser Met Arg Cys Met Val Gln Phe Val Gly Arg Glu Ala Lys Tyr  
 20 25 30

Ser Gly Val Leu Ser Ser Ile Gly Lys Ile Phe Lys Glu Glu Gly Leu  
 35 40 45

Leu Gly Phe Phe Val Gly Leu Ile Pro His Leu Leu Gly Asp Val Val  
 50 55 60

Phe Leu Trp Gly Cys Asn Leu Leu Ala His Phe Ile Asn Ala Tyr Leu  
 65 70 75 80

Val Asp Asp Ser Phe Ser Gln Ala Leu Ala Ile Arg Ser Tyr Thr Lys  
 85 90 95

Phe Val Met Gly Ile Ala Val Ser Met Leu Thr Tyr Pro Phe Leu Leu  
 100 105 110

Val Gly Asp Leu Met Ala Val Asn Asn Cys Gly Leu Gln Ala Gly Leu  
 115 120 125

Pro Pro Tyr Ser Pro Val Phe Lys Ser Trp Ile His Cys Trp Lys Tyr  
 130 135 140

Leu Ser Val Gln Gly Gln Leu Phe Arg Gly Ser Ser Leu Leu Phe Arg



```

<400> 180
Met Gly Pro Val Ser Ala Gly Ser Gln Gly Cys Gly Thr Cys Ala Val
  1          5          10          15
Lys Leu Ala Pro Thr Trp Arg Ala Ala Ala Ala Thr Cys Phe Leu Gln
          20          25          30
His Leu Leu Pro Cys Ser Val Ser Ser Leu Ser Pro Arg Leu Ala Gln
          35          40          45
Glu Cys Trp Lys Ser Ser Arg Leu Gly Leu Gly Ala Trp Pro Leu Asp
          50          55          60
Ile Pro Arg Ala Ser Pro Val Leu Pro Ser Pro Arg Thr Thr Gly Pro
  65          70          75          80

```

**WO 99/57132**

**PCT/US99/09970**

Leu Ala

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :C07H 21/04; C07K 14/705; C12N 15/09, 15/63; C12Q 1/68  
US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 24.3; 435/7.2, 69.1; 530/350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

GENEMBL,N-GENSEQ-34,EST,A-GENSEQ35,PIR60,SWISS-PROT37,SPTREMBL19,APS,MEDLINE,CAPLUS,WPIDS,  
JAPIO, SCISEARCH  
Search terms: secreted protein, bn36553,gage 1, gage 6, rgd

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	Database EST, Accession Number AA242967, HILLIER, L. et al. WashU_Merck EST Project 1997, 06 August 1997.	1-3 ----- 4-7, 9, 11
X - Y	Database EST, Accession Number AA524997, NCI-CGAP <a href="http://www.nci.nlm.nih.gov/ncicgap">http://www.nci.nlm.nih.gov/ncicgap</a> , National cancer Institute, cancer Genome AnDatomy Project (CGAP), Tumor Gene Index. 05 August 1997.	1-3 ----- 4-7, 9, 11
X,P --- Y,P	Database Sptrembl19, Accession Number 060829, STROM, T. M et al. JM27 Protein, Complete CDs (Clone Image 145745 and Image 257878), 01 August 1998.	1-3 ----- 4-7, 9, 11
X,P --- Y,P	Database GenEmbl, Accession Number HSA005894, AJ005894, STROM, T. M. et al. Transcription map in Hp11.23, 15 May 1998.	1-3 ----- 4-7, 9, 11

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* "X" "Y" "&"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
--------------------------------------	---	--------------------------	--

Date of the actual completion of the international search

22 JULY 1999

Date of mailing of the international search report

09 SEP 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NIKMAI S. BASI

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ROBINSON, A.K. et al. A Gene from the Hyperthermophile <i>Pyrococcus furiosus</i> whose Deduced Product is Homologous to Members of the Prolyl Oligopeptidase Family of Proteases. <i>Gene</i> . 1995, Vol. 152, pages 103-106, see Fig. 1, amino acids 358-365.	9, 11
Y	BERGER, S.L. et al. Guide to Molecular Cloning Techniques. <i>Methods in Enzymology</i> . 1987, Vol. 152, pages 661-673, see entire document.	1-7, 9, 11

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-11

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

A. CLASSIFICATION OF SUBJECT MATTER:  
US CL :

536/23.1, 24.3; 435/7.2, 69.1; 530/350

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s)1-11, drawn to polynucleotide comprising SEQ ID NO:1, fragments thereof, expression vector containing said sequence, cell transformed with said vector, polypeptide of SEQ ID NO:2, fragments of the polypeptide of SEQ ID NO:2, process for preparing said polypeptide.

Group II, claim(s)12-13, drawn to polynucleotides comprising SEQ ID NO:3, fragments thereof, polypeptide of SEQ ID NO:4, fragments of the polypeptide of SEQ ID NO:4.

Group III, claim(s)14-15, drawn to polynucleotides comprising SEQ ID NO:5, fragments thereof, polypeptide of SEQ ID NO:6, fragments of the polypeptide of SEQ ID NO:6.

Group IV, claim(s)16-17, drawn to polynucleotides comprising SEQ ID NO:7, fragments thereof, polypeptide of SEQ ID NO:8, fragments of the polypeptide of SEQ ID NO:8.

Group V, claim(s)18-19, drawn to polynucleotides comprising SEQ ID NO:9, fragments thereof, polypeptide of SEQ ID NO:10, fragments of the polypeptide of SEQ ID NO:10.

Group VI, claim(s)20-21, drawn to polynucleotides comprising SEQ ID NO:11, fragments thereof, polypeptide of SEQ ID NO:12, fragments of the polypeptide of SEQ ID NO:12.

Group VII, claim(s)22-23, drawn to polynucleotides comprising SEQ ID NO:13, fragments thereof, polypeptide of SEQ ID NO:14, fragments of the polypeptide of SEQ ID NO:14.

Group VIII, claim(s)24-25, drawn to polynucleotides comprising SEQ ID NO:15, fragments thereof, polypeptide of SEQ ID NO:16, fragments of the polypeptide of SEQ ID NO:16.

Group IX, claim(s)26-27, drawn to polynucleotides comprising SEQ ID NO:17, fragments thereof, polypeptide of SEQ ID NO:18, fragments of the polypeptide of SEQ ID NO:18.

Group X, claim(s)28-29, drawn to polynucleotides comprising SEQ ID NO:19, fragments thereof, polypeptide of SEQ ID NO:20, fragments of the polypeptide of SEQ ID NO:20.

Group XI, claim(s)30-31, drawn to polynucleotides comprising SEQ ID NO:21, fragments thereof, polypeptide of SEQ ID NO:22, fragments of the polypeptide of SEQ ID NO:22.

Group XII, claim(s)32-33, drawn to polynucleotides comprising SEQ ID NO:23, fragments thereof, polypeptide of SEQ ID NO:24, fragments of the polypeptide of SEQ ID NO:24.

Group XIII, claim(s)34-35, drawn to polynucleotides comprising SEQ ID NO:25, fragments thereof, polypeptide of SEQ ID NO:26, fragments of the polypeptide of SEQ ID NO:26.

Group XIV, claim(s)36-37, drawn to polynucleotides comprising SEQ ID NO:27, fragments thereof, polypeptide of SEQ ID NO:28, fragments of the polypeptide of SEQ ID NO:28.

Group XV, claim(s)38-39, drawn to polynucleotides comprising SEQ ID NO:29, fragments thereof, polypeptide of SEQ ID NO:30, fragments of the polypeptide of SEQ ID NO:30.

Group XVI, claim(s)40-41, drawn to polynucleotides comprising SEQ ID NO:31, fragments thereof, polypeptide of SEQ ID NO:32, fragments of the polypeptide of SEQ ID NO:32.

Group XVII, claim(s)42-43, drawn to polynucleotides comprising SEQ ID NO:33, fragments thereof, polypeptide of SEQ ID NO:34, fragments of the polypeptide of SEQ ID NO:34.

Group XVIII, claim(s)44-45, drawn to polynucleotides comprising SEQ ID NO:35, fragments thereof, polypeptide of SEQ ID NO:36, fragments of the polypeptide of SEQ ID NO:36.

Group XIX, claim(s)46-47, drawn to polynucleotides comprising SEQ ID NO:37, fragments thereof, polypeptide of SEQ ID NO:38, fragments of the polypeptide of SEQ ID NO:38.

Group XX, claim(s)48-49, drawn to polynucleotides comprising SEQ ID NO:39, fragments thereof, polypeptide of SEQ ID NO:40, fragments of the polypeptide of SEQ ID NO:40.

Group XXI, claim(s)50-51, drawn to polynucleotides comprising SEQ ID NO:41, fragments thereof, polypeptide of SEQ ID NO:42, fragments of the polypeptide of SEQ ID NO:42.

Group XXII, claim(s)52-53, drawn to polynucleotides comprising SEQ ID NO:43, fragments thereof, polypeptide of SEQ ID NO:44, fragments of the polypeptide of SEQ ID NO:44.

Group XXIII, claim(s)54-55, drawn to polynucleotides comprising SEQ ID NO:45, fragments thereof, polypeptide of SEQ ID NO:46, fragments of the polypeptide of SEQ ID NO:46.

Group XXIV, claim(s)56-57, drawn to polynucleotides comprising SEQ ID NO:47, fragments thereof,

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

polypeptide of SEQ ID NO:48, fragments of the polypeptide of SEQ ID NO:48.

Group XXV, claim(s)58-59, drawn to polynucleotides comprising SEQ ID NO:49, fragments thereof, polypeptide of SEQ ID NO:50, fragments of the polypeptide of SEQ ID NO:50.

Group XXVI, claim(s)60-61, drawn to polynucleotides comprising SEQ ID NO:51, fragments thereof, polypeptide of SEQ ID NO:52, fragments of the polypeptide of SEQ ID NO:52.

Group XXVII, claim(s)62-63, drawn to polynucleotides comprising SEQ ID NO:53, fragments thereof, polypeptide of SEQ ID NO:54, fragments of the polypeptide of SEQ ID NO:54.

Group XXVIII, claim(s)64-65, drawn to polynucleotides comprising SEQ ID NO:55, fragments thereof, polypeptide of SEQ ID NO:56, fragments of the polypeptide of SEQ ID NO:56.

Group XXIX, claim(s)66-67, drawn to polynucleotides comprising SEQ ID NO:57, fragments thereof, polypeptide of SEQ ID NO:58, fragments of the polypeptide of SEQ ID NO:58.

Group XXX, claim(s)68-69, drawn to polynucleotides comprising SEQ ID NO:59, fragments thereof, polypeptide of SEQ ID NO:60, fragments of the polypeptide of SEQ ID NO:60.

Group XXXI, claim(s)70-71, drawn to polynucleotides comprising SEQ ID NO:61, fragments thereof, polypeptide of SEQ ID NO:62, fragments of the polypeptide of SEQ ID NO:62.

Group XXXII, claim(s)72-73, drawn to polynucleotides comprising SEQ ID NO:63, fragments thereof, polypeptide of SEQ ID NO:64, fragments of the polypeptide of SEQ ID NO:64.

Group XXXIII, claim(s)74-75, drawn to polynucleotides comprising SEQ ID NO:65, fragments thereof, polypeptide of SEQ ID NO:66, fragments of the polypeptide of SEQ ID NO:66.

Group XXXIV, claim(s)76-77, drawn to polynucleotides comprising SEQ ID NO:67, fragments thereof, polypeptide of SEQ ID NO:68, fragments of the polypeptide of SEQ ID NO:68.

Group XXXV, claim(s)78-79, drawn to polynucleotides comprising SEQ ID NO:69, fragments thereof, polypeptide of SEQ ID NO:70, fragments of the polypeptide of SEQ ID NO:70.

Group XXXVI, claim(s)80-81, drawn to polynucleotides comprising SEQ ID NO:71, fragments thereof, polypeptide of SEQ ID NO:72, fragments of the polypeptide of SEQ ID NO:72.

Group XXXVII, claim(s)82-83, drawn to polynucleotides comprising SEQ ID NO:73, fragments thereof, polypeptide of SEQ ID NO:74, fragments of the polypeptide of SEQ ID NO:74.

Group XXXVIII, claim(s)84-85, drawn to polynucleotides comprising SEQ ID NO:75, fragments thereof, polypeptide of SEQ ID NO:76, fragments of the polypeptide of SEQ ID NO:76.

Group XXXIX, claim(s)86-87, drawn to polynucleotides comprising SEQ ID NO:77, fragments thereof, polypeptide of SEQ ID NO:78, fragments of the polypeptide of SEQ ID NO:78.

Group XL, claim(s)88-89, drawn to polynucleotides comprising SEQ ID NO:79, fragments thereof, polypeptide of SEQ ID NO:80, fragments of the polypeptide of SEQ ID NO:80.

Group XLI, claim(s)90-91, drawn to polynucleotides comprising SEQ ID NO:81, fragments thereof, polypeptide of SEQ ID NO:82, fragments of the polypeptide of SEQ ID NO:82.

Group XLII, claim(s)92-93, drawn to polynucleotides comprising SEQ ID NO:83, fragments thereof, polypeptide of SEQ ID NO:84, fragments of the polypeptide of SEQ ID NO:84.

Group XLIII, claim(s)94-95, drawn to polynucleotides comprising SEQ ID NO:85, fragments thereof, polypeptide of SEQ ID NO:86, fragments of the polypeptide of SEQ ID NO:86.

Group XLIV, claim(s)96-97, drawn to polynucleotides comprising SEQ ID NO:87, fragments thereof, polypeptide of SEQ ID NO:88, fragments of the polypeptide of SEQ ID NO:88.

Group XLV, claim(s)98-99, drawn to polynucleotides comprising SEQ ID NO:89, fragments thereof, polypeptide of SEQ ID NO:90, fragments of the polypeptide of SEQ ID NO:90.

Group XLVI, claim(s)100-101, drawn to polynucleotides comprising SEQ ID NO:91, fragments thereof, polypeptide of SEQ ID NO:92, fragments of the polypeptide of SEQ ID NO:92.

Group XLVII, claim(s)102-103, drawn to polynucleotides comprising SEQ ID NO:93, fragments thereof, polypeptide of SEQ ID NO:94, fragments of the polypeptide of SEQ ID NO:94.

Group XLVIII, claim(s)104-105, drawn to polynucleotides comprising SEQ ID NO:95, fragments thereof, polypeptide of SEQ ID NO:96, fragments of the polypeptide of SEQ ID NO:96.

Group XLIX, claim(s)106-107, drawn to polynucleotides comprising SEQ ID NO:97, fragments thereof, polypeptide of SEQ ID NO:98, fragments of the polypeptide of SEQ ID NO:98.

Group L, claim(s)108-109, drawn to polynucleotides comprising SEQ ID NO:99, fragments thereof, polypeptide of SEQ ID NO:100, fragments of the polypeptide of SEQ ID NO:100.

Group LI, claim(s)110-111, drawn to polynucleotides comprising SEQ ID NO:101, fragments thereof, polypeptide of SEQ ID NO:102, fragments of the polypeptide of SEQ ID NO:102.

Group LII, claim(s)112-113, drawn to polynucleotides comprising SEQ ID NO:103, fragments thereof, polypeptide of SEQ ID NO:104, fragments of the polypeptide of SEQ ID NO:104.

Group LIII, claim(s)114-115, drawn to polynucleotides comprising SEQ ID NO:105, fragments thereof, polypeptide of SEQ ID NO:106, fragments of the polypeptide of SEQ ID NO:106.

Group LIV, claim(s)116-117, drawn to polynucleotides comprising SEQ ID NO:107, fragments thereof, polypeptide of SEQ ID NO:108, fragments of the polypeptide of SEQ ID NO:108.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/09970

Group LV, claim(s)118-119, drawn to polynucleotides comprising SEQ ID NO:109, fragments thereof, polypeptide of SEQ ID NO:110, fragments of the polypeptide of SEQ ID NO:110.

Group LVI, claim(s)120-121, drawn to polynucleotides comprising SEQ ID NO:111, fragments thereof, polypeptide of SEQ ID NO:112, fragments of the polypeptide of SEQ ID NO:112

Group LVII, claim(s)122-123, drawn to polynucleotides comprising SEQ ID NO:113, fragments thereof, polypeptide of SEQ ID NO:114, fragments of the polypeptide of SEQ ID NO:114.

Group LVIII, claim(s)124-125, drawn to polynucleotides comprising SEQ ID NO:115, fragments thereof, polypeptide of SEQ ID NO:116, fragments of the polypeptide of SEQ ID NO:116.

Group LVIII, claim(s)126-127, drawn to polynucleotides comprising SEQ ID NO:117, fragments thereof, polypeptide of SEQ ID NO:118, fragments of the polypeptide of SEQ ID NO:118.

The inventions listed as Groups I-LVIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The main invention is Group I, which is first product, first method of making the product and first method of using the product. Pursuant to 37 CFR 1.474 (d), these claims are considered by the ISA/US to constitute the main invention and none of the related Groups II-LVIII correspond to the main invention. The products of Groups II-LVIII do not share the same or corresponding special technical feature with Group I because they are drawn to products having materially different structures and functions, each defines a separate invention over the art. Therefore, the claims are not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification <sup>6</sup> : <b>C07H 21/04, C07K 14/705, C12N 15/09, 15/63, C12Q 1/68</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/57132</b>
			(43) International Publication Date: 11 November 1999 (11.11.99)
(21) International Application Number: <b>PCT/US99/09970</b>		(72) Inventors: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). McCOY, John, M.; 56 Howard Street, Reading, MA 01867 (US). LaVALLIE, Edward, R.; 113 Ann Lee Road, Harvard, MA 01451 (US). COLLINS-RACIE, Lisa, A.; 124 School Street, Acton, MA 01720 (US). EVANS, Cheryl; 18801 Bent Willow Circle, Germantown, MD 20874 (US). MERBERG, David; 2 Orchard Drive, Acton, MA 01720 (US). TREACY, Maurice; 12 Foxrock Court, Dublin 18 (IE). AGOSTINO, Michael, J.; 26 Wolcott Avenue, Andover, MA 01810 (US). STEININGER, Robert, J., II; 100 Reed Street, Cambridge, MA 02140 (US). BOWMAN, Michael, R.; 50 Aldrich Road, Canton, MA 02021 (US). DiBLASIO-SMITH, Elizabeth; 17 Chestnut Road, Tyngsboro, MA 01879 (US). WIDOM, Angela; 19 Cherokee Road, Acton, MA 01720 (US).	
(22) International Filing Date: 7 May 1999 (07.05.99)			
(30) Priority Data:			
60/084,564	7 May 1998 (07.05.98)	US	
60/087,645	2 June 1998 (02.06.98)	US	
60/093,712	22 July 1998 (22.07.98)	US	
60/094,935	31 July 1998 (31.07.98)	US	
60/095,880	10 August 1998 (10.08.98)	US	
60/096,068	11 August 1998 (11.08.98)	US	
09/306,111	6 May 1999 (06.05.99)	US	
(71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US).		(74) Agent: MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).	
		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM			
(57) Abstract			
Novel polynucleotides and the proteins encoded thereby are disclosed.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

FIGURE 1A




Plasmid name: pED6dpc2

Plasmid size: 5374 bp

Comments/References: pED6dpc2 is derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning. SST cDNAs are cloned between EcoRI and NotI. pED vectors are described in Kaufman et al.(1991), NAR 19: 4485-4490.

FIGURE 1B



Plasmid name: pNOTs

Plasmid size: 4529 bp

Comments/References: pNOTs is a derivative of pMT2 (Kaufman et al, 1988, Mol. Cell. Biol. 8:1741-1750). DHFR was deleted and a new polylinker was inserted between EcoRI and HpaI. M13 origin of replication was inserted in the ClaI site. GST cDNAs are cloned between EcoRI and NotI

## SEQUENCE LISTING

<110> Jacobs, Kenneth  
 McCoy, John M.  
 LaVallie, Edward R.  
 Collins-Racie, Lisa A.  
 Evans, Cheryl  
 Herberg, David  
 Treacy, Maurice  
 Agostino, Michael J.  
 Steininger II, Robert J.  
 Bowman, Michael R.  
 DiBlasio-Smith, Elizabeth  
 Widom, Angela  
 Genetics Institute, Inc.

<120> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM

<130> GI 6069-74A

<140>

<141>

<160> 160

<170> PatentIn Ver. 2.0

<210> 1

<211> 571

<212> DNA

<213> Homo sapiens

<400> 1

```

ttttttggcca ggtctcttgc tgaactcaagt ttttcagttc agcatctttct agttgcagcg 60
atgagtgcac gagtgcagtc aagatccaga ggaagaggag atggtcagga agtcccgat 120
gtggttgcat tegtgtcttc cgggtgaatct cagcaagaag aaccacccac tgacaatcag 180
gatattgaac ctggacaaga gagagaagga acacctccga tcgaagaacg taaggtagaa 240
ggtgattgac aggaatgga tctggaaag ectcggagtg agcgtggaga tggctctgat 300
gtaaagaga agactccacc taatctaac catgctaaga ctaaagaagc agagatggg 360
cagcctatag ctaaaaagaa gacaaagctg agctacacac atggttgatg tcacattgaa 420
aatgtgactg aaattttgaa aattctctca ataaagtgtg agttttctct gaaaaaaaaa 480
aaaaaaagaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 540
aaaaaaagaa aaaaaaaaaa aaaaaaaaaa a 571

```

<210> 2

<211> 102

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Ser Ala Arg Val Arg Ser Arg Ser Arg Gly Arg Gly Asp Gly Gln
  1                      5                      10                      15

Glu Ala Pro Asp Val Val Ala Phe Val Ala Pro Gly Glu Ser Gln Gln
                20                      25                      30

Glu Glu Pro Pro Thr Asp Asn Gln Asp Ile Glu Pro Gly Gln Glu Arg
                35                      40                      45

Glu Gly Thr Pro Pro Ile Glu Glu Arg Lys Val Glu Gly Asp Cys Gln

```

50	55	60	
Glu Met Asp Leu Glu Lys Thr Arg Ser Glu Arg Gly Asp Gly Ser Asp			
65	70	75	80
Val Lys Glu Lys Thr Pro Pro Asn Pro Lys His Ala Lys Thr Lys Glu			
	85	90	95
Ala Gly Asp Gly Gln Pro			
100			

<210> 3  
 <211> 2709  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 gaggaaaccc ctccgtcggg ctaggagttc gggggggccc ggcggggggg ctgcggagct 60  
 ggcagggtgc aagcgtctgc acctggcgga ccatggcgcc ccatgcgggc gcccggggat 120  
 agcgtcgggg aggcctgcgg gccccggcgc gcacggccgc acctctcccc agccctggcg 180  
 tgggcccgag ccggcccgag cagcaatggg gttccctgcg ctgctggtcg tagcggtgct 240  
 ggcatccgaa caccgggtgg ctggtgcagg cgaggtcttc gggaattcca gcgagggtct 300  
 tattgaattt tctgtgggga aatttagata cttcgagctc aataggccct ctccagagga 360  
 agctattttg catgatattt caagcaatgt gaattttctt atcttccaaa taacttcaca 420  
 gtatcagcat acacctgttt cctttctctc gactctcctt tccaatctct cgggaactagg 480  
 cactgcgcgt ggactggttt tctctcttag accagagcag agtacatgca cttggtactt 540  
 ggggaacttca cgcataccgc ctgtccagaa tatggctatc ctactctctt actcagaaag 600  
 agatectgtc cctggagggt gtaatttggg gttcgattta gatattgac ccaacattta 660  
 ctggagatgt aatttttttg aaacgactat caagtttgcc ccagcaaacg taggctatgc 720  
 ggcagggtta gatcccccac catgtacgc tgggaacgac caggacttca ggtggagggt 780  
 gcagtatgat gtctatccgt atttctgtcc tgagaattgc ctcaatgagg agatgttgc 840  
 gcagcatctg cagaggatgg tcagtgtgac ccaggtgag gccagtgctc tcaaggtggt 900  
 taacctaaac gctaattgta agacaagtgt tctcttctcc tccctccggg gacaaggtgt 960  
 catatataat gtcattgttt gggacccggt tctaaataca tctgctgccc accttctgc 1020  
 tccacataac gcttgcaggt ttgaggcagg agagggtcgt tgtgcttccc taggaagagt 1080  
 gtcttccaaa gtgttcttca cctcttttgc cctgcttggg ttcttcaatt gtctcttgg 1140  
 acacagattc ttgaaaacag acttattctt cataggctt accatcctgg gattcttctt 1200  
 ttatatactg attacaagac tgacaactat caagtatgat gtgaatctga ttctgacgc 1260  
 tgcactggga agcgtcggtg gaattgttct ggtagctgtg ctcatctcgt cagtgcattt 1320  
 ctcgatctgc atgctctgtg ttggactagt gctgggggtc ctcatctcgt cagtgcattt 1380  
 ctttactcca ctgggaaacc taagatttt tcatgatgat ggtgtattct ggtcactttt 1440  
 ctcttgcata gctatctcca ttccagtagt ttccatgggc tgcctaagaa tactgaacct 1500  
 actgaattgt ggagtcattg gctcctatct ggtggtttta gccattgca gttactgggt 1560  
 cacaagctct tctcactca ctttgacgt actcaagaga ggcctcaca aggatttcca 1620  
 cagagcttcc acaaatgtgc ctttccaac taatgactc attctctgg cagtatggg 1680  
 catgctgggt gtaagtggaa ttacgttaca gattcganga gggagaggac gccgttctt 1740  
 cctccccac ccatcaggt tatggaagca agagagagag cgcagagtga caaacattct 1800  
 ggcctctagc taacctatcc ctccattgag agcagggctc tatggccgat taacctagat 1860  
 taaggggctc ttctagaagg agccgccagc tggagagaga accgctttgc ttctgtagat 1920  
 gcccgggggt ttggtcagtg tgcctcagct ttggagtcca tgcctggagt ggttcacag 1980  
 tctctggtgc aggtctcata agagatcagg catatatatc tcttcttgc ataactattt 2040  
 ggtgccttca ttgalatct gtaaggggtg actaggagat taggatgatt gtaagagaat 2100  
 gagaagatg accaanaggt tggtagtag ggggttttt ctatttcca aatanttag 2160  
 aaattacctt ttgggttaca aatctatgat caacttattc attaanatag atacattaaa 2220  
 aaattaaaa actgattctt ctgcagagca ctggtgttcc tttttataac ccttgaaac 2280  
 aagtcctcca cctgagcctg tctaaacttt cggagggagt ttattattga gtctttatct 2340  
 gtgacagtat ttggagattt agggatttga tacttaggcc cttgaaattt agaatacaaa 2400  
 aagagaagca agccagacat ggtgcctcac acctgtaatc ccaatactgg gaggcccgag 2460  
 tggggtatc gcttgggccc aggagtttga gaccgacatg ggcacactga caagaccca 2520

```

tctctgcaga agattaaaa agttggccag gcttggtgag acatgcctgc tccagctcc 2580
cgggagagact gagatggggg gatccctgg agccctgag attgagctg cagtgaacct 2640
tgattgcttc actgactcc agcttgggag acagagacc tctctcgaga aattaaaaaa 2700
aaaaaaaaa 2709

```

&lt;210&gt; 4

&lt;211&gt; 570

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

```

Met Gly Phe Leu Gln Leu Leu Val Val Ala Val Leu Ala Ser Glu His
  1              5              10              15

Arg Val Ala Gly Ala Ala Glu Val Phe Gly Asn Ser Ser Glu Gly Leu
      20              25              30

Ile Glu Phe Ser Val Gly Lys Phe Arg Tyr Phe Glu Leu Asn Arg Pro
      35              40              45

Phe Pro Glu Glu Ala Ile Leu His Asp Ile Ser Ser Asn Val Thr Phe
      50              55              60

Leu Ile Phe Gln Ile His Ser Gln Tyr Gln Asn Thr Thr Val Ser Phe
      65              70              75              80

Ser Pro Thr Leu Leu Ser Asn Ser Ser Glu Thr Gly Thr Ala Ser Gly
      85              90              95

Leu Val Phe Ile Leu Arg Pro Glu Gln Ser Thr Cys Thr Trp Tyr Leu
      100              105              110

Gly Thr Ser Gly Ile Gln Pro Val Gln Asn Met Ala Ile Leu Leu Ser
      115              120              125

Tyr Ser Glu Arg Asp Pro Val Pro Gly Gly Cys Asn Leu Glu Phe Asp
      130              135              140

Leu Asp Ile Asp Pro Asn Ile Tyr Leu Glu Tyr Asn Phe Phe Glu Thr
      145              150              155              160

Thr Ile Lys Phe Ala Pro Ala Asn Leu Gly Tyr Ala Arg Gly Val Asp
      165              170              175

Pro Pro Pro Cys Asp Ala Gly Thr Asp Gln Asp Ser Arg Trp Arg Leu
      180              185              190

Gln Tyr Asp Val Tyr Gln Tyr Phe Leu Pro Glu Asn Asp Leu Thr Glu
      195              200              205

Glu Met Leu Leu Lys His Leu Gln Arg Met Val Ser Val Pro Gln Val
      210              215              220

Lys Ala Ser Ala Leu Lys Val Val Thr Leu Thr Ala Asn Asp Lys Thr
      225              230              235              240

Ser Val Ser Phe Ser Ser Leu Pro Gly Gln Gly Val Ile Tyr Asn Val
      245              250              255

Ile Val Trp Asp Pro Phe Leu Asn Thr Ser Ala Ala Tyr Ile Pro Ala

```

260					265					270						
His	Thr	Tyr	Ala	Cys	Ser	Phe	Glu	Ala	Gly	Glu	Gly	Ser	Cys	Ala	Ser	
275					280					285						
Leu	Gly	Arg	Val	Ser	Ser	Lys	Val	Phe	Phe	Thr	Leu	Phe	Ala	Leu	Leu	
290					295					300						
Gly	Phe	Phe	Ile	Cys	Phe	Phe	Gly	His	Arg	Phe	Trp	Lys	Thr	Glu	Leu	
305					310					315					320	
Phe	Phe	Ile	Gly	Phe	Ile	Ile	Met	Gly	Phe	Phe	Phe	Tyr	Ile	Leu	Ile	
325					330					335						
Thr	Arg	Leu	Thr	Pro	Ile	Lys	Tyr	Asp	Val	Asn	Leu	Ile	Leu	Thr	Ala	
340					345					350						
Val	Thr	Gly	Ser	Val	Gly	Gly	Met	Phe	Leu	Val	Ala	Val	Trp	Trp	Arg	
355					360					365						
Phe	Gly	Ile	Leu	Ser	Ile	Cys	Met	Leu	Cys	Val	Gly	Leu	Val	Leu	Gly	
370					375					380						
Phe	Leu	Ile	Ser	Ser	Val	Thr	Phe	Phe	Thr	Pro	Leu	Gly	Asn	Leu	Lys	
385					390					395					400	
Ile	Phe	His	Asp	Asp	Gly	Val	Phe	Trp	Val	Thr	Phe	Ser	Cys	Ile	Ala	
405					410					415						
Ile	Leu	Ile	Pro	Val	Val	Phe	Met	Gly	Cys	Leu	Arg	Ile	Leu	Asn	Ile	
420					425					430						
Leu	Thr	Cys	Gly	Val	Ile	Gly	Ser	Tyr	Ser	Val	Val	Leu	Ala	Ile	Asp	
435					440					445						
Ser	Tyr	Trp	Ser	Thr	Ser	Leu	Ser	Tyr	Ile	Thr	Leu	Asn	Val	Leu	Lys	
450					455					460						
Arg	Ala	Leu	Asn	Lys	Asp	Phe	His	Arg	Ala	Phe	Thr	Asn	Val	Pro	Phe	
465					470					475					480	
Gln	Thr	Asn	Asp	Phe	Ile	Ile	Leu	Ala	Val	Trp	Gly	Met	Leu	Ala	Val	
485					490					495						
Ser	Gly	Ile	Thr	Leu	Gln	Ile	Arg	Arg	Glu	Arg	Gly	Arg	Pro	Phe	Phe	
500					505					510						
Pro	Pro	His	Pro	Tyr	Lys	Leu	Trp	Lys	Gln	Glu	Arg	Glu	Arg	Arg	Val	
515					520					525						
Thr	Asn	Ile	Leu	Asp	Pro	Ser	Tyr	His	Ile	Pro	Pro	Leu	Arg	Glu	Arg	
530					535					540						
Leu	Tyr	Gly	Arg	Leu	Thr	Gln	Ile	Lys	Gly	Leu	Phe	Gln	Lys	Glu	Gln	
545					550					555					560	
Pro	Ala	Gly	Glu	Arg	Thr	Pro	Leu	Leu	Leu							
565					570											



<210> 5  
 <211> 3063  
 <212> DNA  
 <213> Homo sapiens

<400> 5

```

cgaggccggc gttgtgcccg tggcgggggc gggggagggc gggacaggag gcttcgggga 60
agatggaccc ggcggccctc ctgggctgca gcttcaaggc tgtgaagtgg agctcgggtg 120
ccgtgcccgt cgaacctctg gtcagcactt accggctgac ccagatcggg cgcctgggaa 180
acggagagtg cgtagaaggc ctgggggaaa atgactatct gctgattcat tccctggccc 240
agtggaccac catcaactgc caccagcttg agggagggtc ctatgtcatt gggccaaaga 300
tagagattcc ggtacattat gccgggcaat tcaagctgct ggaaccaagc ccagatatga 360
aggagccagt gcaatatttc aacagtcttg agggaggtgc taaggcattt cctgaacggc 420
tgtacgtcat ggaagatctc acctcaaccg cgaaggctgc ttcagggtga tccaatgaag 480
accttgaaat ttacaaactc accctgtcta ctggggatga actcaactct aatcttcaa 540
gcagaaatcc ttcatgcgaa gcaattonag gaasagtcac gactcaacac aatcttcaa 600
aagattggga agctcaatcc catcagcaag ctgggaaaaa gcamaatgac gtgccttatt 660
tgtatgaatc accggaccac cgaaggcatt agccttccat tccagtgcac gggcagattt 720
agcaccccca agtcccccgg accttcagat gcaaaaaggg ccaacaccca tcccccacat 780
tgtggagaaa accaggcttc ctgtgaatgt gactgtgcaa agccctccac cgggaacccc 840
atacgaacct cacttcatcc gtgaggggca ccgtataag tttgtgaaca tccagaccaa 900
gacgggtggt gtttgtctg tgctggggga caacagatc ctcccctatg ctgaaggagg agagctgggc 1020
gcacttgact gtcaccaagt tcagctccac agaacacctg gtgaaggagg utgaaggagg agagctgggc 1080
cgaacacctg gtccatcact ggctaggtat ctgccaaga cagtccgaca tccatgagta 1140
ttcccgggct gtccgtgatc tgaanaacga ctggaatgaa gaatgcagga gccccagaa 1200
gggtcgggtg tctggccata acccgtgccc caattcgctc agctaagccc ggcagtgaat 1260
caccacgtcc ctccacggac tctcggcttg tgtgtatggc aaccaatctc atggcaacag 1320
tgaggtgaac ctccatggtt gcaaggccct gggggagagt tgggtccctc tacctttccc cagaagctag 1380
catcccgccc catcaggact ctggagatag tggagcgacc agaacttccc taaggagagc tctggctgga 1440
tgagaatcca cccagccatc agcctctcac tgcctctctg agcgaagaga cctcttccc 1500
tcagtttga gggtctgtcc gatccaatg tggacttct cctcttccc 1560
tctgggagca ccagtgaag ctccagatc tccctacct ccaactccc agcctccc 1620
atctgaagcc gtccagagag aatgcaggct cctgaacgcc ccaactgttc cacccccag 1680
cgcaaggcct ttgtccacca gtccctccat cctccctgct ccaactgttc cagcgggga 1740
acagactgca tctcccagcc ccaccktgct ctactattct tccaggctac acacatcgt 1800
cactaaaaat gacttgaag acctgaaatc cactctgtt tccctgtatc catgtaacgc 1860
agtgaacct gattctgtgg acctgaaatc acctctgtt agtctctctg ctgactctgt 1920
gtcctctggg ctctcatgga cttaaccatta ttcaggagca tcagaaagcc agaccaggag 1980
tgacttctct ctggatocaa gcaggagtta tagttacct agacaaaaga cggcaggcac 2040
accaaagaga acctgcccag ccccttttga ttttgatgga tgtgagctcc tggccaggcc 2100
cactagccca gtcaactgag aattcagtag cagcgtctct ggtgtgacaa agtcagccc 2160
ctactctctg gagagccacg atgtgaatc tottgagct ggtgtgacaa agtcagccc 2220
gtcctgccc gtcttaccac ccaggctccc aactctgtg gaagagaggg tgcctccc 2280
aacatctctc ttgcctctga aatttgatgg tgcctgagga gaccccaggt ctgggtcacc 2340
agatctctcg gaggaccagt attttgtaa aaggggcatg caggacatct tctctgctc 2400
ctacccttcc ccatctccgc tccatctcca gctggcccc agatcctctg ggcagggttc 2460
ccctggcag ccactgtctg acctatcagg actctctata paggaagtgt ccaagtccat 2520
acggttccat ggtttgtccg aagatgtcat atcattcttt gttactgaaa agattgatg 2580
gaacctgctt gttcagctaa cggagaagat cctctcagag gatttcaaat tgagcaant 2640
gcagggggag aagataatgc aattcattaa tggctggagg cccaaaatat ngccaaata 2700
ccccgggcca gcatggaaca aactgatac atcctgtgca tagaagggtt gggctgggac 2760
accttccat gttttgcac taanaacct ctctgaaat agggacagga gaactctta 2820
ctatgcagat taagtttttg aatgggtaac aggcattttt gtacatcact acaatgctg 2880
tacagaacac ttgaggtgt gcttgtgac tcaactcaaa aactcagc agctgctaaa 2940
agaaaaaag gctgtgag agaatcatt ctaccccaag taagtttatg cgagaggtta 3000
tgatatttat taaaaaatg ccaaggctga aatctttaa adaaaaaaa 3060
aaa 3063

```

<210> 6

&lt;211&gt; 647

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

```

Met Gln Lys Gly Glu His Thr Ile Arg His Ile Val Glu Lys Thr Arg
 1           5           10           15

Leu Pro Val Asn Val Thr Val Pro Ser Pro Pro Pro Arg Asn Pro Tyr
          20           25           30

Asp Leu His Phe Ile Arg Glu Gly His Arg Tyr Lys Phe Val Asn Ile
 35           40           45

Gln Thr Lys Thr Val Val Val Cys Cys Val Leu Arg Asp Asn Lys Ile
 50           55           60

Leu Pro Met His Phe Pro Leu His Leu Thr Val Pro Lys Phe Ser Leu
 65           70           75           80

Pro Glu His Leu Val Lys Gly Glu Ser Trp Pro Glu Thr Leu Val His
          85           90           95

His Trp Leu Gly Ile Cys Gln Glu Gln Phe Asp Ile Asp Glu Tyr Ser
100           105           110

Arg Ala Val Arg Asp Val Lys Thr Asp Trp Asn Glu Glu Cys Lys Ser
115           120           125

Pro Lys Lys Gly Arg Cys Ser Gly His Asn His Val Pro Asn Ser Leu
130           135           140

Ser Tyr Ala Arg Asp Glu Leu Thr Gln Ser Phe His Arg Leu Ser Val
145           150           155           160

Cys Val Tyr Gly Asn Asn Leu His Gly Asn Ser Glu Val Asn Leu His
165           170           175

Gly Cys Arg Asp Leu Gly Gly Asp Trp Ala Pro Phe Pro His Asp Ile
180           185           190

Leu Pro Tyr Gln Asp Ser Gly Asp Ser Gly Ser Asp Tyr Leu Phe Pro
195           200           205

Glu Ala Ser Glu Glu Ser Ala Gly Ile Pro Gly Lys Ser Glu Leu Pro
210           215           220

Tyr Glu Glu Leu Trp Leu Glu Glu Gly Lys Pro Ser His Gln Pro Leu
225           230           235           240

Thr Arg Ser Leu Ser Glu Lys Asn Arg Cys Asp Gln Phe Arg Gly Ser
245           250           255

Val Arg Ser Lys Cys Ala Thr Ser Pro Leu Pro Ile Pro Gly Thr Leu
260           265           270

Gly Ala Ala Val Lys Ser Ser Asp Thr Ala Leu Pro Pro Pro Pro Val
275           280           285

Pro Pro Lys Ser Glu Ala Val Arg Glu Glu Cys Arg Leu Leu Asn Ala

```

290				295				300							
Pro	Pro	Val	Pro	Pro	Arg	Ser	Ala	Lys	Pro	Leu	Ser	Thr	Ser	Pro	Ser
305					310					315					320
Ile	Pro	Pro	Arg	Thr	Val	Lys	Pro	Ala	Arg	Gln	Gln	Thr	Arg	Ser	Pro
				325					330					335	
Ser	Pro	Thr	Leu	Ser	Tyr	Tyr	Ser	Ser	Gly	Leu	His	Asn	Ile	Val	Thr
			340						345					350	
Lys	Thr	Asp	Thr	Asn	Pro	Ser	Glu	Ser	Thr	Pro	Val	Ser	Cys	Tyr	Pro
		355					360						365		
Cys	Asn	Arg	Val	Lys	Thr	Asp	Ser	Val	Asp	Leu	Lys	Ser	Pro	Phe	Gly
	370					375					380				
Ser	Pro	Ser	Ala	Glu	Ala	Val	Ser	Ser	Arg	Leu	Ser	Trp	Pro	Asn	His
				390						395					400
Tyr	Ser	Gly	Ala	Ser	Glu	Ser	Gln	Thr	Arg	Ser	Asp	Phe	Leu	Leu	Asp
			405						410					415	
Pro	Ser	Arg	Ser	Tyr	Ser	Tyr	Pro	Arg	Gln	Lys	Thr	Pro	Gly	Thr	Pro
			420						425				430		
Lys	Arg	Asn	Cys	Pro	Ala	Pro	Phe	Asp	Phe	Asp	Gly	Cys	Glu	Leu	Leu
		435					440					445			
Ala	Ser	Pro	Thr	Ser	Pro	Val	Thr	Ala	Glu	Phe	Ser	Ser	Ser	Val	Ser
		450				455					460				
Gly	Cys	Pro	Lys	Ser	Ala	Ser	Tyr	Ser	Leu	Glu	Ser	Thr	Asp	Val	Lys
	465				470					475					480
Ser	Leu	Ala	Ala	Gly	Val	Thr	Lys	Gln	Ser	Thr	Ser	Cys	Pro	Ala	Leu
			485						490					495	
Pro	Pro	Arg	Ala	Pro	Lys	Leu	Val	Glu	Glu	Lys	Val	Ala	Ser	Glu	Thr
			500						505				510		
Ser	Pro	Leu	Pro	Leu	Lys	Ile	Asp	Gly	Ala	Glu	Glu	Asp	Pro	Lys	Ser
		515					520					525			
Gly	Ser	Pro	Asp	Leu	Ser	Glu	Asp	Gln	Tyr	Phe	Val	Lys	Lys	Gly	Met
		530				535					540				
Gln	Asp	Ile	Phe	Ser	Ala	Ser	Tyr	Pro	Phe	Ser	Ser	Pro	Leu	His	Leu
	545				550					555					560
Gln	Leu	Ala	Pro	Arg	Ser	Cys	Gly	Asp	Gly	Ser	Pro	Trp	Gln	Pro	Pro
			565						570					575	
Ala	Asp	Leu	Ser	Gly	Leu	Ser	Ile	Glu	Glu	Val	Ser	Lys	Ser	Leu	Arg
			580						585				590		
Phe	Ile	Gly	Leu	Ser	Glu	Asp	Val	Ile	Ser	Phe	Phe	Val	Thr	Glu	Lys
		595				600						605			
Ile	Asp	Gly	Asn	Leu	Leu	Val	Gln	Leu	Thr	Glu	Glu	Ile	Leu	Ser	Glu

610

615

620

Asp Phe Lys Leu Ser Lys Leu Gln Val Lys Lys Ile Met Gln Phe Ile  
 625 630 635 640

Asn Gly Trp Arg Pro Lys Ile  
 645

&lt;210&gt; 7

&lt;211&gt; 892

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

```

ggcaccgagct cgtgcactca tggcgaccog gaacccccc ccccaagact atgaagtgga 60
cgaggactct tatgaagtgt tggatttaac tgagtatgcc agaagccacc agtcgtggaa 120
tcgagctgtt ggccacagth cgggacctat ggtagaaaa tactcagtag ctaccagat 180
tgtaatgggt ggcgttaactg cctggtgtgc aggtttctc tcccgaaag ttggaaacct 240
tgcagcaact gcagtaggtg gtggctttct tctcttcag attgctagtc atagtggcta 300
tgtcagatt gactggaga gattgaaaa agatgtaat aaagcaaaa gacagattaa 360
gacacgagcg aacaaagcag caactgaat caacaattta attgaagag caacagaatt 420
tatcagcag aacattgtga tatccagtga atttgtggga ggtttttgc cgggacttgc 480
atcttaagga catgaatatt ctccataac ggaattcaat atgagaaga agtcggcagc 540
ataaaggcag tctctcaaa gtacatactc cagagtctct agagcaggga gaaacaccta 600
gctggacaat actcaattca caacttagca ttctgacct tgaagcttgg caaactagta 660
tgtgtgtaa aacaccttat atggtatgtg aacagttaga ttcttgagca aaacgtggcl 720
ttcatcgctt tgtaaaaatt tgcacttgtt tagaaactag cctataaat atcacattg 780
gatgtagata tggagagaaa agaatatgt tgggtttatt gcttagcgaa atattctctt 840
tttatttasa taaatgttc tctcttgtgt tttaaaaaa aaaaaaaaa aa 892

```

&lt;310&gt; 8

&lt;311&gt; 155

&lt;312&gt; PRT

&lt;313&gt; Homo sapiens

&lt;400&gt; 8

```

Met Ala Thr Arg Asn Pro Pro Pro Gln Asp Tyr Glu Ser Asp Asp Asp
  1          5          10          15

Ser Tyr Glu Val Leu Asp Leu Thr Glu Tyr Ala Arg Arg His Gln Trp
          20          25          30

Trp Asn Arg Val Phe Gly His Ser Ser Gly Pro Met Val Glu Lys Tyr
          35          40          45

Ser Val Ala Thr Gln Ile Val Met Gly Gly Val Thr Gly Trp Cys Ala
          50          55          60

Gly Phe Leu Phe Gln Lys Val Gly Lys Leu Ala Ala Thr Ala Val Gly
          65          70          75          80

Gly Gly Phe Leu Leu Leu Gln Ile Ala Ser His Ser Gly Tyr Val Gln
          85          90          95

Ile Asp Trp Lys Arg Val Glu Lys Asp Val Asn Lys Ala Lys Arg Gln
          100          105          110

Ile Lys Lys Arg Ala Asn Lys Ala Ala Pro Glu Ile Asn Asn Leu Ile
          115          120          125

```

Glu Glu Ala Thr Glu Phe Ile Lys Glu Asn Ile Val Ile Ser Ser Gly  
130 135 140

Phe Val Gly Gly Phe Leu Leu Gly Leu Ala Ser  
145 150 155

<210> 9  
<211> 1850  
<212> DNA  
<213> Homo sapiens

<400> 9  
caactactact gggggtgcta tgaagcttac tggttgtcat gtgtttataat ttagtctgtt 60  
tttttgattg aatgcagttt aatgtttcca gaagggcaaa gtaattttct ttccagatat 120  
gcaagggttt ggtgggtcca aaaaatgtct atcacaagcc atttttttct ttccctctct 180  
cgaaaagtta aatatctat gtgtttatcc caaacctct taccatagta tctgctctgc 240  
tgtccatcat ctcccttctt ccttatctct gtgtatcttg atggcagccg ctgccacagg 300  
gaatgctctg ggggagggca ggtactgtct ttgctctgtg gtcacagctg gccatccctg 360  
ctgggtgatg ctgggcaaga ccttgggccc gtctgggccc tggcttctct acttctgaaa 420  
tgagcgggaa gatgaactct agttccctcc acctctttag catggtgagg taacagacat 480  
caaaagcttt tctgaactct tcagaagaaa tagttccatt ccagaaaaact ctccaaaaa 540  
aatagtagtg aaaaatttta aaaaactctc ttggagtaag ctttktcaag atgactctcc 600  
acaatggagg cagcgttctt ccttgctcct acacagctga agcatttgtt tcttaggtgt 660  
gaaatcgggg ataaaggaaa aacagagaca cagggcattg ttcataggag gaatcgtcac 720  
ctctctgggt gttctgtggg aatttctctg gtgaggaaaa cgtggcaca ggttctgtct 780  
gtaccaacc tccccaggag agatggccct cggcctgtgc cgtctgttcc acctcgcca 840  
ctccatggaa gtttttggtc tgtttccgga tctgcccctt gccctgaact ctccatcagg 900  
ttgtacctgc ctgctggacc cctccacctg gagggcagcc catgtctcag gccacgccct 960  
agctctctct cctcaaatcc taagtgtttt ccttttaggt ttccctgggt ttgtgaatgg 1020  
atcatgtgtc tctaggtata aacctgacat cactctctca cccggcttac ctccaccaga 1080  
tctcccaagt tctgtctcca tcttctact gcagctgtct tgttctctg gtcactgtct 1140  
catcactgag tctgacccct tgttatcatt tcaaaactgg cctccttccc tegtcccca 1200  
cttctaaag tcaactgtcc attgccacca gattaagctt tctccagcca gatcacctct 1260  
ctctgagaaa cctccattga catggaaaca ccattgtctg gcacacatac tccatactct 1320  
acctctcctt ctgtatccc acacatcttt ccagcctccc cctctactcc actcctgct 1380  
cctctctcca cctcccccct ctcttgtctc cctccctctc tgaatccagc ccagcggggc 1440  
ttctctgccc tccatcacat cacagaagta cctcctgctt ctggttttaa ttagagcctt 1500  
cccgatttac attttctctt gaatttttct ctatctacat ttgctctgtc atgttcaaac 1560  
ccctacttct taagggaact tctctaatct ctatctctca tccccaata gtgttttctt 1620  
cctctggggt cttaaatgtt tggatcctat ctccacagct ttagtcttcc ctgctctgtg 1680  
tgacagttac ctgtgtgcat gtgcaatttc taatttccca ccttagactg tgagcttctt 1740  
aaggcaagaa tcatgcttct tgggtttctg tactctctat ggtgccaac acagtgcctt 1800  
ctacattgca ggcgtgaat aaacattttt aaggcaaaa aaaaaaaga 1850

<210> 10  
<211> 206  
<212> PRT  
<213> Homo sapiens

<400> 10  
Met Ala Leu Gly Leu Cys Arg Cys Phe His Pro Arg His Ser Met Ala  
1 5 10 15  
Ala Phe Gly Leu Phe Pro Ala Leu Pro Ser Ala Leu Asn Ser His Pro  
20 25 30  
Ala Cys Thr Cys Leu Leu Asp Pro Ser Thr Trp Arg Pro Ala His Val  
35 40 45

Ser Gly Pro Ala Leu Ala Ser Ser Pro Gln Ile Leu Ser Val Phe Ser  
 50 55 60  
 Leu Gly Phe Pro Gly Phe Val Asn Gly Ser Cys Val Ser Arg Tyr Lys  
 65 70 75 80  
 Pro Asp Ile Ile Ser Pro Pro Gly Leu Pro Pro Pro Asp Leu Pro Ser  
 85 90 95  
 Ser Val Ser Ile Phe Tyr Leu Gln Leu Leu Cys Ser His Gly His Cys  
 100 105 110  
 Cys His Thr Glu Ser Gly Pro Leu Leu Ser Phe Ser Asn Trp Pro Pro  
 115 120 125  
 Ser Leu Val Pro His Phe Leu Lys Ser Pro Val His Cys His Gln Ile  
 130 135 140  
 Lys Leu Ser Pro Ala Arg Ser Pro Leu Ser Glu Lys Pro Pro Leu Thr  
 145 150 155 160  
 Trp Lys His His Cys Leu Ala His Ile Leu Thr Tyr Ser Pro Ser Arg  
 165 170 175  
 Leu Asp Pro His Thr Ser Phe Gln Pro Pro Leu Pro Leu His Ser Leu  
 180 185 190  
 Leu Pro Pro Pro Pro Pro His Pro Leu Val Ser Pro Pro Leu  
 195 200 205

<210> 11  
 <211> 2216  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 cttgttaagt ttgttttagt aattgttttt taactttcat ttantasttg ctgctaaagg 60  
 tcatgttttc tgataaaca ttttaaaatt tttttgtttt gaadagtaa tttatccccc 120  
 atgatgttag atacatttaa attatttagt ctcttcagag atgagatggg gacagggaag 180  
 tatcttgagg cttaacatat tatttagccc aatanaagat gcattgaagc tottatatat 240  
 tatgagtttg aaaaattttg aaggttagcat attgaagtga totataaata tcttcagttc 300  
 tctctgaagt gtgggtattt cttctatcta aaaaatacat acagtgaactg tottcaaatc 360  
 taatttggtc ttgacaaat argagctaat gggtctcaact cttttgccca ggcctggagt 420  
 tgtttgtttg tttttgttt ttttttttaa ggtctcaact cttttgccca ggcctggagt 480  
 cagtggcaca atcagggttc ccaggetaat gtttttattt ttaatttgta attttttttt 540  
 tttttttttt gttgagacgg agttgctcca tgttgcaacg gctgttctca aactcctang 600  
 ctcaaggcat ctgctgcat tggctccca aagtgtggg attgtagaca taagccact 660  
 caacagagct atgaatatct tcttaactk gtaagaatga ggtatgttt ccatcagttc 720  
 aatacagata tttttcttcc ctccaaaaca gtttattttg attgtttatt ttattttgat 780  
 tgtacatccg tctaatcttg acatggaaaa tgcatacatc tatgaasact tagctgaag 840  
 ggaagaattg ttttagaag acatatttta aacacccgca ctgcaatat attgtatcct 900  
 tatagttatt tcttaaatg ctgttttcca aacatttcct tttcaccctg ttgtgtggct 960  
 tagaccatc tegttaattg ttaattggaa agaggctaca gacaccagca gtgtggttc 1020  
 tgcaggtaca cgtgcaaaa gtaattcctg ctcatccatg cctgtctct gtctctctta 1080  
 ggtcctatcc ttatttgagt ataggtggct taattttgct agacttctg aaaaactaa 1140  
 ggtggagtat cagaagtgat tttagtcaca gttctgagg agagcttaga ataacatcct 1200  
 ctttggag gtgtcttgg gtgcgtggat ctgtgtatca agtctttatt gtaagtctg 1260  
 tccaaaatgc taataaattt aatgttttcc ttccttaatt tcttgcaata gtcttcagg 1320

```

tagcacctca ttttattaa tgatattggg attacataag aacaaagctat atgtagacat 1340
ttgcatttaa ggaatttga gtgtttcaaa gatcccatca ttgcagcttg tatcctttag 1440
atccaatggg aaactttctg agtcttacct taatgtctat ttgagctaat tagtaacttg 1500
tttaaacaga tttagcaata ctttaaaagt actgtagact atttatgtat agatagatca 1560
tattacccat taaaagtctg ggggaaaaaa ttttttaatt ttactcttct tatgtactga 1620
aactttttt taaaaagggt gatgatgaag tttctctgt agcagcagcg cagctatgct 1680
ttaaacaca caaaaggctg tgtccaggtg cagctcctt cacccttctt gcccacgggtg 1740
aggattgaat aaccaggact tggggatatk gtttgtgtc agggttattt tgtgtggtga 1800
ggatattttg ttacacattt atacattttc tttttccact caagtgaagt tctatcttga 1860
ggcctatagc caaagtgaat aacttggtgt ttcaaggaa aattgtcttc cagaactcca 1920
ctgtcatcac tttaaccaca gtggaaagtt gcattgaat gcacagatc taatattcaa 1980
tgttctgcta ctttgttaag gaagtccagg taaaaaatag atttaataa ttgaatttat 2040
ttgtacatat gcagagtaag gtaattctgt atggaaactg ctttatctct ttttttcca 2100
actctgatga gtgaattatt aaatgtgttg ttatggaaat acagattatt gcttctatag 2160
gaagataatt atgaaataa aacctgaac tatataata taaaaaana aaaaaa 2216

```

&lt;210&gt; 12

&lt;211&gt; 126

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

```

Met Leu Phe Ser Lys His Ser Phe Phe Thr Leu Leu Cys Gly Leu Asp
  1                      5                      10                      15

```

```

Pro Ser Arg Asn Leu Leu Ile Gly Lys Arg Leu Gln Thr Pro Ala Val
          20                      25                      30

```

```

Cys Val Leu Gln Val His Ala Ala Lys Val Ile Pro Ala His Pro Cys
          35                      40                      45

```

```

Pro Val Ser Val Ser Phe Arg Val Ile Pro Tyr Leu Ser Ile Gly Gly
          50                      55                      60

```

```

Leu Ile Leu Leu Asp Phe Leu Lys Thr Leu Arg Trp Ser Ile Arg Ser
          65                      70                      75                      80

```

```

Asp Phe Ser His Ser Ser Ala Gly Glu Leu Arg Ile Thr Ser Ser Phe
          85                      90                      95

```

```

Gly Arg Trp Ser Trp Val Arg Gly Ser Trp Tyr Thr Val Phe Ile Val
          100                      105                      110

```

```

Ser Leu Ile Gln Asn Ala Asn Lys Phe Asn Val Phe Leu Pro
          115                      120                      125

```

&lt;210&gt; 13

&lt;211&gt; 1426

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

```

ctgggtctcc agggggagag cctggcctcg tcttttgcta cccagggctg ccccaggcc 50
catgaagcca ataggagagc gtgtggcact gggccacaaa ctgtccctgt cctgtcttcc 120
tcccagacca tggcctctgc tagctccacc ttgaaggagc ccccacatc cctccctcca 180
tcccagagat gccaccactt gtgtctccac aatgtgtccc tgcacccccc ggttcgcgac 240
tgtccgaccc ctgcacacca ctcatgtcac cagggcgtgc atcatgttca tcccactcta 300
tttatctcag cctttctttg ctgtctgggc attttgtatg tagagcagtt gaaacagaa 360
cctcagacct taacatctgt cctgatgtta agtgccttt catgaccac cgttatcta 420

```

```

tgtatatgta aagtttaagga tggagatctta agtttaccat taataaactca gtaactaata 480
tttaatatcc tactegaget ttatgggaagc aaatcatgac atgtatgtgt gtcggtgtgt 540
gcaagctttg aaactccttc cacagccgca tcttctcatg acacaaaggt tttgataagt 600
actttcctgt gggctgctca gggcctcata gcatctcatt caattacaag aatagaggcc 660
agacacgggt gggcatgcct gtatgccag ataactggga ggtcaggga ggaggatcac 720
tgagagccag gagattgagg ctgcagtga catgatcgcg acactgcaat ccagcctggg 780
tgacgggtgag actttgtctc aaaaaaaca aaaaaacaa tggagggcag acagcaagtc 840
cttgaggaca cctcacacag tgtcctgtag ctaagtgtct aggsaaaaac aaaaactcca 900
aacctctcag tggatgaggc caaggtcgca gaaaggcatt ctgttgacag atgaaacgac 960
gaaagctggc cagacccctc tgtatgcctc tgccttctc ctgtgggttg aggggtctc 1020
acccggaggc cactacagc aggaagtga gctgccatgt tctcttgaga cacagctgcc 1080
ctcctccagc tctgtccctg tctcacctg cgggtggggc agaatcctct cctgggata 1140
agcctccaca gcccgttta tcagaacac aggcaggaa attggaaact ccccccagcc 1200
cagcatgtg gtcaattgg ttggttgcgc tgcagttgt ctcttgctt tgttaaggct 1260
tttaataagt acgtttggca caatgtctt taatggatt gtaataattg taacggtttt 1320
agcagcctat aactttcag ctggtgctt tacttagga aaaaacaa ttgttaatac 1380
agaacattgt ttataagata taacctaga aaaaaaaa aaaaaa 1426

```

&lt;210&gt; 14

&lt;211&gt; 80

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

```

Met Pro Pro Leu Val Ser Pro Gln Cys Ala Pro Ala His Pro Gly Ser
  1                      5                      10                      15

```

```

Ala Leu Ser Asp Pro Cys Thr Pro Leu Met Ser Pro Arg Arg Ala Ser
      20                      25                      30

```

```

Cys Ser Ser Pro Ser Ile Tyr Leu Ser Leu Ser Leu Leu Val Gly His
      35                      40                      45

```

```

Phe Val Cys Arg Ala Val Glu Asn Arg Thr Ser Glu Leu Asn Ile Cys
      50                      55                      60

```

```

Pro Asp Val Lys Val Leu Phe Met Thr Thr Leu Leu Ser Met Tyr Met
      65                      70                      75                      80

```

&lt;210&gt; 15

&lt;211&gt; 2364

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

```

gaagcggctg ctgtaggcgc ggaaggagcg agcgggctg cggagcgggc gacagtggcg 60
tggaatctgc ctctctgaga gaggctggga gggcgggcgg cggcggcatg agcgggggca 120
ccctttacat cggcagcaag atcagcctca tctccaagga ggagatccgc tacgagggca 180
tctctacac catcgacccc gaaactcca cgttagccct tccccagtt cgtacctttg 240
gtacagaaga cagaccggca gatcgtccaa taaccctcgc agatgaagtc ttgcaataca 300
tttatctcgg tgggagtga acataagacc ttactgtttg tgagcaccac aaaccatagt 360
gttctttgac tcaagaccca gctattgttc agtctcact aggttcacgc acttcttcat 420
tccagtcctt gggttcttat ggaactttcg gcaggatgcc acatcacagt cagttcagtc 480
cgagttcctt agttgggcag cagtttgggt ctgttggtgt tcttggaagc tctttgacac 540
cctttgggac agaaccatca accagtggta ccttaccaca aagtagtgcg gttggttctg 600
cctttacaca ggatacaga tctctaaaa cacagttaac taagggtcgc tcaagccctc 660
agttagcccc tttgagaaaa agcccaccca tggaaacagg agtcagagac gctcagccc 720
actcacctgc tccagcagct gttggggaga gggctcctgt atcaaccagg cctttgcctt 780
ctgcacagca aaaggcagga gagaatcagg agcacaggca agcttgaagta cacaaagttt 840

```



```

caaggccaga aaatgagcaa ctccgaaatg ataacagag acaagtagct ccagggtgctc 900
cttcagctcc aaggagaggg cgtgggggac atcggggtgg caggggaaga tttgggtatc 960
ggcgagatgg gccaatgaaa tttaggaag actttgactt tgaaagtgc aatgcacast 1020
tcacacagga agagattgac agagaggttc atataaact caattaaaa gaagstaaa 1080
ttgagaaaaa ggagaagcct gtaaatggtg aagataaagg agactcagg gttgataccc 1140
aaaacagtga aggaatggc gatgaagag atccacttg acctaatgc tattatgaca 1200
aaactaaetc cttctttag atattttctt gtgatgaca tagagagcg agaccaact 1260
gggctgagga aagaagatta aatgctgaa catttggaat cccacttct ccaaacctg 1320
ggcggtgggg atacagaggg agaggaggtc ttgggttcgg tggtagcaga gggcggtgtg 1380
gtggcagagg tggtaacctc actgcccccc gaggatttc cggtaggttc agcggaggtc 1440
gtggggggcg ggagtttgg gattttgaat ataggaaaac cacagctttt ggaccctaaa 1500
aggtctggat tgatctact gcttctgaa agaaagacaa caagttgct gcatagtcta 1560
caacaaagtc tctgaaaaa ggtgaatttc tagctctca tggctctga cattgattec 1620
agcttctgca aagaatgaag agtgaaatc gctgtacat tgtcaccagc actgggtttt 1680
tggtttttgt ttgtttttcc gtttaatttc aaagataaaa tgcagttact ttgggggtg 1740
gaaggttcat cttaaacat gagcattaaa tatatttga atagcagaag gtttaagta 1800
ttcttatgta tagttaaact aaagcagtac ttcaagtgga ctlaacaagt attttttcat 1860
cactgaagg tttttttttt ttatcccta dattgtattt ggcatttgc agttgctgc 1920
agataaggcc gtgatctgt gttttgagc acagaaggtt gtgtgtgtgt gtgtgtgtgt 1980
gtgtgtgtgt gtgtgtgtgt gtatgtgtgt gtctcttcc tctttcttct tggggatoc 2040
tgtatatga gtagcttat ttgtcaatc nattaggtg ctggttggtt gagsatttt 2100
ttagtcaact atgtacacac agtaataact gtttcttag caaggtaac tttttatat 2160
agttgtaaaa ttccattata ttccattgac aaagaacat taagccctt gtatagctgt 2220
ctaaaaagca actaattttt taagaataa acattttta agtccaaaa aaaaaaaaa 2280
aaaaaanaaa aaaaaanaaa aaaaaanaaa aaaaaanaaa aaaaaanaaa aaaaaanaaa 2340
aaaaaanaaa aaaaaanaaa aaaa 2360

```

&lt;210&gt; 15

&lt;211&gt; 463

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

```

Met Ser Gly Gly Thr Pro Tyr Ile Gly Ser Lys Ile Ser Leu Ile Ser
  1             5             10             15

```

```

Lys Ala Val Ile Arg Tyr Glu Gly Ile Leu Tyr Thr Ile Asp Thr Glu
          20             25             30

```

```

Asn Ser Thr Val Ala Leu Ala Lys Val Arg Ser Phe Gly Thr Glu Asp
      35             40             45

```

```

Arg Pro Thr Asp Arg Pro Ile Pro Pro Arg Asp Glu Val Phe Glu Tyr
      50             55             60

```

```

Ile Ile Phe Arg Gly Ser Asp Ile Lys Asp Leu Thr Val Cys Glu Pro
      65             70             75             80

```

```

Pro Lys Pro Gln Cys Ser Leu Pro Gln Asp Pro Ala Ile Val Gln Ser
          85             90             95

```

```

Ser Leu Gly Ser Ser Thr Ser Ser Phe Gln Ser Met Gly Ser Tyr Gly
      100             105             110

```

```

Pro Phe Gly Arg Met Pro Thr Tyr Ser Gln Phe Ser Pro Ser Ser Leu
      115             120             125

```

```

Val Gly Gln Gln Phe Gly Ala Val Gly Val Ala Gly Ser Ser Leu Thr
      130             135             140

```

Ser Phe Gly Thr Glu Thr Ser Asn Ser Gly Thr Leu Pro Gln Ser Ser  
 145 150 155 160  
 Ala Val Gly Ser Ala Phe Thr Gln Asp Thr Arg Ser Leu Lys Thr Gln  
 165 170 175  
 Leu Ser Gln Gly Arg Ser Ser Pro Gln Leu Asp Pro Leu Arg Lys Ser  
 180 185 190  
 Pro Thr Met Glu Gln Ala Val Gln Thr Ala Ser Ala His Leu Pro Ala  
 195 200 205  
 Pro Ala Ala Val Gly Arg Arg Ser Pro Val Ser Thr Arg Pro Leu Pro  
 210 215 220  
 Ser Ala Ser Gln Lys Ala Gly Glu Asn Gln Glu His Arg Gln Ala Glu  
 225 230 235 240  
 Val His Lys Val Ser Arg Pro Glu Asn Glu Gln Leu Arg Asn Asp Asn  
 245 250 255  
 Lys Arg Gln Val Ala Pro Gly Ala Pro Ser Ala Pro Arg Arg Gly Arg  
 260 265 270  
 Gly Gly His Arg Gly Gly Arg Gly Arg Phe Gly Ile Arg Arg Asp Gly  
 275 280 285  
 Pro Met Lys Phe Glu Lys Asp Phe Asp Phe Glu Ser Ala Asn Ala Gln  
 290 295 300  
 Phe Asn Lys Glu Glu Ile Asp Arg Glu Phe His Asn Lys Leu Lys Leu  
 305 310 315 320  
 Lys Glu Asp Lys Leu Glu Lys Gln Glu Lys Pro Val Asn Gly Glu Asp  
 325 330 335  
 Lys Gly Asp Ser Gly Val Asp Thr Gln Asn Ser Glu Gly Asn Ala Asp  
 340 345 350  
 Glu Glu Asp Pro Leu Gly Pro Asn Cys Tyr Tyr Asp Lys Thr Lys Ser  
 355 360 365  
 Phe Phe Asp Asn Ile Ser Cys Asp Asp Asn Arg Glu Arg Arg Pro Thr  
 370 375 380  
 Trp Ala Glu Glu Arg Arg Leu Asn Ala Glu Thr Phe Gly Ile Pro Leu  
 385 390 395 400  
 Arg Pro Asn Arg Gly Arg Gly Gly Tyr Arg Gly Arg Gly Gly Leu Gly  
 405 410 415  
 Phe Arg Gly Gly Arg Gly Arg Gly Gly Gly Arg Gly Gly Thr Phe Thr  
 420 425 430  
 Ala Pro Arg Gly Phe Arg Gly Gly Phe Arg Gly Gly Arg Gly Gly Arg  
 435 440 445  
 Glu Phe Ala Asp Phe Glu Tyr Arg Lys Thr Thr Ala Phe Gly Pro  
 450 455 460

<210> 17  
 <211> 2760  
 <212> DNA  
 <213> Homo sapiens

<400> 17  
 tgaagatgcc tccctctgat cctactgctt tgagctgctc tctatggtt tagcactgag 60  
 tggetctaac gttggcgggc aatctctggc tcaacagcta accctgcttc aggatctctt 120  
 ctgcctgctt caccacagct ctcttagagc ccagagacag gtaacctctt tactaagaag 180  
 agttttgctt gaagtaaccc ctagtctgtc gccagcctc ataggagtg aatctctccc 240  
 cccagcagat atcagtgata tcaatcactc aacagtgaaa gtagattgga atagctggg 300  
 tatcttggac atgtttctag gatgcattgc caaagcactc actgtacagc taanagccaa 360  
 aggaacccac atcaactgga cagctggtac cactgtgggc aaaggagta caacagttac 420  
 tcttcagatg attttcaatt ccagttatct ccgacgaggt gaaagtcatt agtggatgaa 480  
 ggtctcaacc cctacccaga tctcagagat catcattaa cctaccaagg atatggcagg 540  
 aggtcatctg tccgaagctt ggtcccgagt gataaaaat gctattgcag aacctatct 600  
 tgccttgacc aagatggag agaatcttag gtctccagtc agatgtattg caacaactag 660  
 actctggctt gctctcgcat cctatgtgt tcttgatcag gaccacgtag atcgtctctc 720  
 ctgggggaga tggatgggaa aggatggaca acaaaaacca atgctatgt gtgataccac 780  
 tgatgatggg gaaactgcag caatcatttt atgcaatgtc tctggaact tatgtacaga 840  
 ctgtgacaga tctctcacc tctcagag atcaaaact catcaagac aggtcttcaa 900  
 agaagaagaa gaagctataa aggttgacct tcatgaaggt tctggtagaa ccaattggt 960  
 ctggttgatg gcactggcag attctaaaac atgaaggca atgttggaat tccgagaaca 1020  
 caccggcaaa cccatccaga gtatgtcaga agcatgtcgc tctctggtt cccggagtgg 1080  
 aacagctta tctgtctgt gacgtctttg tctgtatgca gattgcacgg aatacgttaa 1140  
 gatagcctgt agtaagagc atccttctgt ccattccatgc gggggtgtta aaacgaaga 1200  
 gcactgctc cctgtctac acggtctgtg caaangtgc acaagctctg agcaagacgc 1260  
 cgtgacatg tgcattgat gttcaccga agcgtctctg gcgcacacag ccatteagct 1320  
 ggattgtagt caactattcc acttaccgtg ctgtgggaga gtattagaaa ccaattggt 1380  
 tggcccaagg ataacattt gatcttatcc ttgatccaat aaagaacct tatgaggatg tcagaagaaa 1440  
 catagtacta aaagaacct agatttgaa atgaaggtct gcataagagt gaagctatca caactcttg 1500  
 agccttaatt agatttgaa atgaaggtct gcataagagt agatattgat agatattgat 1560  
 tgtgaggctt tataatgac cagctggcta tgcattgaat agatattgat agatattgat 1620  
 ctaceaattg agaaaggcat attttggttg tgaagctctg tgcattgatg aggtctggag 1680  
 gggagatgat tatgatcca gagagctcat ctgtgtgctg tcttctgatg ttccaggggc 1740  
 tcatgtgtt cccaaacatg gcccagactt tttggaatat aatgtctgct actgtctt 1800  
 agtggctgtt tttttctgt ttggaacaa cactttttgt aatgtctgtc atgtgattt 1860  
 tcaagaaatg actagcatt ctaaggaaag actaccacac tptcttgcag gtcccaagg 1920  
 caagcagtta gaaggcaatg aatgtccact ccatgttgtt catccacca ctgggggaag 1980  
 gtttgccttg ggtgtgtgag tctgcagaaa tgcacacact ttttagaaga cgcagatcct 2040  
 ttgtctacag agagaasat tgccttcatc ccccaaggag atgctgtgaa gtttaacct 2100  
 tgtccaggat aaggacggga ccattttttac atocatgaaa atgmacpatt caccagtgc 2160  
 gaaggatacc aaataccatg tacataattc ttgtatgaa aggtttccc attattttg 2220  
 tttatctctt tttgacaaa tgaactcaa ctgtgaggt ttgtgcatgt ggcattacc 2280  
 gtcattggcc tgtgaagcat tggacattta tagataattg atataaaag atcgcattg 2340  
 ccatggacta agaacgatg tggctttcaa gcaaaaaaga aaatatntca ttgtttatt 2400  
 tatactgctt tttgtatc ctgtacattc gcatcacggg tgggcataaa aagaggata 2460  
 ttttggttta tttcttagac tgtattttaa aaaaaaaa acattgtgtt aggacagct 2520  
 ataaatgtaa caagtatcac actgtatata acatatcaa tgtttgtcct gtataagaat 2580  
 tactaaatta caaatgcaat ttcatttcaa cctctaggtt aggtttgagc ctgaatttt 2640  
 aatgaagtgc aatactgagt gtgcctcatt atcttcagc cgtaaaata ttggaatgta 2700  
 catgtcaata aaacactgt acatttttat acagtgaaca agtctaaaaa 2760

<210> 18  
 <211> 660  
 <212> PRT  
 <213> Homo sapiens

<400> 18

Met Val Leu Ala Leu Ser Gly Ser Asn Val Gly Arg Gln Tyr Leu Ala  
 1 5 10 15  
 Gln Gln Leu Thr Leu Leu Gln Asp Leu Phe Ser Leu Leu His Thr Ala  
 20 25 30  
 Ser Pro Arg Val Gln Arg Gln Val Thr Ser Leu Leu Arg Arg Val Leu  
 35 40 45  
 Pro Glu Val Thr Pro Ser Arg Leu Ala Ser Ile Ile Gly Val Lys Ser  
 50 55 60  
 Leu Pro Pro Ala Asp Ile Ser Asp Ile Ile His Ser Thr Glu Lys Gly  
 65 70 75 80  
 Asp Trp Asn Lys Leu Gly Ile Leu Asp Met Phe Leu Gly Cys Ile Ala  
 85 90 95  
 Lys Ala Leu Thr Val Gln Leu Lys Ala Lys Gly Thr Thr Ile Thr Gly  
 100 105 110  
 Thr Ala Gly Thr Thr Val Gly Lys Gly Val Thr Thr Val Thr Leu Pro  
 115 120 125  
 Met Ile Phe Asn Ser Ser Tyr Leu Arg Arg Gly Glu Ser His Trp Trp  
 130 135 140  
 Met Lys Gly Ser Thr Pro Thr Gln Ile Ser Glu Ile Ile Ile Lys Leu  
 145 150 155 160  
 Ile Lys Asp Met Ala Ala Gly His Leu Ser Glu Ala Trp Ser Arg Val  
 165 170 175  
 Thr Lys Asn Ala Ile Ala Glu Thr Ile Ile Ala Leu Thr Lys Met Glu  
 180 185 190  
 Glu Glu Phe Arg Ser Pro Val Arg Cys Ile Ala Thr Thr Arg Leu Trp  
 195 200 205  
 Leu Ala Leu Ala Ser Leu Cys Val Leu Asp Gln Asp His Val Asp Arg  
 210 215 220  
 Leu Ser Ser Gly Arg Trp Met Gly Lys Asp Gly Gln Gln Lys Gln Met  
 225 230 235 240  
 Pro Met Cys Asp Asn His Asp Asp Gly Glu Thr Ala Ala Ile Ile Leu  
 245 250 255  
 Cys Asn Val Cys Gly Asn Leu Cys Thr Asp Cys Asp Arg Phe Leu His  
 260 265 270  
 Leu His Arg Arg Thr Lys Thr His Gln Arg Gln Val Phe Lys Glu Glu  
 275 280 285  
 Glu Glu Ala Ile Lys Val Asp Leu His Glu Gly Cys Gly Arg Thr Lys  
 290 295 300  
 Leu Phe Trp Leu Met Ala Leu Ala Asp Ser Lys Thr Met Lys Ala Met  
 305 310 315 320

Val Glu Phe Arg Glu His Thr Gly Lys Pro Thr Thr Ser Ser Ser Glu  
 325 330 335  
 Ala Cys Arg Phe Cys Gly Ser Arg Ser Gly Thr Glu Leu Ser Ala Val  
 340 345 350  
 Gly Ser Val Cys Ser Asp Ala Asp Cys Gln Glu Tyr Ala Lys Ile Ala  
 355 360 365  
 Cys Ser Lys Thr His Pro Cys Gly His Pro Cys Gly Gly Val Lys Asn  
 370 375 380  
 Glu Glu His Cys Leu Pro Cys Leu His Gly Cys Asp Lys Ser Ala Thr  
 385 390 395 400  
 Ser Leu Lys Gln Asp Ala Asp Asp Met Cys Met Ile Cys Phe Thr Glu  
 405 410 415  
 Ala Leu Ser Ala Ala Pro Ala Ile Gln Leu Asp Cys Ser His Ile Phe  
 420 425 430  
 His Leu Gln Cys Cys Arg Arg Val Leu Glu Asn Arg Trp Leu Gly Pro  
 435 440 445  
 Arg Ile Thr Phe Gly Phe Ile Ser Cys Pro Ile Cys Lys Asn Lys Ile  
 450 455 460  
 Asn His Ile Val Leu Lys Asp Leu Leu Asp Pro Ile Lys Glu Leu Tyr  
 465 470 475 480  
 Glu Asp Val Arg Arg Lys Ala Leu Met Arg Leu Glu Tyr Glu Gly Leu  
 485 490 495  
 His Lys Ser Glu Ala Ile Thr Thr Pro Gly Val Arg Phe Tyr Asn Asp  
 500 505 510  
 Pro Ala Gly Tyr Ala Met Asn Arg Tyr Ala Tyr Tyr Val Cys Tyr Lys  
 515 520 525  
 Cys Arg Lys Ala Tyr Phe Gly Gly Glu Ala Arg Cys Asp Ala Glu Ala  
 530 535 540  
 Gly Arg Gly Asp Asp Tyr Asp Pro Arg Glu Leu Ile Cys Gly Ala Cys  
 545 550 555 560  
 Ser Asp Val Ser Arg Ala Gln Met Cys Pro Lys His Gly Thr Asp Phe  
 565 570 575  
 Leu Glu Tyr Lys Cys Arg Tyr Cys Cys Ser Val Ala Val Phe Phe Cys  
 580 585 590  
 Phe Gly Thr Thr His Phe Cys Asn Ala Cys His Asp Asp Phe Gln Arg  
 595 600 605  
 Met Thr Ser Ile Pro Lys Glu Glu Leu Pro His Cys Pro Ala Gly Pro  
 610 615 620  
 Lys Gly Lys Gln Leu Glu Gly Thr Glu Cys Pro Leu His Val Val His  
 625 630 635 640

Pro Pro Thr Gly Glu Glu Phe Ala Leu Gly Cys Gly Val Cys Arg Asn  
645 650 655

Ala His Thr Phe  
660

<210> 19  
<211> 1649  
<212> DNA  
<213> Homo sapiens

<400> 19  
gattgtacat agtcttctgg ggcattgggg agccggctgg aggttggagac cctccccctct 60  
ccccccccc cccgggggag gcaaatgtaa aactactaat ttttgtgctt tatatattct 120  
atataaatat atctattttc tttttacaaa accagtttat aaatggtagg ggggtgtggg 180  
ggggaacacat ggagctcccc ttgtgggggg gccccctcca ttaccogacc taacgccctt 240  
ttcttcaccc cccacccccc tccccacccc ctggctgtga ctgctgtgag atgggggtat 300  
agaggttggg caattccccc cccctgttgt atagtggac tatgtttata cgcacaaaaag 360  
agagctgacc ccaggggggg ccagaggggt atgggttctt tgcctccctt tcttccctct 420  
ttttgcccga gcttctgtcg cagttgaacc tcttcttggg ggtgggagta ggttaagggt 480  
gggtgagggc ccaaacctct ctctggtagg gaacgttggg gatgaagatg aagcttatat 540  
gcagttctct ctatgggggt gtggggcaag ggcatttgtt aattaatatt tccaagaatc 600  
agctgtcttg agtctagggg tgggcttggc ggtggtggac gggcgggcct gctggagggg 660  
gaacttgggc gctgttctga ttttaggttt gtttttgttt tgttttgaat ttggggggtt 720  
gtgatttgtt gggggtaggg agattttttt ttttttaagg ctgctccctc aactgttttc 780  
agctgcacat gtttaagaga ataacagccc ccactccccc aggaacccgt gtaattaat 840  
cagacagtag gaagcctggg ctgctgccc ccaagccaca gcccttggat gttccttttc 900  
cgaagagcag aggtctaggc tccagggagg gggagatttg ctccctgtag tccggctgtg 960  
tttggggctt gggccctggg atgggcaaaa ggggatgggg cagactttgt aagcatatgc 1020  
caggtatccg atagtctgtt agaatttagt gtagaacc ttatcagttt ttaattttta 1080  
tatcaactat aactcagacc caagctacaa ggttgggaatt ttggttgggt ttttttttaa 1140  
gtaccttgcg tgtatatttg ctctcagacc cccaccccca ccccgagccc cagtgtttgt 1200  
attttgggtt gttttatatt cgcacatact cagttttcag ttttccctt tacagtcttc 1260  
tccccctacc tccaggaccc tcccccttt taaaaaataa atcgctgaca agtgtgaatc 1320  
cagtgaagac ttatatttgt gttgtgtgta tctgtacag caaggttggg ccttcgtaac 1380  
aacggatgaa atggttccct ttttttaagg ggcctctctc cctccaccct cagcgccctt 1440  
gtccttggca tgttttgtat cagcgatcat ctggaactgt acatatttat gttgcgagag 1500  
gcaaaaggga agtttcggat ttgtcttctt ccaagtttgt ttttaacgta caaataaana 1560  
aaqaacattt taatataaaa aaaaaaanaa aaaaaaanaa aaaaaaanaa aaaaaaanaa 1620  
aaaaaaabaa aaaaaaanaa aaaaaaanaa 1649

<210> 20  
<211> 92  
<212> PRT  
<213> Homo sapiens

<400> 20  
Met Gly Glu Pro Ala Gly Gly Glu Asn Pro Pro Leu Ser Pro His Pro  
1 5 10 15  
Pro Gly Arg Ala Asn Val Lys Leu Leu Ile Phe Val Leu Tyr Ile Phe  
20 25 30  
Tyr Ile Asn Ile Ser Ile Phe Phe Leu Gln Asn Gln Phe Ile Asn Gly  
35 40 45  
Arg Gly Val Trp Gly Gly His Met Glu Leu Pro Leu Trp Gly Gly Pro  
50 55 60

Leu His Tyr Pro Thr Tyr Arg Pro Phe Pro His Pro Pro Pro His Ser  
65 70 75 80

Pro Pro Pro Gly Cys Asp Cys Cys Lys Met Gly Val  
85 90

<210> 21  
<211> 2644  
<212> DNA  
<213> Homo sapiens

<400> 21  
gttgaggatg gctgacattc tctctcagtc agagaccctg gctcgcgaag acctcagtg 60  
ggaattcaag aagccagctc tgcgggtgtc cccagggcg cggagtaagg ccccgccag 120  
cagttcttca aacctgaggg aggtacagaa ggaaggccce actgcgttgc aggaatccaa 180  
ttctgaggag cccgacatcc ctctctctca gccggactgc ggtgatttta ggagtctaca 240  
ggaggagcag tcgggcctca cggcagcggg ttcttccctt ggcggtccag cccgggctcc 300  
ccctaccacg gagcctccat ggggtggccc tgcacagcc cctacagct tagagacct 360  
gaaggggcgc actatecttg gccccgtag cttagaaggg acgagttact gctttttgg 420  
gaggtgtgtc ggtcggaag tgtgcttga gcccccttc gctctcggg accacgcagt 480  
gctgcagcac agggcgtccg gccctgacgg agaattccgc agcaacgggc cgggcttcta 540  
cctctacgat ctgggaagca cccatggcac tttctcaac aaactcgca tcccacctc 600  
ccctactgt cagttccag ttgggcattg tgttcgctt ggaggcagca cccggctctt 660  
tatcctgcag ggaacaggg aagaccggga ggcagaatcc gagttaacag taacacagtt 720  
gaaggaaatg cgaagcagc agcaaatatt gttggrgaag aagatgctag gagaagactc 780  
agatgaagaa gaggaaatgg atacctctga aaggaagata aatgctgga gccagatga 840  
tgagatgggt tgcacctggg gaatgggaga agatgcagta gaggatgatg ctgaagaga 900  
ccctattgtc tttagatttc agcaggaaag gtaggccttt tatataagg atcccaaaa 960  
ggctctccaa ggcttttttg accggaagg agaagaatta gaatatgaat ttgatgaac 1020  
gggacatagc acttggtctt gcagggtgag attacctgtg gacgattcaa ctggaaaaca 1080  
actggtgggt gaggccattc actcaggaaa gaaaaaagaa gcaatgatcc agtgcctatt 1140  
ggaagcttgt cggaktcttg acacttttgg abtgcctcgg caggaaagcag tatctggga 1200  
aaggaaagcc aagaactggg aagatgaaga cttttatgat agtgatgatg acacatttct 1260  
tgataggagt cctctgattt aggaagagcc cctgaacaga atgaagaagg ctggcagat 1320  
tgatggaag ccagagacct ttgatcatt ggttgcaaa ttaantgatg ctgaaggaa 1380  
actttctgaa atctctgaga gattgaagc ctcaagccaa gttctatcag agtctccatc 1440  
tcaggattct ttgatgctt tcatgtcaga aatgaatca ggcagtacat tagatgggt 1500  
gtcccggaag aactctcacc tgagaacttt tgaactgagg aaagaacaa acagacttaa 1560  
agggttaata aaattgttaa agccagcaga gattccagaa ctaaaanga ctgaactcca 1620  
gactacaggt gcagaaacca aagctaaaaa gcttacattg cctctatttg gtgccatgaa 1680  
aggaggaagc aacttcaaat taaaacttg aacagtaggg aagttacccc ccaagcgtcc 1740  
agaactccct ccaactctaa tgagaatgaa agatgagcct gaagtgaagc agggaggagg 1800  
aggaggaagc gaagaagaga aaagaaagga ggcagatgaa aagaaaaac tggaggatgg 1860  
aagctcagt aggcacagc cagagataga gccagagca gcagtgcagg aatggaggcc 1920  
tcccacagat ctccacatt tlaaggaac ccaaacccat ggtaatatct ttcttctct 1980  
tctgtgttg tkeagtggg agttacattg attgtggta ggtttttaa agcagggcca 2040  
gttcttctct ctctcatttg cttgtatgt gatatactga ctctgtagca aggaacata 2100  
ctttcttggg ctctctctct tgaccggcag ttattatttg tottctatgc aattaaggc 2160  
cagttatttc caatccattc cagaattcca gaaattgaa gggctatgga atttgaacc 2220  
atagctgctg tggataatc ctgagctgct gccactgtgt gacttggagg cagtggaac 2280  
agggtacatg atggggcctg atcagggtgt ctcgggata gtcacccct attcattttt 2340  
tctccatcc cttaaacaga ggcacaaaca taattgtact cattggacta aagttctca 2400  
gaaggatctt gcttcaattc tttttgtgt tttggaaact agcacaaac ctgacacata 2460  
tcccccggc tcagcaata ttgatgaa aatgttgaac gacggatag attgtcttc 2520  
atatagatct atgcatcat taattctgt atctctatat atctattcta attacaagg 2580  
gttatgtgtt cattttagaa actatagatc atacataaaa gtccaaaggc aaaaaaaa 2640  
aaa 2644

<210> 22

<211> 667  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (250)

<400> 22

```
Met Ala Asp Ile Leu Ser Gln Ser Glu Thr Leu Ala Ser Gln Asp Leu
  1                      5                      10                      15

Ser Gly Asp Phe Lys Lys Pro Ala Leu Pro Val Ser Pro Ala Ala Arg
      20                      25                      30

Ser Lys Ala Pro Ala Ser Ser Ser Ser Asn Pro Glu Glu Val Gln Lys
      35                      40                      45

Glu Gly Pro Thr Ala Leu Gln Asp Ser Asn Ser Gly Glu Pro Asp Ile
      50                      55                      60

Pro Pro Pro Gln Pro Asp Cys Gly Asp Phe Arg Ser Leu Gln Glu Glu
      65                      70                      75                      80

Gln Ser Arg Pro Thr Thr Ala Val Ser Ser Pro Gly Gly Pro Ala Arg
      85                      90                      95

Ala Pro Pro Tyr Gln Glu Pro Pro Trp Gly Gly Pro Ala Thr Ala Pro
      100                     105                     110

Tyr Ser Leu Glu Thr Leu Lys Gly Gly Thr Ile Leu Gly Thr Arg Ser
      115                     120                     125

Leu Lys Gly Thr Ser Tyr Cys Leu Phe Gly Arg Leu Ser Gly Lys Asp
      130                     135                     140

Val Cys Leu Glu His Pro Ser Val Ser Arg Tyr His Ala Val Leu Gln
      145                     150                     155                     160

His Arg Ala Ser Gly Pro Asp Gly Glu Cys Asp Ser Asn Gly Pro Gly
      165                     170                     175

Phe Tyr Leu Tyr Asp Leu Gly Ser Thr His Gly Thr Phe Leu Asn Lys
      180                     185                     190

Thr Arg Ile Pro Pro Arg Thr Tyr Cys Arg Val His Val Gly His Val
      195                     200                     205

Val Arg Phe Gly Gly Ser Thr Arg Leu Phe Ile Leu Gln Gly Pro Glu
      210                     215                     220

Glu Asp Arg Glu Ala Glu Ser Glu Leu Thr Val Thr Gln Leu Lys Glu
      225                     230                     235                     240

Leu Arg Lys Gln Gln Gln Ile Leu Leu Xaa Lys Lys Met Leu Gly Glu
      245                     250                     255

Asp Ser Asp Glu Glu Glu Glu Met Asp Thr Ser Glu Arg Lys Ile Asn
      260                     265                     270
```



Ala Gly Ser Gln Asp Asp Glu Met Gly Cys Thr Trp Gly Met Gly Glu  
 275 280 285  
 Asp Ala Val Glu Asp Asp Ala Glu Glu Asn Pro Ile Val Leu Glu Phe  
 290 295 300  
 Gln Gln Glu Arg Glu Ala Phe Tyr Ile Lys Asp Pro Lys Lys Ala Leu  
 305 310 315 320  
 Gln Gly Phe Phe Asp Arg Glu Gly Glu Glu Leu Glu Tyr Glu Phe Asp  
 325 330 335  
 Glu Gln Gly His Ser Thr Trp Leu Cys Arg Val Arg Leu Pro Val Asp  
 340 345 350  
 Asp Ser Thr Gly Lys Gln Leu Val Ala Glu Ala Ile His Ser Gly Lys  
 355 360 365  
 Lys Lys Glu Ala Met Ile Gln Cys Ser Leu Glu Ala Cys Arg Ile Leu  
 370 375 380  
 Asp Thr Leu Gly Leu Leu Arg Gln Glu Ala Val Ser Arg Lys Arg Lys  
 385 390 395 400  
 Ala Lys Asn Trp Glu Asp Glu Asp Phe Tyr Asp Ser Asp Asp Asp Thr  
 405 410 415  
 Phe Leu Asp Arg Thr Gly Leu Ile Glu Lys Lys Arg Leu Asn Arg Met  
 420 425 430  
 Lys Lys Ala Gly Lys Ile Asp Glu Lys Pro Glu Thr Phe Glu Ser Leu  
 435 440 445  
 Val Ala Lys Leu Asn Asp Ala Glu Arg Glu Leu Ser Glu Ile Ser Glu  
 450 455 460  
 Arg Leu Lys Ala Ser Ser Gln Val Leu Ser Glu Ser Pro Ser Gln Asp  
 465 470 475 480  
 Ser Leu Asp Ala Phe Met Ser Glu Met Lys Ser Gly Ser Thr Leu Asp  
 485 490 495  
 Gly Val Ser Arg Lys Lys Leu His Leu Arg Thr Phe Glu Leu Arg Lys  
 500 505 510  
 Glu Gln Gln Arg Leu Lys Gly Leu Ile Lys Ile Val Lys Pro Ala Glu  
 515 520 525  
 Ile Pro Glu Leu Lys Lys Thr Glu Thr Gln Thr Thr Gly Ala Glu Asn  
 530 535 540  
 Lys Ala Lys Lys Leu Thr Leu Pro Leu Phe Gly Ala Met Lys Gly Gly  
 545 550 555 560  
 Ser Lys Phe Lys Leu Lys Thr Gly Thr Val Gly Lys Leu Pro Pro Lys  
 565 570 575  
 Arg Pro Glu Leu Pro Pro Thr Leu Met Arg Met Lys Asp Glu Pro Glu  
 580 585 590

Val Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu Lys Glu Lys Glu  
 595 600 605

Glu His Glu Lys Lys Lys Leu Glu Asp Gly Ser Leu Ser Arg Pro Gln  
 610 615 620

Pro Glu Ile Glu Pro Glu Ala Ala Val Gln Glu Met Arg Pro Pro Thr  
 625 630 635 640

Asp Leu Thr His Phe Lys Glu Thr Gln Thr His Gly Asn Ile Phe Leu  
 645 650 655

Leu Leu Pro Val Leu Phe Ser Gly Gln Leu His  
 660 665

&lt;210&gt; 23

&lt;211&gt; 2402

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

```

gagcgagag accagaggag cggccggcggc tgcagagctg cagcgaggga tctcttcgag 60
ctgtctgtgt cgggscagcc agcgcgcaac tgagccagag gacagcgcat cctttcgagg 120
cgggscggca gggccctctg ggtcggaag ctggtctccc ggttggcac cgggacccac 180
gagcccaatg cggggggcgg cggcaaatc ccaacactg tagagatcac ccccaactcc 240
aacggacagg tggggacct cggagatgag gtgcacagg agcagctgca ggttgagcag 300
gagcgcgagc gggaggggga gggagcgcg gscggcgacg gactgggcag cagctctgtg 360
ctggcgctgc ccccgaggcc cctcagcttt gaggcgctg tggccagggt gggggcgctg 420
ggcgcgcgcc agcagctgca gctcgccctc tgcgctctg cgggtgctct cgtggctctg 480
ggcattggct cggactccat ctccagctg tgcacccagc tgcattgcca ctacggggcc 540
ttccccccta atgcctctgg ctgggagcag cctcccaatg ccaggggcgt cagcgctgcc 600
agcgctgccc tagcagctag cggcgccagc cgtctgccc ccagtaccga cccctcgctg 660
agcggtctcg ccccgccggg ctccaactat tgcctcaagg atctgggacta taatggcctt 720
cctgtgtctc ccaccaactg cctcgccag tgggactctg tgtgtgacct ggggtggcag 780
gtgactctgg agcagatctt cttcatcttg ggttttgctt cgggtacct gtccctgggt 840
taccggcgag acagatttgg cgtcgcggg attgtgctg tgaccttgag cctgggtggg 900
ccctgtggag tagggggggc tctcgaggc tctccacag gctcattgga cctccgaltc 960
ctcttggggt ttctgcttgc cgtgttgac ctgggtgtct acctgatgag cctgggagctg 1020
tgggacccaa cccagagggt cgggttgccc ctggcagggg ggttggtggg ggtgggaggg 1080
cacttctgtt tcttgggcct gggccttgct tctaaaggat ggcgattcct acagcgaaatg 1140
atcaccgctc cctgcactct ctctctttt tatggctagg ctggtttgtt cctgggagctc 1200
gcacgggtgg tgatagtga gggacagatt gaggaggttc agtgtgtct ggggatccctg 1260
gctgagcgaa accggcccca tgggcagatg ctggggggag agggccaggg ggccttgacg 1320
gacctggaga atacctgcc tctcctgca acatctctct ttctcttgc tctctctc 1380
aactaccgca acctctggga aatctgctt atcttgggt tcccaactt cattgccc 1440
gcattctgcc actgctacca gctgtggga ggaaggagg ggcctcggg ctctcactg 1500
tgctctctgc tggccagcgg caccgcagcc ctggcctgtg tctctctggg ggtcacctg 1560
gacgagattg ggcgcgggg cctccttctt ctctccatga ccttaccagg cattgcttc 1620
ctggtctctg tgggctctg ggtatctctg aacgaggtct cctcaccac ctctctctg 1680
cttgggctct tctctctcct agctgcccgc atctcagca cctccttgc tgcgagctc 1740
atctccacca ctgtccgggg cgttggcctg ggcctgata tggctctagg ggcgcttggg 1800
ggactgagcg gcccgggcca ggcctccac atggggccat ggcctctct gacgacgtg 1860
gtgttggggg cctgcgctct cctctgctt ctgagctta tgcgtctgc gtcgcccggc 1920
cgcaagctcc cggccagggt gctcgggac ggggagctgt gtcgcccggc ttctctgctg 1980
cggcagccac cctctaccgg ctgtgaccac gtcccgctgc ttgcaacccc caacctgctc 2040
ctctgagcgg cctctgagta ccttgggggg aggcctggcc acacagagag gtggcagaa 2100
gacggggag actgagtagg gaaggcaggg ctgcccagaa gttcagagg cactcacgc 2160
cagccatgca gtagagctca gagggcgtc cccacccctg ctctccctg ctgctttgca 2220
ttcacttctt tggccagagt caggggacag ggagagagct ccaactgta accactgggt 2280

```

```

ctggggtcca tctggggccc aaagacatcc acccagacct cattattttct tgcctctatca 2340
ctctgtttca ataaagacat ttgggtatca cgaataaaca aaaaataaaa aaaaataaaa 2400
aa                                         2402

```

&lt;210&gt; 24

&lt;211&gt; 520

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 24

```

Met Ala Ser Asp Pro Ile Phe Thr Leu Ala Pro Pro Leu His Cys His
  1              5              10              15

Tyr Gly Ala Phe Pro Pro Asn Ala Ser Gly Trp Glu Gln Pro Pro Asn
          20              25              30

Ala Ser Gly Val Ser Val Ala Ser Ala Ala Leu Ala Ala Ser Ala Ala
          35              40              45

Ser Arg Val Ala Thr Ser Thr Asp Pro Ser Cys Ser Gly Phe Ala Pro
          50              55              60

Pro Asp Phe Asn His Cys Leu Lys Asp Trp Asp Tyr Asn Gly Leu Pro
          65              70              75              80

Val Leu Thr Thr Asn Ala Ile Gly Gln Trp Asp Leu Val Cys Asp Leu
          85              90              95

Gly Trp Gln Val Ile Leu Glu Gln Ile Leu Phe Ile Leu Gly Phe Ala
          100              105              110

Ser Gly Tyr Leu Phe Leu Gly Tyr Pro Ala Asp Arg Phe Gly Arg Arg
          115              120              125

Gly Ile Val Leu Leu Thr Leu Gly Leu Val Gly Pro Cys Gly Val Gly
          130              135              140

Gly Ala Ala Ala Gly Ser Ser Thr Gly Val Met Ala Leu Arg Phe Leu
          145              150              155              160

Leu Gly Phe Leu Leu Ala Gly Val Asp Leu Gly Val Tyr Leu Met Arg
          165              170              175

Leu Glu Leu Cys Asp Pro Thr Gln Arg Leu Arg Val Ala Leu Ala Gly
          180              185              190

Glu Leu Val Gly Val Gly Gly His Phe Leu Phe Leu Gly Leu Ala Leu
          195              200              205

Val Ser Lys Asp Trp Arg Phe Leu Gln Arg Met Ile Thr Ala Pro Cys
          210              215              220

Ile Leu Phe Leu Phe Tyr Gly Trp Pro Gly Leu Phe Leu Glu Ser Ala
          225              230              235              240

Arg Trp Leu Ile Val Lys Arg Gln Ile Glu Glu Ala Gln Ser Val Leu
          245              250              255

Arg Ile Leu Ala Glu Arg Asn Arg Pro His Gly Gln Met Leu Gly Glu
          260              265              270

```

Glu Ala Gln Glu Ala Leu Gln Asp Leu Glu Asn Thr Cys Pro Leu Pro  
 275 280 285  
 Ala Thr Ser Ser Phe Ser Phe Ala Ser Leu Leu Asn Tyr Arg Asn Ile  
 290 295 300  
 Trp Lys Asn Leu Leu Ile Leu Gly Phe Thr Asn Phe Ile Ala His Ala  
 305 310 315 320  
 Ile Arg His Cys Tyr Gln Pro Val Gly Gly Gly Gly Ser Pro Ser Asp  
 325 330 335  
 Phe Tyr Leu Cys Ser Leu Leu Ala Ser Gly Thr Ala Ala Leu Ala Cys  
 340 345 350  
 Val Phe Leu Gly Val Thr Val Asp Arg Phe Gly Arg Arg Gly Ile Leu  
 355 360 365  
 Leu Leu Ser Met Thr Leu Thr Gly Ile Ala Ser Leu Val Leu Leu Gly  
 370 375 380  
 Leu Trp Asp Tyr Leu Asn Glu Ala Ala Ile Thr Thr Phe Ser Val Leu  
 385 390 395 400  
 Gly Leu Phe Ser Ser Gln Ala Ala Ala Ile Leu Ser Thr Leu Leu Ala  
 405 410 415  
 Ala Glu Val Ile Pro Thr Thr Val Arg Gly Arg Gly Leu Gly Leu Ile  
 420 425 430  
 Met Ala Leu Gly Ala Leu Gly Gly Leu Ser Gly Pro Ala Gln Arg Leu  
 435 440 445  
 His Met Gly His Gly Ala Phe Leu Gln His Val Val Leu Ala Ala Cys  
 450 455 460  
 Ala Leu Leu Cys Ile Leu Ser Ile Met Leu Leu Pro Glu Thr Lys Arg  
 465 470 475 480  
 Lys Leu Leu Pro Glu Val Leu Arg Asp Gly Glu Leu Cys Arg Arg Pro  
 485 490 495  
 Ser Leu Leu Arg Gln Pro Pro Pro Thr Arg Cys Asp His Val Pro Leu  
 500 505 510  
 Leu Ala Thr Pro Asn Pro Ala Leu  
 515 520

<210> 25  
 <211> 2377  
 <212> DNA  
 <213> Homo sapiens

<400> 25  
 ttcatctctc agcgggaatc catcagttga aatagttcat ggtattatgc acctatataa 60  
 gacaaataag atgacctcct caaagaaga tctgaggcgc agtgcctatgc tgtgtattct 120  
 cacagtcctt gctgcaatga ccaattatga ccttatgaag ttgtttgccc catthaaaga 180  
 agtaattgaa caaatgaaa ttatcagaga ctctaactccc aaccaatata tgggtcctgat 240

```

aasgttttccgt gcaacaggctg atgcgggtag tttttatatg acctgcaatg gccggccagtt 300
caactcaata gaagatgacg tttgccagct agtqtatcgt gaaagaggctg aagtgcctaa 360
atctgaagat ggagccagcc tcccagtgat ggacctgact gaactcccc aagtgcacgg 420
gtgtctggag ccgatggacg agtctgtgaa tgggcattctt acsaagttat gtaaccacag 480
cttccacagc cagtgtctac agcgctggga cgataccacg tgtcttgttt gccgggtactg 540
tcaaacggcc gagccagtag aagaaaataa gtgttttggg tgtggtgttc aggaatatct 600
ttggattttt ltaatatgag gccacatagg atgtggacgg tatgtcagtc gaactgttta 660
taagcacttt gaggaacgct agcaacagta tggcaatgag cttaaccaac atcaggtctg 720
ggactatgct ggagataact atgttccatc actggttgca agtaaaacag atggaasat 780
agtacagtat gaatgtgag gggatacttg ccaggaaag agaatagatg ccttacagtt 840
agagtattca tttttctaa caagccagct ggaatctcag cgaactctac gggaaaaaca 900
gatagttcgg atagagaag agacagcag ggaactaac aacatgaaga ccaagtttan 960
agaaacactt gagaagtgtg ataactcaga gcaaaaacta aatgatctcc taaaagaaa 1020
gcactctgtg gaagaaaagt gaactcagct aaacacaaa gtggccaaac tcaatctca 1080
cagtgggkat cctagctct agcaagctg agtggggaga ttctcctcc gtgtgaaat 1140
gtcagtgag gctctgact agctaattgt gtattttgtt gggtttagta tttctaat 1200
gtttacaaa tttgggctg catgttcagg ttgcagctag agggagctcg ggcagatttt 1260
caattacgct ttcaagat aaccaaagc ttttctana tctaaaaat agattttca 1320
cagagccccc tttagaacg tcatataacg cttgtgtcgg ccaacagagg ggtgtgtac 1380
tctctctgga accataaatg tcaataaatt tateacctgc agtaattgag caactttaa 1440
ataagccctg tgttggaaat tagtttcttg aagaggtaga gggataggtt agtaagatgt 1500
attgttaaac aacagpttt agtttttgc ttaataatag ccacagtttt tcaaatgac 1560
acattkaga ataggtttt agcctgtcat taggcctcat cccctttgac ctasatgtct 1620
gacatgttac ttgttagtac atcaactgta tcaataatc scatctgttt ttgtgggatg 1680
tgcctcagca tttcccaaaa aacttkacgt gtaatgttgc aasatgaatg tactcagaca 1740
ttcttaattt ttacttggg ccgaccaact ctltgagtet ctcttggact tatatctaca 1800
gatactttaa gagtgggaat gtaagcata acctaatct ctltccata gagattctat 1860
tttattttaa atctatttt acactagtta gaactcgtct gttttggcca agtaattgtc 1920
ttgcatgtct gaccttgag aagctgggtt ggtatnagc atactaatga agagaattag 1980
aagtagttta caagctcgc tcaactctca tttctctgt gtacaggtcg gataaattgt 2040
ccaccaccac ctgggaaac agatttttca gtaccaggtg tamttggagg tttaaaaaa 2100
gtgccagag caatgagca atagggcagt gtttccaaac aagataagta aagaaaaaa 2160
atatgtccca gaaaaaaa aatcttacc aagataagta aagaaaaaa aattttttt 2220
taaacagtc aagagtcatg ttgaatttc aaaaaatc atcagacaga agttgttttc 2280
ttcaggaggg aatgaacca cttaatatc ccaactcacc ttgaacaatg aattgaatt 2340
aaatagcca aactttgaaa ttaaaaaaa aaaaaaa 2377

```

&lt;210&gt; 26

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

```

Met His Leu Tyr Lys Thr Asn Lys Met Thr Ser Leu Lys Glu Asp Val
  1             5             10             15

```

```

Arg Arg Ser Ala Met Leu Cys Ile Leu Thr Val Pro Ala Ala Met Thr
  20             25             30

```

```

Ser His Asp Leu Met Lys Phe Val Ala Pro Phe Asn Glu Val Ile Glu
  35             40             45

```

```

Gln Met Lys Ile Ile Arg Asp Ser Thr Pro Asn Gln Tyr Met Val Leu
  50             55             60

```

```

Ile Lys Phe Arg Ala Gln Ala Asp Ala Asp Ser Phe Tyr Met Thr Cys
  65             70             75             80

```

```

Asn Gly Arg Gln Phe Asn Ser Ile Glu Asp Asp Val Cys Gln Leu Val
  85             90             95

```

Tyr Val Glu Arg Ala Glu Val Leu Lys Ser Glu Asp Gly Ala Ser Leu  
 100 105 110  
 Pro Val Met Asp Leu Thr Glu Leu Pro Lys Cys Thr Val Cys Leu Glu  
 115 120 125  
 Arg Met Asp Glu Ser Val Asn Gly Ile Leu Thr Thr Leu Cys Asn His  
 130 135 140  
 Ser Phe His Ser Gln Cys Leu Gln Arg Trp Asp Asp Thr Thr Cys Pro  
 145 150 155 160  
 Val Cys Arg Tyr Cys Gln Thr Pro Glu Pro Val Glu Glu Asn Lys Cys  
 165 170 175  
 Phe Glu Cys Gly Val Gln Glu Asn Leu Trp Ile Cys Leu Ile Cys Gly  
 180 185 190  
 His Ile Gly Cys Gly Arg Tyr Val Ser Arg His Ala Tyr Lys His Phe  
 195 200 205  
 Glu Glu Thr Gln His Thr Tyr Ala Met Gln Leu Thr Asn His Arg Val  
 210 215 220  
 Trp Asp Tyr Ala Gly Asp Asn Tyr Val His Arg Leu Val Ala Ser Lys  
 225 230 235 240  
 Thr Asp Gly Lys Ile Val Gln Tyr Glu Cys Glu Gly Asp Thr Cys Gln  
 245 250 255  
 Glu Glu Lys Ile Asp Ala Leu Gln Leu Glu Tyr Ser Tyr Leu Leu Thr  
 260 265 270  
 Ser Gln Leu Glu Ser Gln Arg Ile Tyr Trp Glu Asn Lys Ile Val Arg  
 275 280 285  
 Ile Glu Lys Asp Thr Ala Glu Glu Ile Asn Asn Met Lys Thr Lys Phe  
 290 295 300  
 Lys Glu Thr Ile Glu Lys Cys Asp Asn Leu Glu His Lys Leu Asn Asp  
 305 310 315 320  
 Leu Leu Lys Glu Lys Gln Ser Val Glu Arg Lys Cys Thr Gln Leu Asn  
 325 330 335  
 Thr Lys Val Ala Lys Leu Lys Ser Gln Ser Gly Tyr Pro Ser Ile  
 340 345 350

&lt;210&gt; 27

&lt;211&gt; 460

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

ccgagatgaag ccggcggttg acgagatggt ccccgaggcc gccgggacct acgtggacct 60  
 ggacgaggcg ggagggcgca cggcgctctt gatggacttg gcggccaatg aaagacggtt 120  
 catgcagact tctttacga ttttgaagat cttttgatg atgatgacct ccagtgagat 180  
 gccctcttgc tgcaggcggg gccaaacct tggtaacag cgcagtggtg agcctggcca 240

```

ggacagtttc aggtgggtttt aaagaacacg tggaaatccc ttgsattttag gacctggtta 300
accagaaaga taagactgtt cttaacgacg tgaatgattc tgttcacoto tgaacgggat 360
caggtttttgt cctcactcca attaaagaa agcaatgtca catgassana aaaaaaanaa 420
aaaaaaaaaa aaaaaaanaa aaaaaaanaa aaaaaaanaa 460

```

<210> 28  
 <211> 85  
 <212> PRT  
 <213> Homo sapiens

```

<400> 28
Met Lys Pro Ala Val Asp Glu Met Phe Pro Glu Gly Ala Gly Pro Tyr
 1             5             10             15
Val Asp Leu Asp Glu Ala Gly Gly Ser Thr Gly Leu Leu Met Asp Leu
          20          25          30
Ala Ala Asp Glu Lys Pro Phe Met Gln Thr Phe Leu Thr Ile Leu Lys
      35          40          45
Ile Phe Leu Met Met Met Thr Ser Ser Glu Met Pro Ser Gly Cys Arg
      50          55          60
Arg Gly Gln Ala Leu Gly Thr Glu Pro Gln Cys Glu Pro Ala Gln Asp
      65          70          75          80
Ser Phe Arg Trp Phe
          85

```

<210> 29  
 <211> 3204  
 <212> DNA  
 <213> Homo sapiens

```

<400> 29
gtttggcacc tctggccgag tctctgttgc cgggtgcatg ttggagcgga gacttagcac 60
aatggcagaa cctgtttctc cactgaagca ctctgtgtg gctaagaggg cgattactgc 120
aatctttgac cagttactgg agtttgttac tgaaggatca cattttgttg aagcaacata 180
taagaatccg gaacttgatc gaatagccac tgaagatgat ctggttagaa tgcgaaggata 240
taagacaaag cttcccatca ttggtgaggg gctatctcgg agacacatga aggtggcatt 300
ttttggcagg scasgcagtg ggaagagctc tgttatcaat gcaatgttgtt gggataaagt 360
ctctcctagt gggattggcc ctacaaccaa ttgcttctca agtgttgaag gaactgatgg 420
agataaagcc tatcttatga cagaaggatc agatgaaaaa aagagtgtga agacagttaa 480
tcactggcc catgccttcc acatggacaa agatttgaaa gctggctgtc ttgtacgtgt 540
gttttgacca aaagcaaat gtgcctctct gagagatgac ctggtgttag tagacagfcc 600
agggccagat gtcactacag agctggatag ctggattgat aagttttgac tagatgttga 660
tgtctttgtt ttggtcgcaa actctgaatc aacactaatg aatacggaaa aacacttttt 720
tcacaaggty aatgaagggc ttcccaagcc taatatcttc attctcaata atcgttggga 780
tgctcttgc tcagagccag aatatatgga agacgtacgc agacagcaca tggaaagatg 840
cctgcatttc ttggtggagg agctcaaatg tgtcaaatgt ttagaagcac agaatcgtat 900
cttctttgtt tcagcaagg aagttcttag tcttagaaag caaaaagcac aggggatgcc 960
agaaagtgt gtggcaattg ctgaaggatt tcatgcaaga ttacaggcat ttcaaatctt 1020
tgaaacaatc tttagggagt gtatctcgca gtcagtagtg aaaaacaaat tggaaacaga 1080
cactatcaga gctaaacaga tactagctac tgtgaaaaac ctatgggatt cagtaaacct 1140
ggcagctgaa gataaaagge attactcagt ggaagagagg gaagaccaca ttgatatagt 1200
ggactttatt cgaaacacaga tgaactcttt aacactggat gttaagaaaa aatcaaggga 1260
ggttacggag gaggtagcaa acaaagtctc atgtgcantg scagatgaaa ttgttcgact 1320
gtctgttttg gttgatgaat ttgttcaga gtttcatct aatccagatg kattaanaat 1380
atataaaagt gaactaata agcacataga ggtgtgtatg ggaagaaatt tggctgatcg 1440

```

```

atgcacacgat gaagtaaaag ccttagtgcc tcagaccag caagaaakta ttgaaaattt 1500
gaagcattta cctcagctg gtatacagga taactacat acactgctc cttccagaa 1560
attctgacetc agttataatc taacttaaca caagttatgt tcagattttc aaagagatat 1620
tgtatttctg ttttcccttg gctggtcttc ccttgtagat cgtatttttg gctcagaaa 1680
tgctcaaaag gtgctctag gattatacag gctatcttt cagctcccta gatcttttag 1740
ttctactccc actgctctca ccactccagc aacgcacagc aatgcctcac aggaagaact 1800
catgattaca ttagtaacag gattggcgct cgttacatct agaacttcta tgggcatcct 1860
tattgttgga ggaagtattt ggaaaactat aggttggaac ctctatcttg tttcattaac 1920
tatgtatgga gctttgtatc ttatgaaag actgagctgg accaccatg ccaaggagcg 1980
agccttttaa cagcagtttg taactatgc aactgaaaaa ctgaggatga ttgttagctc 2040
cagcagtgca aactgcagtc actaagtaaa acnacsaaat gctaccactt ttgtctgctc 2100
gtgcacacaa gttgatatta ctcaaaaaaa gcttggaaga gaacttgcta gattacccaa 2160
agaaatagat cagttggaga aaatacaaaa caattcaaaag ctcttaagaa ataaagctgt 2220
tcacgttcaa aatgagctgg agaattttac taagcagttt ctacttcaa gcaatgaaga 2280
ctcttaaaaa tagagattgc ttgtgtgccc atgacagag gaaacgaaac ttgtaagatt 2340
ggacaggttg ttattttat gaaattactt taatatgaa ttgtactaac tgtacctaaa 2400
tagcaaaagc ctgtgtagat tctgttaatg atctgtctca ggttatgtgt tttttgaa 2460
agtgttatgt ccttagtttt aattttgagt aaagaaaagg ctcaatcatg aattagttac 2520
aagcaacagt accaacttat gtgacccctg aggtgtgggg ctgtgagctc ttaatttgtt 2580
tttgattctg aaaaactctg ctctctggca tccaggagtt agagattgag ccttctatct 2640
tctttctcaa aactagtttt tgatgcttcc ttctatggga atagtcactt tttatttag 2700
taaatggcat tgttggaaac accaaggagt gtggaatgtc cttagagtga ttatttatgc 2760
aagttacagt cagcttgcca tcatggcagc tatgtgaac actaataaat gtgtttttac 2820
tttttatccc cgttaaaact gatgtaaaac aggtataaag ctgtttatag tcaattata 2880
gtatctgggt ctaagtaatt tcttagatg tttctaaaga aacattttca gctttgtccc 2940
cattatgatt ccaataagga accgtttctt agtgcattt taggagtaaa gtttgaagag 3000
ataaaaaatg ccaagatag ggaagctctg aattttgaat gataaacagt gatgttttaa 3060
aaaagctgtt gttcttcagg aggcatttgc ctaggatatt gctgagattat accccattgg 3120
aggcttttaa tttatttgt atgaatttct caggatttca ttaaaaatta ttattgtatt 3180
tttaacctta aaaaaaaaaa aaaa

```

&lt;210&gt; 30

&lt;211&gt; 741

&lt;212&gt; PAT

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

```

Met Ala Glu Pro Val Ser Pro Leu Lys His Phe Val Leu Ala Lys Lys
1 5 10 15

```

```

Ala Ile Thr Ala Ile Phe Asp Gln Leu Leu Glu Phe Val Thr Glu Gly
20 25 30

```

```

Ser His Phe Val Glu Ala Thr Tyr Lys Asn Pro Glu Leu Asp Arg Ile
35 40 45

```

```

Ala Thr Glu Asp Asp Leu Val Glu Met Gln Gly Tyr Lys Asp Lys Leu
50 55 60

```

```

Ser Ile Ile Gly Glu Val Leu Ser Arg Arg His Met Lys Val Ala Phe
65 70 75 80

```

```

Phe Gly Arg Thr Ser Ser Gly Lys Ser Ser Val Ile Asn Ala Met Leu
85 90 95

```

```

Tyr Asp Lys Val Leu Pro Ser Gly Ile Gly His Ile Thr Asn Cys Phe
100 105 110

```

```

Leu Ser Val Glu Gly Thr Asp Gly Asp Lys Ala Tyr Leu Met Thr Glu
115 120 125

```



Gly Ser Asp Glu Lys Lys Ser Val Lys Thr Val Asn Gln Leu Ala His  
 130 135 140  
 Ala Leu His Met Asp Lys Asp Leu Lys Ala Gly Cys Leu Val Arg Val  
 145 150 155 160  
 Phe Trp Pro Lys Ala Lys Cys Ala Leu Leu Arg Asp Asp Leu Val Leu  
 165 170 175  
 Val Asp Ser Pro Gly Thr Asp Val Thr Thr Glu Leu Asp Ser Trp Ile  
 180 185 190  
 Asp Lys Phe Cys Leu Asp Ala Asp Val Phe Val Leu Val Ala Asn Ser  
 195 200 205  
 Glu Ser Thr Leu Met Asn Thr Glu Lys His Phe Phe His Lys Val Asn  
 210 215 220  
 Glu Arg Leu Ser Lys Pro Asn Ile Phe Ile Leu Asn Asn Arg Trp Asp  
 225 230 235 240  
 Ala Ser Ala Ser Glu Pro Glu Tyr Met Glu Asp Val Arg Arg Gln His  
 245 250 255  
 Met Glu Arg Cys Leu His Phe Leu Val Glu Glu Leu Lys Val Val Asn  
 260 265 270  
 Ala Leu Glu Ala Gln Asn Arg Ile Phe Phe Val Ser Ala Lys Glu Val  
 275 280 285  
 Leu Ser Ala Arg Lys Gln Lys Ala Gln Gly Met Pro Glu Ser Gly Val  
 290 295 300  
 Ala Leu Ala Glu Gly Phe His Ala Arg Leu Gln Glu Phe Gln Asn Phe  
 305 310 315 320  
 Glu Gln Ile Phe Glu Glu Cys Ile Ser Gln Ser Ala Val Lys Thr Lys  
 325 330 335  
 Phe Glu Gln His Thr Ile Arg Ala Lys Gln Ile Leu Ala Thr Val Lys  
 340 345 350  
 Asn Ile Met Asp Ser Val Asn Leu Ala Ala Glu Asp Lys Arg His Tyr  
 355 360 365  
 Ser Val Glu Glu Arg Glu Asp Gln Ile Asp Arg Leu Asp Phe Ile Arg  
 370 375 380  
 Asn Gln Met Asn Leu Leu Thr Leu Asp Val Lys Lys Lys Ile Lys Glu  
 385 390 395 400  
 Val Thr Glu Glu Val Ala Asn Lys Val Ser Cys Ala Met Thr Asp Glu  
 405 410 415  
 Ile Cys Arg Leu Ser Val Leu Val Asp Glu Phe Cys Ser Glu Phe His  
 420 425 430  
 Pro Asn Pro Asp Val Leu Lys Ile Tyr Lys Ser Glu Leu Asn Lys His  
 435 440 445

Ile Glu Asp Gly Met Gly Arg Asn Leu Ala Asp Arg Cys Thr Asp Glu  
 450 455 460  
 Val Asn Ala Leu Val Pro Gln Thr Gln Gln Glu Ile Ile Glu Asn Leu  
 465 470 475 480  
 Lys Pro Leu Leu Pro Ala Gly Ile Gln Asp Lys Leu His Thr Leu Ile  
 485 490 495  
 Pro Cys Lys Lys Phe Asp Leu Ser Tyr Asn Leu Asn Tyr His Lys Leu  
 500 505 510  
 Cys Ser Asp Phe Gln Glu Asp Ile Val Phe Arg Phe Ser Leu Gly Trp  
 515 520 525  
 Ser Ser Leu Val His Arg Phe Leu Gly Pro Arg Asn Ala Gln Arg Val  
 530 535 540  
 Leu Leu Gly Leu Ser Glu Pro Ile Phe Gln Leu Pro Arg Ser Leu Ala  
 545 550 555 560  
 Ser Thr Pro Thr Ala Pro Thr Thr Pro Ala Thr Pro Asp Asn Ala Ser  
 565 570 575  
 Gln Glu Gln Leu Met Ile Thr Leu Val Thr Gly Leu Ala Ser Val Thr  
 580 585 590  
 Ser Arg Thr Ser Met Gly Ile Ile Ile Val Gly Gly Val Ile Trp Lys  
 595 600 605  
 Thr Ile Gly Trp Lys Leu Leu Ser Val Ser Leu Thr Met Tyr Gly Ala  
 610 615 620  
 Leu Tyr Leu Tyr Glu Arg Leu Ser Trp Thr Thr His Ala Lys Glu Arg  
 625 630 635 640  
 Ala Phe Lys Gln Gln Phe Val Asn Tyr Ala Thr Glu Lys Leu Arg Met  
 645 650 655  
 Ile Val Ser Ser Thr Ser Ala Asn Cys Ser His Gln Val Lys Gln Gln  
 660 665 670  
 Ile Ala Thr Thr Phe Ala Arg Leu Cys Gln Gln Val Asp Ile Thr Gln  
 675 680 685  
 Lys Gln Leu Glu Glu Glu Ile Ala Arg Leu Pro Lys Glu Ile Asp Gln  
 690 695 700  
 Leu Glu Lys Ile Gln Asn Asn Ser Lys Leu Leu Arg Asn Lys Ala Val  
 705 710 715 720  
 Gln Leu Glu Asn Glu Leu Glu Asn Phe Thr Lys Gln Phe Leu Pro Ser  
 725 730 735  
 Ser Asn Glu Asp Ser  
 740

&lt;210&gt; 31

&lt;211&gt; 2483

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

```

cacatgttgc cccaaataca agcacaacac taaccatgag cttcagcaat cagctcaata 60
cagtgcacaa ccaggccagt gttctagctt ccagttctac tgcagcagct gatactcttt 120
ctctggctea ttccagatgtc tcaactactaa actaccagtc agcttctgtac ccactcatctg 180
ctgcaccagc tcctggagtt gccacgcagg gtgtttctct gcagccttga accacccaga 240
tttgcactca gacagatcca ttccaaacga cttttatagt atgtccacct atgtctgtac 300
ctggactaca agcaacaaca agcattctgt gttccctgtg gaggatggat aatgtctgtac 360
cgattgtacc ccaggcacca gctgtctcag ccactacaga ttcagtcagg agttctcagc 420
cagactltga gggaaaaaat atccagacat tcttgagaaa tggctctctg aggaagctgt 480
acaccactaa tggtagcacc tctccacctc cagtagacca catcacaccg cagtctgcgg 540
tgccctttac tctgagctgc gcagccggcc ggccggcgct ggttgaacag actgcogctg 600
tactgcaggc gtgccclyga gggactcagc aaattctctt gcttccact tggcaacagt 660
tgcttgggt agctctacac aactctgttc agccacagc aatgtctcca gaggccatgg 720
ggagtggaca gcagctagct gactggagga atgcccactc tcatggcaac cagtacagca 780
ctatcatgca gcagccatcc ttgctgacta cccatgtgac attgccact gctcagctc 840
tgantgttgg tgttgcctat gttgtcagac aacacacac cagttccctc ccttcagaag 900
agaantaaga gtacagctca gtctcttcca agtctctctt agatgttctg ccttcccaag 960
tctattctct ggttggagc agtccctcc gcaccacac ttcttataat tcttgggtcc 1020
ctgtcccaag tcagcatcag cccatcatca ttccagatac tccagccct cctgtgagtg 1080
tcactctat ccgaagtga acatgtgagg agagagaca caantacaag cctagtagct 1140
ctggactgag gccaggctc aatgtctcca gttatgtcac tgtcaatgat tctccagact 1200
ctgactcttc tttaggcagc ccttatctca ctgactcctt gagtctctc cggagcaata 1260
gtggatccgt tctggagggg cctggcagag ttgtggcaga tggcactgga accgcacta 1320
tcattgtgac tccactgaaa actcagcttg gtgactgac tgtagcacc caggctcag 1380
gtctcctgag caataagact aagccagtcg cttcagtgag tgggcagtca tctggatgct 1440
gtatccccc caccaggtat cagctccac gcgggggagc cagtgcagca caaccactca 1500
atcttagcca gaaccagag tcctcggcgg cttccacctc acaggagaga agcagcacc 1560
cagcccccag caggcagcag gcttttctgg cccctctctc ccaggcccc tacaccttcc 1620
agcatggcag cccgtacac tcgacagggc acccacact tgcaccggcc cctgtctacc 1680
tgccaaagca ggcctcatctg tctacgtatg ctgccccgac ttctgtgtgt gcactgggt 1740
caaccagctc cattgtctat cttttctccc cccagggttc ctcaaggcat gctgcagct 1800
atccacctca ccttagcact ttgtgtcacc aggtccctgt cagtgttggg ccagcctcc 1860
tcacttctgc cagctgtgac cctgtcagtc accaacccc gtttgccacc caatctctca 1920
ttgggtcttc ccgaggctca accatttacc ctggatcccc gctgagtcct accaagatca 1980
gcagtatcc ctacttatag ttggtgagca tgagggagga ggaatcatgg ctactctctc 2040
ctggccctgc gttcttaata ctggcctatg gagagatcct cttttacct cttgaaattt 2100
cttagccagc aacttktttt gcaggggccc actgaagcag aaggtttttc tctgggggaa 2160
cttgtctcag tgttgactgc attgttgtag tcttccaaa gtttgcctc tttttaatt 2220
cattattttt gtgacagtaa ttttggtact tggaaaggtt cagatgccc tcttctgag 2280
ttaccaagga agagagattg tcttgaggtt accctctgaa aatatatttg tctctctgac 2340
ttgatttcta taatgcttt taaaaacaa tgagccctct ctttatttca tttctgttta 2400
ttgtgattgc tggtcaggaa aatgtctgat agagggtt gaaatctgat gacaaaaaaa 2460
aaaaaaadad aaaaaadad aad

```

&lt;210&gt; 32

&lt;211&gt; 654

&lt;212&gt; PRF

&lt;213&gt; Homo sapiens

&lt;400&gt; 32

```

Met Ser Phe Ser Asn Gln Leu Asn Thr Val His Asn Gln Ala Ser Val
  1                      5                      10                      15

Leu Ala Ser Ser Ser Thr Ala Ala Ala Thr Leu Ser Leu Ala Asn
                20                25                30

```

Ser Asp Val Ser Leu Leu Asn Tyr Gln Ser Ala Leu Tyr Pro Ser Ser  
 35 40 45  
 Ala Ala Pro Val Pro Gly Val Ala Gln Gln Gly Val Ser Leu Gln Pro  
 50 55 60  
 Gly Thr Thr Gln Ile Cys Thr Gln Thr Asp Pro Phe Gln Gln Thr Phe  
 65 70 75 80  
 Ile Val Cys Pro Pro Ala Phe Gln Thr Gly Leu Gln Ala Thr Thr Lys  
 85 90 95  
 His Ser Gly Phe Pro Val Arg Met Asp Asn Ala Val Pro Ile Val Pro  
 100 105 110  
 Gln Ala Pro Ala Ala Gln Pro Thr Thr Asp Ser Val Arg Ser Ser His  
 115 120 125  
 Ala Asp Leu Gln Gly Lys Asn Ile Gln Thr Phe Leu Arg Asn Gly Leu  
 130 135 140  
 Leu Arg Lys Leu Tyr Thr Thr Asn Gly Ser Asn Ser Pro Pro Ser Ser  
 145 150 155 160  
 Ser His Ile Thr Pro Gln Tyr Ala Val Pro Phe Thr Leu Ser Cys Ala  
 165 170 175  
 Ala Gly Arg Pro Ala Leu Val Gln Gln Thr Ala Ala Val Leu Gln Ala  
 180 185 190  
 Trp Pro Gly Gly Thr Gln Gln Ile Leu Leu Pro Ser Thr Trp Gln Gln  
 195 200 205  
 Leu Pro Gly Val Ala Leu His Asn Ser Val Gln Pro Thr Ala Met Ile  
 210 215 220  
 Pro Gln Ala Met Gly Ser Gly Gln Gln Leu Ala Asp Trp Arg Asn Ala  
 225 230 235 240  
 His Ser His Gly Asn Gln Tyr Ser Thr Ile Met Gln Gln Pro Ser Leu  
 245 250 255  
 Leu Thr Asn His Val Thr Leu Ala Thr Ala Gln Pro Leu Asn Val Gly  
 260 265 270  
 Val Ala His Val Val Arg Gln Gln Gln Ser Ser Ser Leu Pro Ser Lys  
 275 280 285  
 Lys Asn Lys Gln Ser Ala Pro Val Ser Ser Lys Ser Ser Leu Asp Val  
 290 295 300  
 Leu Pro Ser Gln Val Tyr Ser Leu Val Gly Ser Ser Pro Leu Arg Thr  
 305 310 315 320  
 Thr Ser Ser Tyr Asn Ser Leu Val Pro Val Gln Asp Gln His Gln Pro  
 325 330 335  
 Ile Ile Ile Pro Asp Thr Pro Ser Pro Pro Val Ser Val Ile Thr Ile  
 340 345 350

Arg Ser Asp Thr Asp Glu Glu Glu Asp Asn Lys Tyr Lys Pro Ser Ser  
 355 360 365  
 Ser Gly Leu Lys Pro Arg Ser Asn Val Ile Ser Tyr Val Thr Val Asn  
 370 375 380  
 Asp Ser Pro Asp Ser Asp Ser Ser Leu Ser Ser Pro Tyr Ser Thr Asp  
 385 390 395 400  
 Thr Leu Ser Ala Leu Arg Gly Asn Ser Gly Ser Val Leu Glu Gly Pro  
 405 410 415  
 Gly Arg Val Val Ala Asp Gly Thr Gly Thr Arg Thr Ile Ile Val Pro  
 420 425 430  
 Pro Leu Lys Thr Gln Leu Gly Asp Cys Thr Val Ala Thr Gln Ala Ser  
 435 440 445  
 Gly Leu Leu Ser Asn Lys Thr Lys Pro Val Ala Ser Val Ser Gly Gln  
 450 455 460  
 Ser Ser Gly Cys Cys Ile Thr Pro Thr Gly Tyr Arg Ala Gln Arg Gly  
 465 470 475 480  
 Gly Thr Ser Ala Ala Gln Pro Leu Asn Leu Ser Gln Asn Gln Gln Ser  
 485 490 495  
 Ser Ala Ala Pro Thr Ser Gln Glu Arg Ser Ser Asn Pro Ala Pro Arg  
 500 505 510  
 Arg Gln Gln Ala Phe Val Ala Pro Leu Ser Gln Ala Pro Tyr Thr Phe  
 515 520 525  
 Gln His Gly Ser Pro Leu His Ser Thr Gly His Pro His Leu Ala Pro  
 530 535 540  
 Ala Pro Ala His Leu Pro Ser Gln Ala His Leu Tyr Thr Tyr Ala Ala  
 545 550 555 560  
 Pro Thr Ser Ala Ala Ala Leu Gly Ser Thr Ser Ser Ile Ala His Leu  
 565 570 575  
 Phe Ser Pro Gln Gly Ser Ser Arg His Ala Ala Ala Tyr Thr Thr His  
 580 585 590  
 Pro Ser Thr Leu Val His Gln Val Pro Val Ser Val Gly Pro Ser Leu  
 595 600 605  
 Leu Thr Ser Ala Ser Val Ala Pro Ala Gln Tyr Gln His Gln Phe Ala  
 610 615 620  
 Thr Gln Ser Tyr Ile Gly Ser Ser Arg Gly Ser Thr Ile Tyr Thr Gly  
 625 630 635 640  
 Tyr Pro Leu Ser Pro Thr Lys Ile Ser Gln Tyr Ser Tyr Leu  
 645 650

&lt;210&gt; 33

&lt;211&gt; 2731

<212> DNA  
<213> Homo sapiens

<220>

<221> Feature

<222> (2173)

<220>

<221> Feature

<222> (2700)

<400> 33

```

ggcactccac ggcgtgtgaag atggcggggg ctggcgtggc tcagggtgtt cctgtcatte 60
ttctgcttct ggcagctcac cctcaccac tctcgttttt cagtccggga ccggcaaccg 120
tagctgctgc cgaacgggtcc aactggcaca ttccgatacc ctccggggaa aattatttta 180
gttttggaaa gctctctctc agaatacca ctatcttctc gaagtctgat ggagaacctt 240
gtgacctgtc ttgaatata acctggtctc tgaasgctc tgattgttac aatgsaattc 300
ataacttcaa ggcgaagaa gtagagttgt atttggaaa actcaaggaa aaagagggtc 360
tgtctgggaa atatcaaaac tcatcaaat tgttcagaa ctgcagtcaa ctcttcaaaa 420
cacagacctt tctggagat ttatgcac gactgcctct tttagggaaa aaacaggagg 480
ctaaggagaa tggacaaaac cttaccttta ttggagataaacggcaatg catgaacct 540
tgcaaacctt gcaagatgaa ccatacatt tcatgtaca tatlggaatt tcatctcaa 600
aggaactctc aaagaaaat tcaatgggtt atctttttac catgactgtt gaagtgaagg 660
gtccctatga atacctcaaa cttgaagact atcccttgat gatttttttc atgtgtatgt 720
gtattgtata tgtctgtttt ggtgtctgtg ggtcggcatg gtctgcctgc tactggagag 780
atctctgag aattcagttt tggatttgtg ctgtcatctt cctcggaaatg cttgagaaag 840
ctgtcttcta tgggaatttt cagaaatctc gacacaaagg agaattctgc caggctgctc 900
tgtctcttgc agagctgctt tcaagcagtg aacgtctcct ggtctgaact ctggtcctca 960
tgcctgctct gggatattgg accctctctt ctttctgtct ctggcatgga aggggtctct 1020
tagtagcagg agcctctctt tcttggcct ttatccctct taatccctct ggccttctca 1080
ggcctcagac tgatcttgtt agctgactc aaacatgaa gctattaaaa ctccggagga 1140
tgtgtgtgtg gatatttatt acctctcttg tcccaaacac gcttcttttg gcagtggcag 1200
acattgttaa actctctttg ttgaaaacca tgaagttcag aatagtgaac tctcagtcgg 1260
catcattgtt gtttatctct tggaaaacca ctgtcttata aacactgat ggtgttgatg 1320
tgagttatca gcaactctat gaataatgta agttatataa ctatataaaa attagagcta 1380
acagtggtaa ggttcttcta ctctctctaa gtattctctc ctatctatgt gcttggatata 1440
ctgggggtct gatactctcc tctgctgtta ttctccagtt tttgggagag aattttgtaa 1500
gtactttttt gttgtttttt taaccagtr catatgtgta aataatacta taggatttct 1560
gtttataacta cagtgtgcta actatgtgta actgtgacag taggatttct 1620
gagtaactaa gagctcaagg gcaaatgaa attatgaat taacctgtta 1680
cttaagatct gataaataat aactttctct ttgcaaat actctgttcc 1740
tattttgttc tgtatagggt atgtgaagc caaaaactac attcatctac 1800
tctcagaggg atgtgaagc actcaagggt ggcctctagt ttatgtgaca aggggttaag 1860
aacagtaact gacgttagca tttatgaaac gaactggggg agctgtgggt 1920
acagtaactg gacgttagca tttatgaaac gaactggggg agctgtgggt 1980
ttttgttgtt tgtattgtag ttgtctctca ctatcagggc gtatcagggc 2040
acttctcccc tgtgttctc cangatctct gtttactctg ctatcagggc 2100
tgtgttctc tgcctctctt ggtctctctt tcttactctc ctatcagggc 2160
agaggttctt ggcctctctt ggtctctctt tcttactctc ctatcagggc 2220
gggtgtgtga ttcgggggaa ctatcagggc ggtctctctt tcttactctc 2280
tgaatcaga agactgggtt tcttactctc ggtctctctt tcttactctc 2340
tgggctctc cactctctc ggtctctctt tcttactctc ctatcagggc 2400
cagcacaact gaagttctt tcttactctc ggtctctctt tcttactctc 2460
ggaaataaatt tgcaggtttt ggtctctctt tcttactctc ctatcagggc 2520
aatattttgt ttttccaggt ggtctctctt tcttactctc ctatcagggc 2580
tgtgtgcaat agcattatgt ctatcagggc ggtctctctt tcttactctc 2640
ccaaaaaaa aaaaaaaa aaaaaaaa a

```

<210> 34

&lt;211&gt; 441

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Mat Ala Ala Ala Ala Trp Leu Gln Val Leu Pro Val Ile Leu Leu Leu
  1              5              10              15

Leu Gly Ala His Pro Ser Pro Leu Ser Phe Phe Ser Ala Gly Pro Ala
          20              25              30

Thr Val Ala Ala Ala Asp Arg Ser Lys Trp His Ile Pro Ile Pro Ser
          35              40              45

Gly Lys Asn Tyr Phe Ser Phe Gly Lys Ile Leu Phe Arg Asn Thr Thr
  50              55              60

Ile Phe Leu Lys Phe Asp Gly Glu Pro Cys Asp Leu Ser Leu Asn Ile
  65              70              75              80

Thr Trp Tyr Leu Lys Ser Ala Asp Cys Tyr Asn Glu Ile Tyr Asn Phe
          85              90              95

Lys Ala Glu Glu Val Glu Leu Tyr Leu Glu Lys Leu Lys Glu Lys Arg
          100              105              110

Gly Leu Ser Gly Lys Tyr Gln Thr Ser Ser Lys Leu Phe Gln Asn Cys
          115              120              125

Ser Glu Leu Phe Lys Thr Gln Thr Phe Ser Gly Asp Phe Met His Arg
          130              135              140

Leu Pro Leu Leu Gly Glu Lys Gln Glu Ala Lys Glu Asn Gly Thr Asn
          145              150              155              160

Leu Thr Phe Ile Gly Asp Lys Thr Ala Met His Glu Pro Leu Gln Thr
          165              170              175

Trp Gln Asp Ala Pro Tyr Ile Phe Ile Val His Ile Gly Ile Ser Ser
          180              185              190

Ser Lys Glu Ser Ser Lys Glu Asn Ser Leu Ser Asn Leu Phe Thr Met
          195              200              205

Thr Val Glu Val Lys Gly Pro Tyr Glu Tyr Leu Thr Leu Glu Asp Tyr
          210              215              220

Pro Leu Met Ile Phe Phe Met Val Met Cys Ile Val Tyr Val Leu Phe
          225              230              235              240

Gly Val Leu Trp Leu Ala Trp Ser Ala Cys Tyr Trp Arg Asp Leu Leu
          245              250              255

Arg Ile Gln Phe Trp Ile Gly Ala Val Ile Phe Leu Gly Met Leu Glu
          260              265              270

Lys Ala Val Phe Tyr Ala Glu Phe Gln Asn Ile Arg His Lys Gly Glu
          275              280              285

Ser Val Gln Gly Ala Leu Ile Leu Ala Glu Leu Leu Ser Ala Val Lys

```

290 295 300  
 Arg Ser Leu Ala Arg Thr Leu Val Ile Ile Val Ser Leu Gly Tyr Gly  
 305 310 315 320  
 Ile Val Lys Pro Arg Leu Gly Val Thr Leu His Lys Val Val Val Ala  
 325 330 335  
 Gly Ala Leu Tyr Leu Leu Phe Ser Gly Met Glu Gly Val Leu Arg Val  
 340 345 350  
 Thr Gly Ala Gln Thr Asp Leu Ala Ser Leu Ala Phe Ile Pro Leu Ala  
 355 360 365  
 Phe Leu Asp Thr Ala Leu Cys Trp Trp Ile Phe Ile Ser Leu Thr Gln  
 370 375 380  
 Thr Met Lys Leu Leu Lys Leu Arg Arg Asn Ile Val Lys Leu Ser Leu  
 385 390 395 400  
 Tyr Arg His Phe Thr Asn Thr Leu Ile Leu Ala Val Ala Ala Ser Ile  
 405 410 415  
 Val Phe Ile Ile Trp Thr Thr Met Lys Phe Arg Ile Val Thr Cys Gln  
 420 425 430  
 Ser Val Ser Tyr Lys His Ile Tyr Glu  
 435 440

&lt;210&gt; 35

&lt;221&gt; 1670

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 35

aatcaggctc acccccaagt tgggggggtc attgacaagt cgaagagttg ggtccttgtg 60  
 tatgcatggg tgggatgga aggggaagag cccatggctg gatgtgctgg gaaccccggg 120  
 aagcctcttc agccattgtt gggcctagcc tgggacccga cagcaactcc gggatgggga 180  
 ctggggagtg ggcacacagt ggaagccatc tgggcagacc gaccccatgt gcagtcacct 240  
 ggacaggctt ctcctctctg agcaattcta gctccctctg agggccagtt ccagagacag 300  
 gccgagggtg gggagctccc accccatgct ctcttcacga cctcctacga gatgatgatg 360  
 cagtgtgtgt cccgcatgtt ggcctacccc ctgcatgtca tctcaatggc ctgcattgtc 420  
 cagtttgttg gacgggaggg caagtacagt ggtgtgttga gctccattgg gaagcttttc 480  
 aaagaggag ggcctgtggg attcttctgt ggattaatcc ctaactctcc gggcagatgtg 540  
 gttttcttgt ggggtctgaa cctgctggcc caatttcaca atgcttacct ggtggatgac 600  
 agcttcagcc agggcctggc catccggagc tataccaagt tegtgatggg gattgcagtg 660  
 agcatgttga cctacccctt cctgctagtt ggcagctcca tggctgtgaa caactggggg 720  
 ctgcaagctg ggtccccccc ttactcccca gtgttccaat cctggattca ctgctggag 780  
 tacctagtg tgcaggggca gctcttcaga ggtccagcc tgcctttctg ccgggtgtca 840  
 tcaggatcat gctttgacct ggagtaacct gaatctctca aaaaacacgg tctcaacctg 900  
 gccacgtgg gtgaggctg accaccttgg gacacctgca agcagactcc aacccaacaa 960  
 caaccagatg tgcctcagcc cagccggggt tggggctgca cccagtggat tgggtcacc 1020  
 gatgtggggt tgggcggggg tggggctgca ggaatcccat acrtgcaga tttgctgagt 1080  
 gggaaagtga ggcgaggttg ggagttggca tatgggggcc caggttggat ggggaagggc 1140  
 ctgtcttctg cagagggccc gagaatggct tctcagtcag cccagggccc atcctgcaga 1200  
 taatgggttc agacccccc cctctacccc ctccagtcag cccagggccc atcctgcaga 1260  
 tcagctggga gcatcattct cctgctttgt acataggggt tgggtccctg gcaagtggcc 1320  
 acctcaatgt ctaggcttat gctaggaggg aatatggccag gctctgctg tgtttttctc 1380  
 aacactactt ttctgatatg agggcagcac ctgctctgta atgggaatc atgcaactac 1440



```

tcagaaatgtg tctctctcat ctactgctca tctgtttaat gctgatgect cgcgtacagg 1500
atctggttac ctctgcagtt gtgaataccc agaggttggg cagatcagtg tctctagttc 1560
taccacagttt taaggttcat ggtcagattt gacctcatct ccgcacaaia atgtatttgg 1620
tgatttggga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1670

```

&lt;210&gt; 36

&lt;211&gt; 164

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

```

Met Gly Gly Met Val Arg Glu Glu Ala Leu Ala Trp Met Cys Arg Glu
  1                      5                      10                      15

Pro Arg Lys Ala Phe Ser Ala Ile Val Gly Pro Ser Leu Gly Pro Asp
      20                      25                      30

Ser Thr Pro Gly Trp Gly Thr Gly Glu Trp Ala Thr Gly Gly Ala Ile
      35                      40                      45

Leu Gly Arg Pro Thr Pro Cys Ala Val Pro Gly Thr Gly Phe Ser Leu
      50                      55                      60

Leu Ser Thr Cys Ser Ser Pro Arg Gly Pro Val Pro Glu Thr Gly Arg
      65                      70                      75                      80

Gly Trp Arg Val Pro Thr Pro Cys Ser Leu Pro Asp Leu Leu Arg Asp
      85                      90                      95

Asp Asp Ala Val Cys Val Pro His Val Gly Pro Pro Pro Ala Cys His
      100                      105                      110

Leu Asn Ala Leu His Gly Pro Val Cys Gly Thr Gly Gly Glu Val Glu
      115                      120                      125

Trp Cys Ala Glu Leu His Trp Glu Asp Phe Gln Arg Gly Arg Ala Ala
      130                      135                      140

Gly Ile Leu Arg Trp Ile Asn Pro Ser Pro Pro Gly Arg Cys Gly Phe
      145                      150                      155                      160

Leu Val Gly Leu

```

&lt;210&gt; 37

&lt;211&gt; 1493

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (1415)

&lt;400&gt; 37

```

ggaatggcggg cgcgcggccc gaactggag gcccggcggg ggccgcgggg agggccggct 60
gtcgaagacc gctgctgccc ccgcgcgggg cgcgcggctt tcaatggcgc catgcgcag 120
gacgggcagg cggcaagatg agaccgccc gccacgcatg tctcacttt tctggcgtt 180
catgatcttg gcaagctctg tcatgccta ctgcagtcag ctggccggcg gccactgtga 240
gattgtgacc ttggaacggg acagcagcca gactcgagg acgacggccc ggcagaccg 300

```

```

ccgctgtgag tctagaaagg ggcagatcgc cggcaccacg agagcccgcc cccctgtgt 360
ggacgcaaga atcatcaaga ccagccagtg gtgtgacatg ctccgtgttc tggaggsgga 420
aggtcgagac ttgttaataa accggtccgg ctggacgtgc acccagcccg gccggaggt 480
aaagaccacc acggtctcct gacaaacaca gccctgagg ggcgccggga gtggccttg 540
ctccctggag agccacgtc tcagccacag ttctccactc gctcggactc tcaccgttc 600
tctgcggccc gcccaatccg ttctccgtg gtccgtgaaq gacggcctca ggccttgga 660
tctgagctt ctgtctgtcc agccgacccc agggggccgg actcagacc ataggccggg 720
ggcgccacct ggcacagca atccgagtc tgtggagcc cggcccgcc cagcccccgc 780
cgacgtggtg gttggccctg ctgtctcccg agggaggagg gtagggagga gctccggcag 840
ccacagaaag ctgcagccca gcccgctga gacacgacgc ctgcccaggg ggaatgtcag 900
gcacagaagg ggcctctccc cgtgccccag actgtccgaa ttgttttat ttctttata 960
tttcagtata ccccatagac caagagacca aatctatctg aacctggag caccctcact 1020
gtcagggtcc ctgggggtcg ttgtgcccgc gggagggcaa tggtagcaga gacatgtgt 1080
ggcccccagg gaggcgagag gggggccgtg gtggagggct ccaccccagg agcaccccc 1140
gcacccctcg aggcagggtt tgggtgcgc gggggccgtg gacacctgc gggaggcagc 1200
gacggccccc acgcagccgc cgggaacgca ggcgccttta tctctctgta cttagatcaa 1260
cttgaccgta ctcaaatccc ttctgtttt aaccagttaa acatgctct tctacagtc 1320
catttttgat agttggatca tccagtatct gccagagcca tgttgggtct ccgtgactg 1380
ctgctcctat gatacccat ttagctccag aaganaaaga aaactcaggt accacttgt 1440
tgaaagagac cttatatgt attttgcaa gcttaaaaa aaaaaaana aa 1493

```

&lt;210&gt; 38

&lt;211&gt; 132

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

```

Met Ala Pro Ser Pro Arg Thr Gly Ser Arg Gln Asp Ala Thr Ala Leu
  1             5             10             15

```

```

Pro Ser Met Ser Ser Thr Phe Trp Ala Phe Met Ile Leu Ala Ser Leu
          20          25          30

```

```

Leu Ile Ala Tyr Cys Ser Gln Leu Ala Ala Gly Thr Cys Glu Ile Val
      35          40          45

```

```

Thr Leu Asp Arg Asp Ser Ser Gln Pro Arg Arg Thr Ile Ala Arg Gln
      50          55          60

```

```

Thr Ala Arg Cys Ala Cys Arg Lys Gly Gln Ile Ala Gly Thr Thr Arg
      65          70          75          80

```

```

Ala Arg Pro Ala Cys Val Asp Ala Arg Ile Ile Lys Thr Lys Gln Trp
          85          90          95

```

```

Cys Asp Met Leu Pro Cys Leu Glu Gly Glu Gly Cys Asp Leu Leu Ile
      100          105          110

```

```

Asn Arg Ser Gly Trp Thr Cys Thr Gln Pro Gly Gly Arg Ile Lys Thr
      115          120          125

```

```

Thr Thr Val Ser
      130

```

&lt;210&gt; 39

&lt;211&gt; 3693

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (108)

&lt;400&gt; 39

```

cctggcccaa ggaatgcccggt ttctgtctctaa agggagggtct ggccataaag cccctggatt 60
ggtggttaggg cggggtcgggc cccagtcctcc accctgaagc ggaagtgnag gaaagtctga 120
ggaccatcac cccgtgccca tggacatcca gccagccag ctgctcgaat ggttggtaga 180
cagaaggcac tgcagcctga aatggcagag tctgtgtctg agatcccggt agaagctcaa 240
tgctgccatc caggacatgc cagagagcga agagatcgcc cagctgctgt ctgggtccctg 300
tgagtgttta ggggtctgca ctggagtcgc ctcttctgtg aggtagtatg tatcagctga 360
gtactctctt tcttatctag ccaccagggc tggcctttag agaggagagc atttttattt 420
gaggtataga tctgtgcttc ttccactatg aatgtattta gggaaagct acacattcaa 480
gttaactggc gtatattata catgaatcag ccacctgcag tgcacagagt aggtgaagaa 540
gatacttttag ttctcgaaat cccgtctctt gctttctaga ctttcactac ttccactgac 600
taagaatcct ggaacttctc aaaggccacg aggcctccac gaagaatatt tttggccgat 660
actcttcaca gtcgatgaag gattcgccag agattatagc tctgtatgag aaggcaca 720
cctactctagg taagtggcc cggcctggga gccctggtat ccattgggaa gccactctc 780
agagttctga cctaccaggc ttataggagg caccgtctgt gactgggagc agactggagt 840
gtagatgttg cccatttcta ggtgttaaa tcaattgttt tagatggaat tgattttccc 900
tgagtggagt gctgggggaa ggaggaggtc caggccggtg gtggccattc gccgtgctc 960
agcagagcgg tgtgtgtggg tcttccacca ctcaactctt ggttagcggg agtgtgctgc 1020
ccccacccc accccgtac ccccatgtta caacaggcag aaggagccatg ggttttccctg 1080
ggagcgaata tcaagtgcct gaggacaaat acaggactca ctgtgtttgg gttgggtgta 1140
gtataaataa taataetggc taatatttcc tgaactctca cttaattgca ggaattgtgc 1200
ttgtgtgttc atgtgattc tctctgcat ttcaatgata catgttattg tegtgtttt 1260
accgatgagg gtgggattag aggggttaaa caactgtct taggtccac actgtggaa 1320
aagtggggtt gggagctga ctctgtgctc ttccaccaca caaaggatgt ggtgtgcatc 1380
tggggcctgc ctgctcctg tgggggtgtc ctgggtgaa ttctgtggc acttctcagt 1440
ggaactctct agcctcctgg ttccgaatgt caactatgag atccctcacc tgaagagcga 1500
gattgccaa ggcgcagc gcagtgccgg agcagttcta cccctcctgc agaggaggag agtgcacggc 1560
aggggttccc ggcctcttcg cagccagagg acacctgggc cctgtctgt ctactctga 1620
ggcggcgccg cccctcagcc tggttgcgcc ccttttgggc cagtgtctta cttttctctg 1680
gtctttggat gtttcttca atctgttgg ctccacctct tctccctct ctagggcga 1740
aatgtctcag gaaactgct ggcctgtgt agggacctgc ctgctcagc aggtgagatt 1800
ggggcagcgg ctcaagcctc cctgggggaa gccattgacg tgtaccagc gtctgtggg 1860
tttgtgtgtg agaggtagag aggcctcagc ttctcctggt ttctccata taccatttga actctttaca 1920
ccccagctca tgacctctc ttggccattt gaccatgttg gaccatgttg ctgttgagc ctgttgagc 1980
cactgtgacc catgtctggc atgtctggg ttctgtcaga agcgggggaa gacccacct ctcaacggc 2040
tgaccttacc atgtctggg agggagctc gttgtgtgt agcgggggaa gacccacct ctcaacggc 2040
aggtgttggc atgtctggg ttctgtcaga agcgggggaa gacccacct ctcaacggc 2100
ggacagggac agagccctct gtgttggaac gttgttgga gacccacct ctcaacggc 2160
tgccagaaga tgcgattgac tgggscagc ttgtgttgga gacccacct ctcaacggc 2220
actctggcat ctctgcagc ggtgttgga ggtgttgga gacccacct ctcaacggc 2280
cabagatcc tggaggtgat ggtatagac ggtgttgga gacccacct ctcaacggc 2340
cagtgttga atcactgag acccggaatc gttgttgga gacccacct ctcaacggc 2400
cactgttga atcactgag acccggaatc gttgttgga gacccacct ctcaacggc 2460
tcttcttagc ccagagagca ggtgttgga ggtgttgga gacccacct ctcaacggc 2520
agttccagc ggtccagcc atctgcagg ttctgttgga gacccacct ctcaacggc 2580
tgtcagtgct ggtgttgga atctgcagg ttctgttgga gacccacct ctcaacggc 2640
tgatcctggc ctccacagg ctctgttgga atctgcagg ttctgttgga gacccacct ctcaacggc 2700
gctctgctc ctctgttgga atctgcagg ttctgttgga gacccacct ctcaacggc 2760
ctggccattg cctctgttgga atctgcagg ttctgttgga gacccacct ctcaacggc 2820
catatacgg aagaatctgc atctgttgga atctgcagg ttctgttgga gacccacct ctcaacggc 2880
ccttaaaag cctctgttgga atctgttgga atctgttgga gacccacct ctcaacggc 2940
gagcctctca cctctgttgga atctgttgga atctgttgga gacccacct ctcaacggc 3000
ctccactgag tgcctgttgga atctgttgga atctgttgga gacccacct ctcaacggc 3060
tgagcagtc ccagctgttgga atctgttgga atctgttgga gacccacct ctcaacggc 3120
cacttgagga gcaggcggt ctgtgttgga atctgttgga gacccacct ctcaacggc 3180

```

```

agctgcagaa gctgattgaa gctgacatct ccaagaggta cagcaggaga cctgtgaacc 3360
tgatgggaa cctctctgtga cscctcctgt gttcttgcct gccctctctt tccctctctg 3420
ggatgaagat gatagccagg gctgttgttt tggggccctt caaggcaaaa gaccaggctg 3480
actggaagat ggaagagcac aggaagggaag cggcacctga tggtagatct ggcactctac 3540
atgtttctta caagaagctg tggtagattgg cctgttggta taccaggaga aacccacaga 3600
ttctctctct agttagtata gggactttaa taaaagagga aaaaactctt gttccaaaa 3660
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa                                     3693

```

&lt;210&gt; 40

&lt;211&gt; 230

&lt;212&gt; RAT

&lt;213&gt; Homo sapiens

&lt;400&gt; 40

```

Met Leu Arg Phe Val Gln Lys Arg Gly Asn Ser Thr Val Tyr Glu Trp
  1                      5                      10                      15

Arg Thr Gly Thr Glu Pro Ser Val Val Glu Arg Pro His Leu Glu Glu
                20                      25                      30

Leu Pro Glu Gln Val Ala Glu Asp Ala Ile Asp Trp Gly Asp Phe Gly
                35                      40                      45

Val Glu Ala Val Ser Glu Gly Thr Asp Ser Gly Ile Ser Ala Glu Ala
  50                      55                      60

Ala Gly Ile Asp Trp Gly Ile Phe Pro Glu Ser Asp Ser Lys Asp Pro
  65                      70                      75                      80

Gly Gly Asp Gly Ile Asp Trp Gly Asp Asp Ala Val Ala Leu Gln Ile
                85                      90                      95

Thr Val Leu Glu Ala Gly Thr Gln Ala Pro Glu Gly Val Ala Arg Gly
                100                      105                      110

Pro Asp Ala Leu Thr Leu Leu Glu Tyr Thr Glu Thr Arg Asn Gln Phe
                115                      120                      125

Leu Asp Glu Leu Met Glu Leu Glu Ile Phe Leu Ala Gln Arg Ala Val
                130                      135                      140

Glu Leu Ser Glu Glu Ala Asp Val Leu Ser Val Ser Gln Phe Gln Leu
                145                      150                      155                      160

Ala Pro Ala Ile Leu Gln Gly Gln Thr Lys Glu Lys Met Val Thr Met
                165                      170                      175

Val Ser Val Leu Glu Asp Leu Ile Gly Lys Leu Thr Ser Leu Gln Leu
                180                      185                      190

Gln His Leu Phe Met Ile Leu Ala Ser Pro Arg Ser Gly Phe Pro Leu
                195                      200                      205

Met Gln Gly Ser Ala Ile Leu Ser Ser Ser Ala Ser Leu Tyr Ser Ser
                210                      215                      220

Ser Cys Ser Met Thr Pro
                225                      230

```

<210> 41  
 <211> 1701  
 <212> DNA  
 <213> Homo sapiens

<400> 41  
 cccctgagat gatcttctct ttccaaacttc ttgaacttgg acatgaagga tgtgggcccc 60  
 gaatcatgtg gccagccccc cccctgttgg ccccaaccag ccttggagtc tcttctaggg 120  
 aaggccctccc agcatcttgg actcgagagt ggttagcccc tctacctctt ggagctgac 180  
 tggggttgaa ctgagtggtt tctttagctt accggggagg cagctgcttg ttctctccc 240  
 accagccctc tcccacatc ccagctgccc tggcttgggtc ctgaagccct ctgtctacct 300  
 gggagatcag ggaacacagg cctttagggat acagggggtc cctttctgtt accaccccc 360  
 accctctctc aggaacacac taggttggttc tggatgcttg ttctttagcc agccaaggtt 420  
 cactggcatt ctcccattgg gatcttgagg gaccaagctg ctgggatttg gaaggagttt 480  
 caccctgacc gttagcttag ccaggttccc aggaggccct accatactcc ctttcagggtc 540  
 cagggtctcc gnaagcccaa ggaagggttc ctgtgctgct gtctggttga gagcctgcca 600  
 ccgtgtgttg gtaggtgtgg ccagctgag tgcattgggt acaggccctt ggcctatggc 660  
 ctgggtgtgt gtagctcag ccttaggttc gcagtgttga gagggtgtt gtgggggaag 720  
 aggtgtgctt tcaagtggt tgtgtgcaag ggttgggtt gttagggga 780  
 cgtgtgttg cgtgtgttg ggcattgtg atgagtgct gccgtgaa gtgtccacag 840  
 ttgagaggtt ggaagcaggat gagggaatc tgtccctc aataatcact tgtgagctc 900  
 cagctctgcc caagacgcca cctgggggga cagccaggag ctctccatgg ccaggctgcc 960  
 tgtgtgcatg tccctgtct ggtgcccctt tgcctgctc ctgcacact cacagggtcc 1020  
 ctacacacaa gtgcccctca gaagcagccc ctggagga caagggaagg aatggggatg 1080  
 gctggggctc tctccatct cctttctct tgccttgc atgcttggtc tccctctca 1140  
 aaactccat tccctgtct ccagcccctt tgcctagct tgaatttgg gaggaggaag 1200  
 gggcgatttg aggaagagg ggaagagct tatggctgg ggcacacta tgcctgccc 1260  
 ggggttctta ctgttccag gtgctccag ggcaggcagg ggcacacta tgcctgccc 1320  
 ctggtaaaag tgaacctgc catttaccag cagccctggc atgttcttc cccacaggaa 1380  
 tgaataggag ggaagctccg aaactttcca tcccagaagc agtctcctg gttaggcag 1440  
 actggatttt tgcctgccc ctgacccctt gtcctctttt gaggagagg agctatgcta 1500  
 gaactctaac ctgagggaat cgggtgggtt ggcctagct ctttctgata tgaacctt 1560  
 taaggtggga ggttggcag ggaatgtctt aataatcaa tcccaagctt caaaaaaaa 1620  
 aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa 1680  
 aaaaaaaa aaaaaaaa a a 1701

<210> 42  
 <211> 240  
 <212> PRT  
 <213> Homo sapiens

<400> 42  
 Met Lys Asp Val Gly Pro Glu Ser Cys Gly Gln Pro Thr Pro Cys Trp  
 1 5 10 15  
 Pro Ser Pro Ala Leu Glu Ser Val Leu Gly Lys Ala Ser Gln His Leu  
 20 25 30  
 Gly Leu Glu Ser Gly Gln Pro Leu Tyr Leu Leu Glu Leu Asn Trp Gly  
 35 40 45  
 Gly Thr Glu Cys Val Leu Ser Ser Thr Gly Arg Thr Ala Ala Cys Phe  
 50 55 60  
 Leu Pro Thr Ser Leu Leu Pro Thr Ser Pro Ala Ala Trp Leu Gly Pro  
 65 70 75 80  
 Glu Ala Leu Cys Leu Pro Gly Arg Pro Gly Thr Thr Gly Leu Arg Asp  
 85 90 95

Thr Gly Gly Pro Leu Leu Leu Pro Pro Pro Thr Leu Leu Gln Asp Thr  
 100 105 110  
 Thr Arg Trp Cys Trp Met Leu Val Leu Trp Pro Ala Lys Val His Gly  
 115 120 125  
 Asp Ser Pro His Gly Ile Leu Arg Asp Gln Ala Ala Gly Ile Gly Lys  
 130 135 140  
 Glu Phe His Pro Asp Arg Cys Pro Ser Gln Val Pro Arg Arg Pro His  
 145 150 155 160  
 His Thr Pro Phe Gln Gly Gln Gly Ser Ser Lys Pro Arg Ala Arg Ile  
 165 170 175  
 Leu Cys Cys Cys Leu Val Glu Ser Leu Pro Pro Cys Val Gly Ser Val  
 180 185 190  
 Gly Gln Ala Glu Cys Ile Gly Asp Arg Ala Val Ser Met Gly Leu Gly  
 195 200 205  
 Val Cys Glu Leu Arg Pro Arg Cys Ala Val Trp Arg Arg Val Leu Ser  
 210 215 220  
 Gly Lys Arg Cys Gly Phe Lys Val Cys Val Cys Arg Gly Trp Val Cys  
 225 230 235 240

&lt;210&gt; 43

&lt;211&gt; 1784

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

```

aggtctagaa ttcaatcggt aatatctctt aagtttttaa aaactcgga taattatata 60
tatctttttt gccgtttata tttaggggtt ttgttgata aaatcaagtc ttggttggtg 120
cttgctgaat taatatatta tgagtggkgc atttttaagt atagtgaaca agacaccta 180
ttaagtacag tgataaagca tctatatctt gtaaaaaaaa aaaaatctg cctatgcctg 240
ttttttaaga aaaaaaaat gctgtatcg gctgtatgg gactgtaatg cgttagtggt 300
cttgacatat actggaaatg catgtatact ggcgtacttt atattctcta aaatgcttaa 360
tgcccttgaa attttgtaet caaaaaaaag ctttgaaaaa tctaaagggg agagtattct 420
ttaaagtttt taacataagc ctgtcaatgc acatgtcgac ggttagcatg tttagcaaac 480
cttgtagaat tataataagt ttgtagttac ctgtgaact ctcaatgcct ggcactcttt 540
aetgtcaata cagtttagtt attttgttct gtctctgcat gtgccacaaa atatgtactt 600
ttttcacttt tctcccttg tatctcagtt acgggttaca actgggttcat tctgaacaaa 660
acaacacaaa aagtcacatc atatttttta scaattgtat aagtgcacaa gtaattcact 720
acagcctaaa gccctgcctt tgtaakttga cttctgacal gttggcaatc aaagcctgca 780
cttgtaacaa tgaaaaagaa aaagcatttt atcttactac tcaataaaat gtgcataaac 840
ttacagaatt ctctctcttc cactgagtcg gctgaaggga tttatgtgca caacacacat 900
gtgtctctta gctgctggcc caccacacaa catcacaggg ctacttccac tctaagaaag atctttggtg 1020
ctaggggctt cgtgacttga ggggtggtgc ctacttccac cgcagatgcc atcttttggc 1080
gattttgtgc tcaaatcaga taagagaagc ctgtttaaag gctttctctt cattgaacaa taanattaat 1140
ttctcaagg agccagttaa aaaccagag ctgtakttga tgcctatctt atctgctaaa gcaaaaagta 1200
ctaggtactt aagtgacttt tcatcacaga aaagcgttgc atttgtatta acaagaaakt 1260
tgtatacca cgttccagct actatctaat catcacccga agatttaaga tacaacaaat 1320
ttcagtttgt ttgtaacatt gtctactctt agtgcacttt gttttatata ataaagtatg 1380
cttgttatat taataataa gaatatggca attagcgata tagcataccc aaacaaagat 1440
gttctcgata cagtcggca aagactatcc caaggttatt ttaatgaatt cagacatttt 1500
ttctgtgga tatttctcca tctaaaaa aatggcaacc aaggaaata tttagatgca 1560

```

```

acttactaga gtgatgaggt gaaagaaatg gtgattcttg tattatgggt tttattttct 1620
ttcttataac tgcagagaaa atatcttgac taaaaaaat tcaattttttt ggaattccttt 1680
cttttacaaa ttgtctctgg gcaactatgg catagaaata aacatttgac attaaaaata 1740
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 1784

```

<210> 44  
 <211> E2  
 <212> PRT  
 <213> Homo sapiens

<400> 44  
 Met Cys His Lys Ile Cys Thr Phe Phe Thr Phe Phe Pro Leu Tyr Ile  
 1 5 10 15  
 Ser Tyr Gly Leu Gln Leu Val His Ser Glu Asn Asn Asn Asn Lys Ser  
 20 25 30  
 Pro Phe Ile Phe Phe Asn Asn Cys Ile Ser Ala Gln Val Ile His Tyr  
 35 40 45  
 Ser Leu Lys Pro Cys Leu Cys Asn Leu Thr Ser Asp Met Leu Ala Ile  
 50 55 60  
 Lys Ala Cys Thr Cys Asn Asn Glu Lys Glu Lys Ala Phe Tyr Ile Thr  
 65 70 75 80  
 Thr Gln

<210> 45  
 <211> 1034  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (598)

<400> 45  
 ggaagatggc ggcctctggg gcggagcggc aggtctctggc acaatacttg gtgttaagaa 60  
 aggatctatc acaagctcgg ttctcttgac cggggggcgc actggtagcg cagggtttgtc 120  
 acggggccac cggggccttg ccaactcacc ggcactaccc gcacacagcc gcttacctcc 180  
 aagagctggg gcgatcggc aaagtgggtc tggaggcccc agatgagacc accctaaagg 240  
 agctggcggc gaccctgcaa cagaagaaca ttgaccacat gctgtggcct gagcaaccag 300  
 agaatatcgc cacttgattt gctctccggc cctaccacca ggaagagtg ggcaggtatt 360  
 tgaagaaagt ccgattgttc aagtaactgc tgccttgatg tgtttgaata cgcaggccac 420  
 ccattccaaa gcacatgtg ttccttgcag tgtcagcttg ctccgctctt tcagttggtg 480  
 caattttctg aggttcaagc acatgttcat attaaagttg tcatttaata ctactccctc 540  
 ttatttaata gtccaagtgg ggaaggtggg agagcagtat tgtctgggga tcattgcnca 600  
 aatagaagat ttggttagac tctctctgtg ggtccaagga aactcccttc cagtcactcg 660  
 ggtttgaaac ttctgttttg aattccctct tcttccatc cagttatcat atttcattga 720  
 atccnagata aactcaattt caagatggcg tagtatttta tgtattgtta aaaaatatgc 780  
 cggcaaatca aacacttgta ttccaataac aaagatgtta aaatttggcc agtgtgtgtg 840  
 ctacatctg ttaattccag ggttttggga agccaaggcc ggaagatcgc ttgagcccat 900  
 ggttcaagg ttaagtcag ttctaatacg gcaacggcac ttcagcgtgg gcaacagagt 960  
 gagacatggg ttctataaag atttaataca agttaaaaaa aaaaaaaada aaaaaaaada 1020  
 aaaaaaaada aaaa 1034

<210> 46

<211> 126  
 <212> PRT  
 <213> Homo sapiens

<400> 46

```
Met Ala Ala Ser Gly Ala Glu Pro Gln Val Leu Val Gln Tyr Leu Val
  1           5           10           15

Leu Arg Lys Asp Leu Ser Gln Ala Pro Phe Ser Trp Pro Ala Gly Ala
      20           25           30

Leu Val Ala Gln Ala Cys His Ala Ala Thr Ala Ala Leu His Thr His
      35           40           45

Arg Asp His Pro His Thr Ala Ala Tyr Leu Gln Glu Leu Gly Arg Met
      50           55           60

Arg Lys Val Val Leu Glu Ala Pro Asp Glu Thr Thr Leu Lys Glu Leu
      65           70           75           80

Ala Glu Thr Leu Gln Gln Lys Asn Ile Asp His Met Leu Trp Leu Glu
      85           90           95

Gln Pro Glu Asn Ile Ala Thr Cys Ile Ala Leu Arg Pro Tyr Pro Lys
      100          105          110

Glu Glu Val Gly Gln Tyr Leu Lys Lys Phe Arg Leu Phe Lys
      115          120          125
```

<210> 47  
 <211> 1626  
 <212> DNA  
 <213> Homo sapiens

<400> 47

```
caacttctgtt agctgagagt ttgttttgtga cttattacag agcctgtgac ttaaaatcc 60
ttcccacac cacaagctaa agtggggagaa gacaaactac ctacaccttt caaccagag 120
ggaggagcaa aaatcagtgat actttttacag aagaacctgc cagcctgtga tgatcctacc 180
aaagagaaac ctcaatgagc tatggaaatt ccttttttgt gaattgagtg ctgttttttgc 240
ttttctcaga ttccaaatga gagtataca tttcttttgt ttgatgtgct ggttgagatc 300
tgataatana agaccatgct ttgaattctc tcagctaaqt gtaaggatt ccttcagaga 360
tttattttat ccgagaatag agaccatktc gatgatgtat acaaggaaca acctaaactc 420
tgctgagcca ctgtttgaac aaataaactc acttaatgtt aatttcaaca cacaaagaa 480
aacagtcttg cttattcacg gatcaagacc agtaggctcc atcccattat ggcttcagaa 540
cttcgttaag attttgttga atgaagaaga tatgaatgta attgtatgag actgagagcc 600
gggtgctaca acctttattt alaatagagc agttaaaac actagaaag ttgctgtgag 660
tttgagtgtg caccatana atcttttgaa gcatggtgca tctcttgaca attttcatt 720
cataggtgtg agtttagggg ctcatatcag tggatttgtt ggaagatat ttcattgta 780
acttggaaga ataacaggtc ttgacctgc tggpccaggt ttctccagaa aaccacata 840
tagcagatta gattacaagg atgcaagtt tctgcatgct atccattctg actccaatgg 900
aattcaattc attaatgca accaccagag agcagttcac ttgttcattg catctttaga 960
aacaaactgc aattttattt catttccttg tcttcaatc aaagattaca agactagctt 1020
atgtgtggac tgtgactgtt ttaaggaana atcatgtcct cggctgggtt atcaagccaa 1080
gctatttaaa ggtgttttaa aagaaggat ggaaggaga cctcttagga ctactgtgtt 1140
tttgatatac agtgcctatt attttgttct cagataatt gtccagata aaactatgat 1200
ggatggctcg ttttcattta aattatttaa tcagcttggg atgattgaag agccaagct 1260
ctatgaagaa agataacata tgtaaaagag gcacactaac totaacaac tagtgacttt 1320
aaaagtctta agcgtatcag gagaaggaga cctcctggc taactgtgtg aaacctgtc 1380
tctactaaa atkcaaaaa ttagctgggc atggtggcac gtgctgtgag tccagctac 1440
```



```

tcaggagagct gaggaagag gatgcttga acccaggagg tggaggttgc agtgagctga 1500
gattgcacag ctgcctcca gctgggtga cagagcaga cctccattca aataaataa 1550
taataaata aataaataa taataaata aataaataa gtttaagagt aaaaaaaa 1620
aaaaaa 1626

```

&lt;210&gt; 48

&lt;211&gt; 368

&lt;212&gt; FRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

```

Met Ile Leu Pro Lys Arg Asn Leu Asn Glu Leu Trp Asn Phe Leu Phe
  1             5             10             15

```

```

Gly Glu Leu Ser Ala Val Phe Ala Phe Leu Arg Phe Gln Met Arg Val
          20             25             30

```

```

Tyr Ile Phe Leu Cys Leu Met Cys Trp Val Arg Ser Asp Asn Lys Arg
          35             40             45

```

```

Pro Cys Leu Glu Phe Ser Gln Leu Ser Val Lys Asp Ser Phe Arg Asp
          50             55             60

```

```

Leu Phe Ile Pro Arg Ile Glu Thr Ile Leu Met Met Tyr Thr Arg Asn
          65             70             75             80

```

```

Asn Leu Asn Cys Ala Glu Pro Leu Phe Glu Gln Asn Asn Ser Leu Asn
          85             90             95

```

```

Val Asn Phe Asn Thr Gln Lys Lys Thr Val Trp Leu Ile His Gly Tyr
          100            105            110

```

```

Arg Pro Val Gly Ser Ile Pro Leu Trp Leu Gln Asn Phe Val Arg Ile
          115            120            125

```

```

Leu Leu Asn Glu Glu Asp Met Asn Val Ile Val Val Asp Trp Ser Arg
          130            135            140

```

```

Gly Ala Thr Thr Phe Ile Tyr Asn Arg Ala Val Lys Asn Thr Arg Lys
          145            150            155            160

```

```

Val Ala Val Ser Leu Ser Val His Ile Lys Asn Leu Leu Lys His Gly
          165            170            175

```

```

Ala Ser Leu Asp Asn Phe His Phe Ile Gly Val Ser Leu Gly Ala His
          180            185            190

```

```

Ile Ser Gly Phe Val Gly Lys Ile Phe His Gly Gln Leu Gly Arg Ile
          195            200            205

```

```

Thr Gly Leu Asp Pro Ala Gly Pro Arg Phe Ser Arg Lys Pro Pro Tyr
          210            215            220

```

```

Ser Arg Leu Asp Tyr Thr Asp Ala Lys Phe Val Asp Val Ile His Ser
          225            230            235            240

```

```

Asp Ser Asn Gly Ile Gln Phe Ile Lys Cys Asn His Gln Arg Ala Val
          245            250            255

```

```

His Leu Phe Met Ala Ser Leu Glu Thr Asn Cys Asn Phe Ile Ser Phe

```

260	265	270
Pro Cys Arg Ser Tyr Lys Asp Tyr Lys Thr Ser Leu Cys Val Asp Cys		
275	280	285
Asp Cys Phe Lys Glu Lys Ser Cys Pro Arg Leu Gly Tyr Gln Ala Lys		
290	295	300
Leu Phe Lys Gly Val Leu Lys Glu Arg Met Glu Gly Arg Pro Leu Arg		
305	310	315
Thr Thr Val Phe Leu Asp Thr Ser Ala Tyr Tyr Phe Val Leu Ser Ile		
325	330	335
Ile Val Pro Asp Lys Thr Met Met Asp Gly Ser Phe Ser Phe Lys Leu		
340	345	350
Leu Asn Gln Leu Gly Met Ile Glu Glu Pro Arg Leu Tyr Glu Glu Arg		
355	360	365

<210> 49  
 <211> 1221  
 <212> DNA  
 <213> Homo sapiens

<400> 49  
 ggaagaagctg agaataatca cctctgataa agatcacaga agctgccagg gaggtgtttg 60  
 attaatctca tctattgaaa atattgttca gaccccatgt gacataactg gagccagctgc 120  
 agtgcacatga agaactacga gattagcctg gatattaact tctcttctag agaataagatt 180  
 tcatgttcca tctctctgca atggttaatt cacacagaaa ccccatgttt aacattccaa 240  
 gaggaattta ctgcttaaca gccatcttgc ccccaatatg catttgttct cagttctcag 300  
 tgcactctag ctatcacttc actgaggctc ctggggcctt cccagtagcc actaatgggg 360  
 aacgatttcc ttggcaggag ctaaggctcc ctagtgtggt catctctctc cattatgacc 420  
 tctttgtcca ccccaatctc acctctctgg cctttgttgc atctgagaag atcgaggtct 480  
 tggctcagaa tgcacccag tttatcatct tgcacagcaa agatcttgaa atcaggaatg 540  
 ccccccctca gtccagaggaa gattccagat acatgaaacc aggaanaagaa ctgaaagtgt 600  
 tgaagttacc tgcctatgaa caaattgcac tctctgttcc agagaaaact acgctctccc 660  
 tgaataacta tctggtctatg gacttccaaag ccaggttagg tgatggcttt gaagggtttt 720  
 atadaagcac atccagaact ctggtgtgtg aaacaagaat tcttgcaata acagattttg 780  
 agccaaacca ggcacgcctg cctttccctt gctttgatga accgttgttc aaagccaaat 840  
 tttcaatcaa gatcagaaga gagagcaggg atattgcact atccaacatg ccaagggtgt 900  
 ccatctatgc atccccagac aaacgggaat aaacacatta tcttttgcag gcctcactga 960  
 agctacttga tttttatgaa agtactttg atacttacta tccactctcc aaactgggta 1020  
 tgttcanaat ccccatctat ctcttcaatt ttgtctatga aacttgctta gatctcttcc 1080  
 ctcttctctc ttgtatgtga tttaaatgag caactgaggaa ttcagtttagc ccagggaanaa 1140  
 atantttgtt cctcagagat gattcttgag tgtagaaaat aaatatatta tgcactgccc 1200  
 caaananaaa aaaaaaaaaa a 1221

<210> 50  
 <211> 305  
 <212> PRT  
 <213> Homo sapiens

<400> 50  
 Met Phe His Ser Ser Ala Met Val Asn Ser His Arg Lys Pro Met Phe  
 1 5 10 15  
 Asn Ile His Arg Gly Phe Tyr Cys Leu Thr Ala Ile Leu Pro Gln Ile  
 20 25 30

Cys Ile Cys Ser Gln Phe Ser Val Pro Ser Ser Tyr His Phe Thr Glu  
           35                          40                          45  
 Asp Pro Gly Ala Phe Pro Val Ala Thr Asn Gly Glu Arg Phe Pro Trp  
           50                          55                          60  
 Gln Glu Leu Arg Leu Pro Ser Val Val Ile Pro Leu His Tyr Asp Leu  
           65                          70                          75                          80  
 Phe Val His Pro Asn Leu Thr Ser Leu Asp Phe Val Ala Ser Glu Lys  
                           85                          90                          95  
 Ile Glu Val Leu Val Ser Asn Ala Thr Gln Phe Ile Ile Leu His Ser  
                          100                         105                         110  
 Lys Asp Leu Glu Ile Thr Asn Ala Thr Leu Gln Ser Glu Glu Asp Ser  
           115                         120                         125  
 Arg Tyr Met Lys Pro Gly Lys Glu Leu Lys Val Leu Ser Tyr Pro Ala  
           130                         135                         140  
 His Glu Gln Ile Ala Leu Leu Val Pro Glu Lys Leu Thr Pro His Leu  
           145                         150                         155                         160  
 Lys Tyr Tyr Val Ala Met Asp Phe Gln Ala Lys Leu Gly Asp Gly Phe  
                          165                         170                         175  
 Glu Gly Phe Tyr Lys Ser Thr Tyr Arg Thr Leu Gly Gly Glu Thr Arg  
                          180                         185                         190  
 Ile Leu Ala Val Thr Asp Phe Glu Pro Thr Gln Ala Arg Met Ala Phe  
                          195                         200                         205  
 Pro Cys Phe Asp Glu Pro Leu Phe Lys Ala Asn Phe Ser Ile Lys Ile  
           210                         215                         220  
 Arg Arg Glu Ser Arg His Ile Ala Leu Ser Asn Met Pro Lys Val Ser  
           225                         230                         235                         240  
 Ile Tyr Ala Ser Pro Asp Lys Arg Asn Gln Thr His Tyr Ala Leu Gln  
                          245                         250                         255  
 Ala Ser Leu Lys Leu Leu Asp Phe Tyr Glu Lys Tyr Phe Asp Ile Tyr  
                          260                         265                         270  
 Tyr Pro Leu Ser Lys Leu Gly Met Phe Lys Phe His Ile Ile Val Phe  
           275                         280                         285  
 Ile Phe Ala His Lys Thr Cys Leu Asp Leu Phe Pro Leu Ser Leu Cys  
           290                         295                         300

Met  
305

<210> 51  
 <211> 951  
 <212> DNA  
 <213> Homo sapiens

```

<400> 51
gggtggtgag gagtctggg cggttccgc ggctctctcc tcttccccgt tcccttccac 60
cccccccgcc acccttttcc cccctccggc tccgtccacc tcccgctccc cccctccagg 120
acaagaatgc cctgcgggga acaaccacac agcgctcaga tggctttggg cccggtccag 180
cggtaaccta ccccccagac caccctccagc cctctcgcct cggagggcag cagtggggag 240
gaagaatgcc ggtaacagcc caggagcacc agcgagagct ttctaacctg ccaaggtgct 300
gaccttttcc taccacgggg aattggctca tccacaccaa gaatcagcca cagcgggaac 360
aagcatgcag ggatctctca acagccttcc caagcaatgt tccctttact ccgcccagaa 420
gcccacatca ggcctggctgt aagactggaa agtacttacc agaatcgacc acgtatatg 480
gtagtgggtt ccactaatgg tagacaaagc actgaagaaa gtatcgtctt aggaatggat 540
ctctctctta atgacagcac ttgtaccatg ggccttgggt tgcctctctg gagcgacag 600
ctatttcatc tggatggtsa tggctgggtc agtctataga cggataaccg agttcacata 660
ctcaaacctg tatctgtgca gcaatgtgg gtgcacaggg attcaaggaa caaacactgt 720
gatgtactat tgggtggaag atgaactgga gcagccttcc tggagagtga ttgcccaca 780
tgccttatca ttltgcctga tctttgtcct agtaacctta tctctatgga tttactctaa 840
gtttgtaac atggatgtgt gcaagattt tagctctaac aatgtttgtc agtgttctaa 900
taatagcaaa aaataaaaa caaatgattg aaaaaaaaa aaaaaaaa a 951

```

<210> 52

<211> 194

<212> PRT

<213> Homo sapiens

<400> 52

```

Met Ala Leu Val Thr Val Gln Arg Ser Pro Thr Pro Ser Thr Thr Ser
 1             5             10             15

Ser Pro Cys Ala Ser Glu Ala Asp Ser Gly Glu Glu Glu Cys Arg Ser
          20             25             30

Gln Pro Arg Ser Ile Ser Glu Ser Phe Leu Thr Val Lys Gly Ala Ala
          35             40             45

Leu Phe Leu Pro Arg Gly Asn Gly Ser Ser Thr Pro Arg Ile Ser His
          50             55             60

Arg Arg Asn Lys His Ala Gly Asp Leu Gln Gln His Leu Gln Ala Met
          65             70             75             80

Phe Ile Leu Leu Arg Pro Glu Asp Asn Ile Arg Leu Ala Val Arg Leu
          85             90             95

Glu Ser Thr Tyr Gln Asn Arg Thr Arg Tyr Met Val Val Val Ser Thr
          100             105             110

Asn Gly Arg Gln Asp Thr Glu Glu Ser Ile Val Leu Gly Met Asp Phe
          115             120             125

Ser Ser Asn Asp Ser Thr Cys Thr Met Gly Leu Val Leu Pro Leu Trp
          130             135             140

Ser Asp Thr Leu Ile His Leu Asp Gly Asp Gly Gly Phe Ser Val Ser
          145             150             155             160

Thr Asp Asn Arg Val His Ile Phe Lys Pro Val Ser Val Gln Ala Met
          165             170             175

Trp Val Asp Arg Asp Ser Arg Asn Lys His Cys Asp Val Leu Leu Val
          180             185             190

```

Glu Glu

<210> 53  
 <211> 1514  
 <212> DNA  
 <213> Homo sapiens

<400> 53  
 gcacgacatt tttacgggtc acccatattg catgtatcag gaataataac ctttttatka 60  
 ttgagtagtg ttctattgta tgtatatacc acagtttatt tctcccttca tcttttgeta 120  
 gattttcggg ttttttcaca ttgcgctatt cagtataaac ctgctctcaa ctttcattgt 180  
 caagtctttg agtgcacata tatttgcggt tctcttgagt gaatgcacct tgttgggtca 240  
 cgtggcttaa ctttaaaaaa ttttaattca tgtgttgcat atgtagtgat tattaagtat 300  
 tatctcataa ttttatttcc ttgcttaatt atgttgagtg tatttcattt gtattttagt 360  
 ttgcacatgt ttgttcaaat tcttcacctg tttttaaaga agcagtaaga cttatttttg 420  
 tgttctgaac ataggttctt tctcacataa aatgtgctat gaatgttagg tttcaaatac 480  
 tccaaatgaa tggctagaga attactatct gtagaastat ttatctgtca aagggatgct 540  
 aacattttac tttattgctc taaaatagaa aagttgcagg aatgtctgtg agtttttagt 600  
 gaaacacatga tagctgggtt tactagtgaa atttgagttt taantgtcaa tgttagctaa 660  
 cggccagat agggacacct gcagggtggt tacttgagc tgtgactcaa ctggtccttc 720  
 actgcacaa atacctgggg ttgactcatt ggctgaact ttgcaaatg aggaacctta 780  
 ggcacaaatca gtgaatttct gaactgcctt cgtcttcagt tatatgggga tttcccaact 840  
 tttgagakec ttgtaaggat tatatgagat gaagagatga gacaggtat ataaaagtcc 900  
 cagcacagag cgtgtcatat aatatggctt cacaaqtacc ctcatctcct ttcagtcgt 960  
 tttttgttct tgtttttgtt tttttgagac catctcactc tgttggccag gctggagtgc 1020  
 ctcttcattt ttattctctt attcagcaag tacttgatca atgtgctttg taccaggtac 1080  
 tgagctcttc gttgggtat atgtgtgac aeggagattg tagattcttg cagggaacac 1140  
 tgaatcaaa caaggcgacc cgaactagt agacctgtc tctactagaa gaattttaa 1200  
 aatcacctag atgtggcgcg ggcacggctg ctacagcctg tggccccagc actttgggat 1260  
 gctgaggcgg gtggtcacg ggttcaggag atcagagaca tctggnata cagggaaca 1320  
 ccccgctctc actggaaata caaggaaatt ggcggggcgt gggggcgggc atctgtggtc 1380  
 ccaattactc gggaggctgc agcaggagag tggcatgaac cggggaggcg gattttgcat 1440  
 tgagccgaga tcaagccact gcactccagc ctggggcaga gaatagagct ccatctcaaa 1500  
 aaaaaaaaaa aaaa 1514

<210> 54  
 <211> 91  
 <212> PRT  
 <213> Homo sapiens

<400> 54  
 Met Ala Ser Gln Val Pro Ser Ser Pro Phe Gln Ser Phe Phe Val Phe  
 1 5 10 15  
 Val Phe Val Phe Leu Arg Pro Ser His Ser Val Ala Gln Ala Gly Val  
 20 25 30  
 Pro Leu His Phe Tyr Phe Phe Ile Gln Gln Val Leu Ile Lys Cys Ala  
 35 40 45  
 Leu Tyr Gln Val Leu Ser Ser Ser Leu Gly Tyr Asn Gly Asp Gln Gly  
 50 55 60  
 Asp Cys Arg Phe Trp Gln Gly Lys Leu Thr Ser Asn Thr Ala Thr Arg  
 65 70 75 80  
 His Ser Glu Thr Leu Ser Leu Leu Glu Glu Leu

<210> 55  
 <211> 1417  
 <212> DNA  
 <213> Homo sapiens

<400> 55

```

gtccaaatcc tattgtccac agtcagactt ctacaaactc cletgaacaa atgcagcctc 60
caatgtttcc ctctcaaatg accattgctg tgttacaggg ctcttcagtt cctcaagacc 120
agcagtcacac caacctatct ctctccaga gtcccatgaa taatcttcag actaacacag 180
tagcccaaga agcatctttt gcagacccga actcaatttc tccacttcag tcaacatcaa 240
acagtgaaca caaagctgct ttcccaacag aagctccaat atcacacac cagactccta 300
tgctttccc agaacaggca caacccccgc agtcagggtt atttcagcct cagggtggcc 360
tgggtctcct tccacctaat ccaatgcctc aaagccaaac aggaaccatg ttccagtcac 420
agcactcaat agttccatg cagagtaact ctccatccca ggaacagcag cagcagcagc 480
aacagcagca gcaacagcag cagcaacaa acagagcact ttatttcagt aatcagaata 540
ccatggctac aatggctct ccaagcacc cccacccaa catgatctt aacccaatc 600
aaatccaat ggtaatcag gagcaacaga accagtcaat tttccacca caagtaaca 660
tggccccaat gaatcaagc caacagcccc tgcatttca gactcagtc acagtttct 720
cacttcagaa cccgggtcct acctagtcgg aatcatcacc gccccccttg ttcctagct 780
ctctcagat tcagttggta caagggctac ctagtctca agagcagcaa gtaactctct 840
tcttatctcc agcatccatg ctgccttgc agaccagta aatcaacaa gatctgcaa 900
agtctctct ttattctct cagaacaaac tgcctggaa tcaaggagcc acattttgc 960
ctcaccaca ggtacttta ttccacaaa cagcaggagg caaatgaa caactgcag 1020
attctcctgg ctcatctcag cagacatcag gaatgttct atttggcatt caaataaact 1080
gtagtcagct tttaacctct ggaccagcta cattgctga tcagttgat gccataagtc 1140
agccaggcca accacaaac gagggccagc cactgtgac aacacttct tctcagcaa 1200
tgccagagaa tctccactg gcctctctta taacaccaa ccgaacac ccaagagctg 1260
atttgcctgt ttcattgca aacccaggga acacttgac tggctcctt taactggata 1320
taaattccac gaagcaatc ctgattccaa gatctgtga gatctgtg ttccatgaga 1380
atttactctt caaadacaa acaaaaaaa aaaaaaa

```

1417

<210> 56  
 <211> 420  
 <212> PRT  
 <213> Homo sapiens

<400> 56

```

Met Gln Pro Pro Met Phe His Ser Gln Ser Thr Ile Ala Val Leu Gln
  1             5             10             15

Gly Ser Ser Val Pro Gln Asp Gln Gln Ser Thr Asn Ile Phe Leu Ser
      20             25             30

Gln Ser Pro Met Asn Asn Leu Gln Thr Asn Thr Val Ala Gln Glu Ala
      35             40             45

Phe Phe Ala Ala Pro Asn Ser Ile Ser Pro Leu Gln Ser Thr Ser Asn
      50             55             60

Ser Glu Gln Gln Ala Ala Phe Gln Gln Gln Ala Pro Ile Ser His Ile
      65             70             75             80

Gln Thr Pro Met Leu Ser Gln Glu Gln Ala Gln Pro Pro Gln Gln Gly
      85             90             95

Leu Phe Gln Pro Gln Val Ala Leu Gly Ser Leu Pro Pro Asn Pro Met
      100             105             110

```

Pro Gln Ser Gln Gln Gly Thr Met Phe Gln Ser Gln His Ser Ile Val  
 115 120 125  
 Ala Met Gln Ser Asn Ser Pro Ser Gln Glu Gln Gln Gln Gln Gln  
 130 135 140  
 Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Ser Ile Leu Phe Ser  
 145 150 155 160  
 Asn Gln Asn Thr Met Ala Thr Met Ala Ser Pro Lys Gln Pro Pro Pro  
 165 170 175  
 Asn Met Ile Phe Asn Pro Asn Gln Asn Pro Met Ala Asn Gln Glu Gln  
 180 185 190  
 Gln Asn Gln Ser Ile Phe His Gln Gln Ser Asn Met Ala Pro Met Asn  
 195 200 205  
 Gln Glu Gln Gln Pro Met Gln Phe Gln Ser Gln Ser Thr Val Ser Ser  
 210 215 220  
 Leu Gln Asn Pro Gly Pro Thr Gln Ser Glu Ser Ser Gln Thr Pro Leu  
 225 230 235 240  
 Phe His Ser Ser Pro Gln Ile Gln Leu Val Gln Gly Ser Pro Ser Ser  
 245 250 255  
 Gln Gln Gln Gln Val Thr Leu Phe Leu Ser Pro Ala Ser Met Ser Ala  
 260 265 270  
 Leu Gln Thr Ser Ile Asn Gln Gln Asp Met Gln Gln Ser Pro Leu Tyr  
 275 280 285  
 Ser Pro Gln Asn Asn Met Pro Gly Ile Gln Gly Ala Thr Phe Ser Pro  
 290 295 300  
 Gln Pro Gln Ala Thr Leu Phe His Asn Thr Ala Gly Gly Thr Met Asn  
 305 310 315 320  
 Gln Leu Gln Asn Ser Pro Gly Ser Ser Gln Gln Thr Ser Gly Met Phe  
 325 330 335  
 Leu Phe Gly Ile Gln Asn Asn Cys Ser Gln Leu Leu Thr Ser Gly Pro  
 340 345 350  
 Ala Thr Leu Pro Asp Gln Leu Met Ala Ile Ser Gln Pro Gly Gln Pro  
 355 360 365  
 Gln Asn Glu Gly Gln Pro Pro Val Thr Thr Leu Leu Ser Gln Gln Met  
 370 375 380  
 Pro Glu Asn Ser Pro Leu Ala Ser Ser Ile Asn Thr Asn Gln Asn Ile  
 385 390 395 400  
 Glu Lys Ile Asp Leu Leu Val Ser Leu Gln Asn Gln Gly Asn Asn Leu  
 405 410 415  
 Thr Gly Ser Phe  
 420

<210> 57  
 <211> 2297  
 <212> DNA  
 <213> Homo sapiens

<400> 57

```

gaagtgaagg ttgaatgata cccacttaac taataaatga ataagcgtac ttgaantgat 60
tttttaaaagt gtttggtagt ctatacttat gttctttctt tgtttccact atagacagta 120
ttcgtgggcta ctttggggaa acaattgctc tgaacttttg atttttggag tatttcactt 180
ttgcattaat ccccatgggt gtcatgggt taacttaata cttgtttgtg tgggaagact 240
atgacaagta cgtgatcttt gectcggtca acctaatctg gtccacgggtg atctctggaa 300
tctgggaagcg tggctgtgac aacatgaact acaggttggg gacactgctc atgaagagaa 360
agtttgaggga gccccgggca ggatttcatg gtgttttggg tatcaatcac atcactggga 420
agggaggagcc tctgtacccc agctacaaga gacagktgag cttttacctg gtctccctgc 480
cattcgtgtg cctctgcttc tatttctcac tgtatgtcat gatgatttac ttgcacatgg 540
aggtttgggc cttgggtcta catgagaaca ggggtctga gtggaccagt gtccctgtct 600
atgtgccccg catcatctat gccattgtga ttgagatcat gaatcgtctc tatcgtatg 660
ctgcagagtt tttaaactta tgggagaatc acagattgga atctgcctat cagaaccatc 720
taattctgaa agttttagt ttcaacttcc caaattgctt tgcctcactc ttctatatg 780
cctttgtctt gaagatata agcttttgc gccagagctt ggcacctctc cttaattact 840
cccagatcct caaccacatt atggaactct tcttctctta ttggtctcaa aggaagcatg 900
gtgtgcaggt gaaggagaa gctgcaggct taaggctaga cattgatgct ecatttatg 960
aacaagtcat cctggaaaaa gaaatgggaa cttatttggg cactcttgat gattacttgg 1020
agttatctct gcagtttggg tatgtgagcc tttctctctg tgtttacca ttacagctg 1080
cctttgctgt gtttaataac ttcactgaag taatttcaga tgccttaaaa atgtgcaggg 1140
tcttcaaacg tccattctca gaaccttcag ccaatatttg tgtgtggcag atgatatctt 1200
gtttggacac aggtgtaaag agagggtctg attgcagggt catgaagcat cttttggggg 1260
aaatggcaat gtcctgtgct ttgtttggg tgggtgtgtt ttcacaggta aatcacctc 1320
tcanaaggta atgaagtata tcccttaaat gctgcagtt tattgtatgt aattataaac 1380
tcaatkatagg tgetttttaa aaaaacttga agtttagtt acaaacatat tgcaggttca 1440
ggaagccagg cactgtctca tctgtctggt gaaagcatga gctggtcaac ccatggagtg 1500
cagcttctga tttatctatc aaaaatcaaa atgcattgtc cctttgactc agccatttca 1560
ctttccagaa ttttagctta tccacattt gtttaattgt ctttaaatga agaaatggc 1620
tgttagcact tttgtaatag caactaaatg cccaccaata agaaatggc atacataaatt 1680
ctgtatcgtc catgtaatca atgcagaaag cagtgtggca agaatgagg gagctctttt 1740
agtattgaca cagaagctcc ctcaagacac tttaatgac taaggcaagg ggcctgacag 1800
tatgtagtat gctgaaaagg ggttaggaaa ggtgttatat attcaatatg cttattttgc 1860
atacaaaacty cctggaaagg tacataagaa attgcaataa gtgtttgtct totaaggaga 1920
atgggagctg cgaagtggaa gaaaggggga caagggaaaa agacactctt cactatgtac 1980
ttagsatctt ttatttttaa gccatgagaa tataatcaaa agtaaataaa tagaaatttt 2040
aaactaaagt aaagacagtc cttagattct tatgaagaaa actgtggaaa aataatcaga 2100
ctacacaaag attctaat tgtgaagatg cagaagttct ccatgggaca aaatgtakte 2160
atattaaagt tggattcagt catcttttgt ttgtctctt ttaaattea tgtttctaat 2220
agtaactgtg tttcttggaa ttaattggta aattattaaq tgggtgatcc catatacttt 2280
gtaaaaaaan aaaaaaa

```

2297

<210> 58  
 <211> 379  
 <212> PRT  
 <213> Homo sapiens

<400> 58

```

Met Ala Val Ile Gly Leu Pro Tyr Tyr Leu Phe Val Trp Glu Asp Tyr
  1                               5                               10          15
Asp Lys Tyr Val Ile Phe Ala Ser Phe Asn Leu Ile Trp Ser Thr Val
                20                25                30

```



Ile Leu Glu Leu Trp Lys Arg Gly Cys Ala Asn Met Thr Tyr Arg Trp  
 35 40 45  
 Gly Thr Leu Leu Met Lys Arg Lys Phe Glu Glu Pro Arg Pro Gly Phe  
 50 55 60  
 His Gly Val Leu Gly Ile Asn Ser Ile Thr Gly Lys Glu Glu Pro Leu  
 65 70 75 80  
 Tyr Pro Ser Tyr Lys Arg Gln Leu Arg Ile Tyr Leu Val Ser Leu Pro  
 85 90 95  
 Phe Val Cys Leu Cys Leu Tyr Phe Ser Leu Tyr Val Met Met Ile Tyr  
 100 105 110  
 Phe Asp Met Glu Val Trp Ala Leu Gly Leu His Glu Asn Ser Gly Ser  
 115 120 125  
 Gln Trp Thr Ser Val Leu Leu Tyr Val Pro Ser Ile Ile Tyr Ala Ile  
 130 135 140  
 Val Ile Glu Ile Met Asn Arg Leu Tyr Arg Tyr Ala Ala Glu Phe Leu  
 145 150 155 160  
 Thr Ser Trp Glu Asn His Arg Leu Glu Ser Ala Tyr Gln Asn His Leu  
 165 170 175  
 Ile Leu Lys Val Leu Val Phe Asn Phe Leu Asn Cys Phe Ala Ser Leu  
 180 185 190  
 Phe Tyr Ile Ala Phe Val Leu Lys Asp Met Lys Leu Leu Arg Gln Ser  
 195 200 205  
 Leu Ala Thr Leu Leu Ile Thr Ser Gln Ile Leu Asn Gln Ile Met Glu  
 210 215 220  
 Ser Phe Leu Pro Tyr Trp Leu Gln Arg Lys His Gly Val Gln Val Lys  
 225 230 235 240  
 Arg Lys Val Gln Ala Leu Lys Ala Asp Ile Asp Ala Thr Leu Tyr Glu  
 245 250 255  
 Gln Val Ile Leu Glu Lys Glu Met Gly Thr Tyr Leu Gly Thr Phe Asp  
 260 265 270  
 Asp Tyr Leu Glu Leu Phe Leu Gln Phe Gly Tyr Val Ser Leu Phe Ser  
 275 280 285  
 Cys Val Tyr Pro Leu Ala Ala Ala Phe Ala Val Leu Asn Asn Phe Thr  
 290 295 300  
 Glu Val Asn Ser Asp Ala Leu Lys Met Cys Arg Val Phe Lys Arg Pro  
 305 310 315 320  
 Phe Ser Glu Pro Ser Ala Asn Ile Gly Val Trp Gln Met Ile Phe Cys  
 325 330 335  
 Leu Asp Thr Gly Val Lys Arg Gly Leu Asn Cys Lys Val Met Arg Asn  
 340 345 350

Leu Leu Gly Glu Met Glu Met Ser Cys Val Leu Phe Val Val Val Val  
 355 360 365

Val Ser Gln Val Asn Thr Pro Ile Lys Arg  
 370 375

&lt;210&gt; 59

&lt;211&gt; 4145

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 59

```

aggtctcgaa ttcaagatga agtaagaaag gaaagagagg gtctggagaa tgaacttga 60
tctgtgaatt ttgaactgac aagcaagttt ttgaacgccc tggctcaaga tgggtgtgata 120
aatgaagag ctcttttckgt tactgaacta gatcgagttc atggaggtct tataactaaa 180
gtccagaaat ctctaaagaa acagagaggga ctctttaaaa atattcaggt ctccatcag 240
gaattttcaa aaatgaacaa ctctaatat gtactttggt gaacttctag cttaattgaa agttttgaa 300
aatktagata ctgcattatga caactttgtt gaacttctag gtccaggttc agaacaaatg cagtgtata 420
aagttttaca atgagttgac tgaatcctg gtccaggttc ctcttaaaag acttgcaaa aagcattgcc 480
gtttttgcaa ggaagacaga aagagatgaa ctcttaaaag ctctgctatc agtctcacc agcaggagg 540
agagaacctt gtgctctctc aatctctaca ctctgctatc agaacatgc cgtctctac gccccagccc 600
catgcacctt ctctccacc ctctgctatc tgtgcttcca gcaactcag ctctctctgc tactgtctca 660
ctccaggtg gggctgggac tgcctgacaa gtctcctcac aaacgcttg ctccagctct 720
ctctccaggt cgcagggacc acctctctcc acctctctcc gatctctggt gattgtcaa 780
atgcccattg ccatgggcta taatctctat gcttatggcc aghataatct gccatctca 840
ccagtgtatc accagagttc tggacaggct ccatactcgg gcccccagca gctctctca 900
cccttctctc agcccccca gcagtctctc tatccacagc agtaatatgt ctgctcagca 960
gtccagctga ttccagatcag agggaaagaa ataccaacc tgcantagt gtactaaact 1020
ctacgtctct gttaatgtca tgtactctcc tggactgaat gcagtgtata atttctgct 1080
acagctagaa gctgtgccc agttccaat aataggatt ttgattacac atgtgagatt tctgtctgt 1140
gcagtataaa cactaggtat agtgaacat agtgaacat ttgaacagat tatcttttc tacataagac 1260
aatgagaaat taaacctgca atgaacat ttacaaagat ggtttaagta tggatactag agaaaattaa 1320
atggttggga catcagatac ttacaaagat tttaatttcc attatgctat ttgcatat 1380
gtttttcttc tctttgttt atttttaaa taacttctc aagaacagtt gkcttgtta 1440
caaggcactg taactctctt tactgtctc attttttttc tggattatgt ctcttttgt 1500
tgktttgtta ttgattctca tactgttttt aagtgtatga atagtacaag ttattatcna 1560
atgtatataa gtttaacaga taactgtttt tatctactct tctctactat gtggaagaa caaacagttg 1620
ggatgtttta cagggaatc aaagaatct atctataat gtggaagaa caaacagttg 1680
gtanaagatt ttgacattt attttagaa ataaatctgt ttaactatct ttgcatcttt 1740
ctaccaaga ctcttcaaat aaacatacaa ataaatctgt ttaactatct ttgcatcttt 1800
agctctctca gaaatctgat gaaatctctg gtaataatc ctctcataatg kaatggaaat gtttgggaa 1860
ctggttaaaa taactacgac atttgcaat ctctcataatg ctctactgtg ttaactatct ttgcatcttt 1920
cacttactgt acctctctat ctttttctca ttaatttttt ttctctaaa gattggagta 1980
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2040
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2100
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2160
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2220
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2280
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2340
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2400
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2460
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2520
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2580
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2640
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2700
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2760
gtctatgtat taatactaat ctctgcatc agaatatctc atctctgtat aaactctagt 2820

```

```

tttttagccca tttcccatgc tgggcoctgt cacagagcca caggagagac attcagaaac 2880
taggaaagga gggcccccaca gctgatactg ccacagacac cctgactcac toggctctgt 2940
tagtgtaacc tttaaatgt agcaacacaa accctttccc tcttgctcag tcaactatcc 3000
cttggtttct ttttaactac ctgtgtctgg gcacagacaa tcaataaaa tgcagccctt 3060
tattactgtt aaggatcata ctgttggttt ggcgttgaaa ggttactact ctgtgattca 3120
ggtgtgttgt acccatatct ataattagga ttatttatct tctaatcca aggaaggaa 3180
atcatcccca gacctttat gctgagcttt ggcatactat tttaactgg attgtactta 3240
aatatgaag ctctgcatag aggaactagt cagaagtggg gaatacactg tctaattttt 3300
atcagtcctg tataaagtat tgatotaaga gaactctccc tgtgcccctt ggtctttatt 3360
ctcaattaa gaaacacagc acatgtcacg acaaacaaat caatctttat gagatattcc 3420
tgtatccata ccccgcttg ttgcaattt ataaacctcc ccttcaaaac taaggagttg 3480
cagaaaaaaa tggatttcac agagccttgc gtccctaaag ttctgtccc gtacgcagtc 3540
tttatagtc aaacagatta taaaaaatgt ttccatttg aactttacag ttgcgaag 3600
tgtttttata ctttttctaa ttccagaaac aggataattt gtaagtggg ttccagttt 3660
ctaataggga ttttttgtgt ttgtttttt aattttcagc atctcttga gaattttgt 3720
aacgcaaat ggcactctcc ttttaaaaga cgtttgcaat tattagtga ttccagttac 3780
agaacaaggt ataaaggaaa aaacccctgt aggtagtgtt ataattgcta gattcaaat 3840
agactagaa aggttcattt taagatttac ttggaagagc aaagaaaggaa aaatlattt 3900
tttaagaaa gagaatttc aggccttact ctggtatga agtttatatt ttttaaaaa 3960
atctatatt atcacaccag agattttaga ttctttctg gttagaaca ttgctggtag 4020
ttggattata tttttattgt attcatttt cttaggggga acattgtaaa gaaacaaaa 4080
ggtccagatg aatgtatgt agaaataaaa gttgaagat tcttacttca aaaaaaaa 4140
aaaaa 4145

```

&lt;210&gt; 60

&lt;211&gt; 289

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

```

Met Thr Ser Lys Phe Leu Thr Ala Leu Ala Gln Asp Gly Val Ile Asn
  1             5             10             15

Glu Glu Ala Leu Ser Val Thr Glu Leu Asp Arg Val Tyr Gly Gly Leu
      20             25             30

Thr Thr Lys Val Gln Glu Ser Leu Lys Lys Gln Glu Gly Leu Leu Lys
      35             40             45

Asn Ile Gln Val Ser His Gln Glu Phe Ser Lys Met Lys Gln Ser Asn
      50             55             60

Asn Glu Ala Asn Leu Arg Glu Glu Val Leu Lys Asn Leu Ala Thr Ala
      65             70             75             80

Tyr Asp Asn Phe Val Glu Leu Val Ala Asn Leu Lys Glu Gly Thr Lys
      85             90             95

Phe Tyr Asn Glu Leu Thr Glu Ile Leu Val Arg Phe Gln Asn Lys Cys
      100            105            110

Ser Asp Ile Val Phe Ala Arg Lys Thr Glu Arg Asp Glu Leu Leu Lys
      115            120            125

Asp Leu Gln Gln Ser Ile Ala Arg Glu Pro Ser Ala Pro Ser Ile Pro
      130            135            140

Thr Pro Ala Tyr Gln Ser Ser Pro Ala Gly Gly His Ala Pro Thr Pro
      145            150            155            160

```

Pro Thr Pro Ala Pro Arg Thr Met Pro Pro Thr Lys Pro Gln Pro Pro  
 165 170 175

Ala Arg Pro Pro Pro Val Leu Pro Ala Asn Arg Ala Pro Ser Ala  
 180 185 190

Thr Ala Pro Ser Pro Val Gly Ala Gly Thr Ala Ala Pro Ala Pro Ser  
 195 200 205

Gln Thr Pro Gly Ser Ala Pro Pro Pro Gln Ala Gln Gly Pro Pro Tyr  
 210 215 220

Pro Thr Tyr Pro Gly Tyr Pro Gly Tyr Cys Gln Met Pro Met Pro Met  
 225 230 235 240

Gly Tyr Asn Pro Tyr Ala Tyr Gly Gln Tyr Asn Met Pro Tyr Pro Pro  
 245 250 255

Val Tyr His Gln Ser Pro Gly Gln Ala Pro Tyr Pro Gly Pro Gln Gln  
 260 265 270

Pro Ser Tyr Pro Phe Pro Gln Pro Pro Gln Gln Ser Tyr Tyr Pro Gln  
 275 280 285

Gln

<210> 61  
 <211> 1417  
 <212> DNA  
 <213> Homo sapiens

<400> 61  
 ggtgcccagc atggcgagtg tagtgctgcc gagcggatcc cagtctgcga cggcagcggc 60  
 ggcggcgagg cctcccgggc tccggctcag gcttctgctg ttgtctttct ccgcccgggc 120  
 actgatcccc acaggtgatg ggcggaatct gcttcggaan gacgtgacag tgatcgaggg 180  
 agaggttgag accatcagtt gccaaagtcaa taagagtgac gactctgtga ttcagctact 240  
 gaatcccaac aggcagacca tttatttcag ggaacttcag cctttgaagg atagcaggtt 300  
 tcagttgctg aatttttcta gcagtgcaat caaggtatca ttgcacaaag tctcaatttc 360  
 tgatgaagga agatactttt gccagctcta taccgatccc cccagggaaa gttacaccac 420  
 catcacagtc ctggccccc cactgaatct gatgatgat atccagaaag acactgcggt 480  
 ggaaggtgag gagattgaa tcaactgcac tcttatggc agcagccag ccacagctat 540  
 caggtggttc aaagggaaac cagagctaaa aggcgaatcg gaggtggang agtggtcaga 600  
 catgtacact gtgaccagtc agctgatgct gaaggtgcac aagcagatc atggggcccc 660  
 agtgcattgc caggtgagag accttggggt cactgggaac ctgcagaccc agcgggtatct 720  
 agaatcacag tataagcttc aagtgcacat kmgatgact tatctctctc aaggttcaac 780  
 ccgggaaggg gacgcgcttg agttaacatg tgaagccatc ggggaagcccc agcctgtgat 840  
 ggttaacttg gtgagagtcg atgatgaast gcttcaacac gccgtactgt ctgggcccc 900  
 cctgttcctc aataacctaa acaaacaga taatggtaac taagcttccc gaggaggtga 1020  
 catagtgggg aagctcact cggattatat gctgtatgta tccgattccc gaggaggtga 1080  
 agcaggctcg atcaggagag tggatcatgc cgtgatcggt ggcgtcgttg cgttggtggt 1140  
 gtctgtcatg ctgtgcttgc tcatcattct gggcgctat tttgccagac ataaaggtac 1200  
 ctacttcact catgaagcca aaggagcccc kgacgcagca gacgtgagca cagctataat 1260  
 caatgcagaa ggaggacaga atcaactcga agaaagaaa gagtaactca tctagatcag 1320  
 cctttttggt tcaatgaggt gtccaactcg cctatttag atgataaaga gacagtgata 1380  
 ttggaacttg ccagaaattc gtgtgttttt ttatgaatgg gtggaaaggt gtgagactgg 1417  
 gaagucttgg gatttgcctg gtaaaaaaaa aaaaaaa

&lt;210&gt; 62

&lt;211&gt; 414

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 62

```

Met Ala Ser Val Val Leu Pro Ser Gly Ser Gln Cys Ala Ala Ala Ala
 1              5              10              15

Ala Ala Ala Ala Pro Pro Gly Leu Arg Leu Arg Leu Leu Leu Leu
 20              25              30

Phe Ser Ala Ala Ala Leu Ile Pro Thr Gly Asp Gly Gln Asn Leu Phe
 35              40              45

Thr Lys Asp Val Thr Val Ile Glu Gly Glu Val Ala Thr Ile Ser Cys
 50              55              60

Gln Val Asn Lys Ser Asp Asp Ser Val Ile Gln Leu Leu Asn Pro Asn
 65              70              75              80

Arg Gln Thr Ile Tyr Phe Arg Asp Phe Arg Pro Leu Lys Asp Ser Arg
 85              90              95

Phe Gln Leu Leu Asn Phe Ser Ser Ser Glu Leu Lys Val Ser Leu Thr
100              105              110

Asn Val Ser Ile Ser Asp Glu Gly Arg Tyr Phe Cys Gln Leu Tyr Thr
115              120              125

Asp Pro Pro Gln Glu Ser Tyr Thr Thr Ile Thr Val Leu Val Pro Pro
130              135              140

Arg Asn Leu Met Ile Asp Ile Gln Lys Asp Thr Ala Val Glu Gly Glu
145              150              155              160

Glu Ile Glu Val Asn Cys Thr Ala Met Ala Ser Lys Pro Ala Thr Thr
165              170              175

Ile Arg Trp Phe Lys Gly Asn Thr Glu Leu Lys Gly Lys Ser Glu Val
180              185              190

Glu Glu Trp Ser Asp Met Tyr Thr Val Thr Ser Gln Leu Met Leu Lys
195              200              205

Val His Lys Glu Asp Asp Gly Val Pro Val Ile Cys Gln Val Glu His
210              215              220

Pro Ala Val Thr Gly Asn Leu Gln Thr Gln Arg Tyr Leu Glu Val Gln
225              230              235              240

Tyr Lys Pro Gln Val His Ile Gln Met Thr Tyr Pro Leu Gln Gly Leu
245              250              255

Thr Arg Glu Gly Asp Ala Leu Glu Leu Thr Cys Glu Ala Ile Gly Lys
260              265              270

Pro Gln Pro Val Met Val Thr Trp Val Arg Val Asp Asp Glu Met Pro
275              280              285

Gln His Ala Val Leu Ser Gly Pro Asn Leu Phe Ile Asn Asn Leu Asn

```

290	295	300
Lys Thr Asp Asn Gly Thr Tyr Arg Cys Glu Ala Ser Asn Ile Val Gly		
305	310	315 320
Lys Ala His Ser Asp Tyr Met Leu Tyr Val Tyr Asp Ser Arg Ala Gly		
	325	330 335
Glu Glu Gly Ser Ile Arg Ala Val Asp His Ala Val Ile Gly Gly Val		
	340	345 350
Val Ala Val Val Val Phe Ala Met Leu Cys Leu Leu Ile Ile Leu Gly		
	355	360 365
Arg Tyr Phe Ala Arg His Lys Gly Thr Tyr Phe Thr His Glu Ala Lys		
	370	375 380
Gly Ala Asp Asp Ala Ala Asp Ala Asp Thr Ala Ile Ile Asn Ala Glu		
	385	390 395 400
Gly Gly Gln Asn Asn Ser Glu Glu Lys Lys Glu Tyr Phe Ile		
	405	410

<210> 63  
 <211> 1571  
 <212> DNA  
 <213> Homo sapiens

<400> 63  
 ggccggggag actgcgaccc ttttctctca gtctgcctta ctaccatgcc gctctacgag 60  
 ggccctggga gggcgggga gaagacggcg gtctgtgacg acctgggaga ggcctttacc 120  
 aagtgtggat ttgctggaga aactgggtca agatgtataa ttctagtgt gataaaaaga 180  
 gctgggatgc ctaagcctgt cagaattgtt cagtataata tcaatacaga agaattatat 240  
 tcttacctaa aggaattcat ccacatacta tatttcagga atctattggt gaatccaga 300  
 gaccgcccag ttgtgattat cgaatcggta ttatgtcctt ctcaattcag agagacactc 360  
 actcgtgttc ttttcaata ttttgaggtt ccactgtctt tgcctgtctc aagtcactct 420  
 ctggctcttc tgacgcttgg aatttattct gccatgtctc tagattgtgg atataggga 480  
 agcttgggtt tcccatata cgaaggatc ccaattctaa attgttggg agcactacc 540  
 ctaggaggga agctcttca caagcagttg gaaactcacc tattgggaca atgtactgtt 600  
 gacacagctg ttgctaaaga acagagcctt ccttcagtga tgggttcagt tccgggaagt 660  
 gtcttagagg acatttaagg gcgtacttgc tttgtaagtg atctgaagcg aggaactaaa 720  
 atccaaggcg caaatktta tttgtatggg aataatggcg gtccctccc acccccaaat 780  
 gttgactatc cattagatgg agagaaattt ttacatatcc ttggatcaat cagagattca 840  
 gtltgtggaan ttctttttga accagataat gaagagcaat cagttgcac ttaatatltg 900  
 gattccctta tacagtgtcc gatagacacc aggaagcaac tagcagagaa ttltgtgtc 960  
 ataggtggca ctcttatgtt gccaggattt ctccacagat tgcctgcaga aataaggtat 1020  
 ttggtagaan aaccaaaata taanaagca ctltgcacta agacatttcg aattcatact 1080  
 ccacctgcaa aagctaatlg tgtggcctgg ctgggagggg ctatttttgg agcattaca 1140  
 gatatacttg ggagccgttc tpttcaaaag gaatattata atcagacggg cegtatacct 1200  
 gattgggtgt ctctcaataa ccacactltg gaatgtatgt ttgatgtcgg gaaaactcaa 1260  
 ccacctctga tgaagagagc atttccact gagaactaga agtttgatta aaatcaacc 1320  
 ttgcttcata tcaaatattt aaccaattat aagcaattg tacaaagtat gtaggatgtt 1380  
 ttgttataga ggaactatgt ggaagtgaat gcattctgtt cttaactctt gcattaatat 1440  
 ataattcttt tgactttgtt ctctttgtgt agtggtaaaa tggtagcttg tgcattattga 1500  
 gatttgcctgt atttatatca ataaaglate gttaagcaaa aaaaaaaa 1560  
 aaaaaaaa a 1571

<210> 64  
 <211> 417

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 64

```

Met Pro Leu Tyr Glu Gly Leu Gly Ser Gly Gly Glu Lys Thr Ala Val
 1             5             10             15

Val Ile Asp Leu Gly Glu Ala Phe Thr Lys Cys Gly Phe Ala Gly Glu
      20             25             30

Thr Gly Pro Arg Cys Ile Ile Pro Ser Val Ile Lys Arg Ala Gly Met
      35             40             45

Pro Lys Pro Val Arg Val Val Gln Tyr Asn Ile Asn Thr Glu Glu Leu
      50             55             60

Tyr Ser Tyr Leu Lys Glu Phe Ile His Ile Leu Tyr Phe Arg His Leu
      65             70             75             80

Leu Val Asn Pro Arg Asp Arg Arg Val Val Ile Ile Glu Ser Val Leu
      85             90             95

Cys Pro Ser His Phe Arg Glu Thr Leu Thr Arg Val Leu Phe Lys Tyr
      100            105            110

Phe Glu Val Pro Ser Val Leu Leu Ala Pro Ser His Leu Met Ala Leu
      115            120            125

Leu Thr Leu Gly Ile Asn Ser Ala Met Val Leu Asp Cys Gly Tyr Arg
      130            135            140

Glu Ser Leu Val Leu Pro Ile Tyr Glu Gly Ile Pro Val Leu Asn Cys
      145            150            155            160

Trp Gly Ala Leu Pro Leu Gly Gly Lys Ala Leu His Lys Glu Leu Glu
      165            170            175

Thr Gln Leu Leu Glu Gln Cys Thr Val Asp Thr Ser Val Ala Lys Glu
      180            185            190

Gln Ser Leu Pro Ser Val Met Gly Ser Val Pro Glu Gly Val Leu Glu
      195            200            205

Asp Ile Lys Ala Arg Thr Cys Phe Val Ser Asp Leu Lys Arg Gly Leu
      210            215            220

Lys Ile Gln Ala Ala Lys Phe Asn Ile Asp Gly Asn Asn Glu Arg Pro
      225            230            235            240

Ser Pro Pro Pro Asn Val Asp Tyr Pro Leu Asp Gly Glu Lys Ile Leu
      245            250            255

His Ile Leu Gly Ser Ile Arg Asp Ser Val Val Glu Ile Leu Phe Glu
      260            265            270

Gln Asp Asn Glu Glu Gln Ser Val Ala Thr Leu Ile Leu Asp Ser Leu
      275            280            285

Ile Gln Cys Pro Ile Asp Thr Arg Lys Gln Leu Ala Glu Asn Leu Val
      290            295            300

```

Val Ile Gly Gly Thr Ser Met Leu Pro Gly Phe Leu His Arg Leu Leu  
305 310 315 320

Ala Glu Ile Arg Tyr Leu Val Glu Lys Pro Lys Tyr Lys Lys Ala Leu  
325 330 335

Gly Thr Lys Thr Phe Arg Ile His Thr Pro Pro Ala Lys Ala Asn Cys  
340 345 350

Val Ala Trp Leu Gly Gly Ala Ile Phe Gly Ala Leu Glu Asp Ile Leu  
355 360 365

Gly Ser Arg Ser Val Ser Lys Glu Tyr Tyr Asn Glu Thr Gly Arg Ile  
370 375 380

Pro Asp Trp Cys Ser Leu Asn Asn Pro Pro Leu Glu Met Met Phe Asp  
385 390 395 400

Val Gly Lys Thr Glu Pro Pro Leu Met Lys Arg Ala Phe Ser Thr Glu  
405 410 415

Lys

<210> 65  
<211> 1752  
<212> DNA  
<213> Homo sapiens

<400> 65  
ggcgaatcag agggacggcc ccagaatggc atggtagatg gaacgcagct gagaggtctg 60  
acaagatgta ccaggtccca ctaccactgg atcgggatgg gacctggta cggctccgct 120  
tcccaatggc ggccttggtc acgggtctgt gtccacttgt cgccttcttc ttctgcattc 180  
tctggctccct gctcttccac tccaaggaga caacggccat acactgtggg gtgcaccaat 240  
acctgccttc ggtgaagctc gccatcggcg gggagggtgc ccagcgctac gtgtggcggt 300  
tctgcctcgg cctgcactcg gcgcctcgct tcttggtggc ctgcgcctac tggaaacct 360  
acctcagctg caactccctg tgttctgtgt atcgcgcgt ctgcgccttc aacttcggcc 420  
tcaatgttgt gggaacctc ggttgcctag tgcctactta tgtctctctc tccgaggact 480  
tcaacctcca cgaacatgct tcaattgtgt tcaattgtgt atcctcctc atcctcctgg 540  
tcaactgcat tctctggcgg ttgactaaga agcacacagt aagtcaggag gatcgcaagt 600  
cctacagctg gaacacagcg ctcttcatca tcaacttcat ctcttcttcc tggcgctgg 660  
ctgtctactt tgggcacaa atgtatttgg agctcggagt gtacacccac ttgtctatcc 720  
tggagtacac tgttgtctta accaacatgg cgttccacat gacggcctgg tgggacttcc 780  
ggacacagga gctgtctata acctctcagg ctgaggaaaa gcgattctga accttcagt 840  
cctgcttggg aggaagcagg ccactgcctc gaacaaagaa acacgatacc attctggcch 900  
tcccaacccc acatcctctc ttggccttac tgaagatggg ggaagggtaa gaaggagg 960  
tgtaggccaa ggtcaccccc agtgtgtgtg gttctctctc tccacccctc atatggcgt 1020  
gggtctctca aacatcactt tcaactgaga ggcctcaaga agctgagctg gcagagagct 1080  
ccaccatttg gtgtcaaaa aaaaaacgtc ctgaggttca tgaccacat ccagtttctg 1140  
gcctttacac agtcaccttt caactgaggtc aggggcctct gaggagtggc tgcctccctg 1200  
caaccacagg cctttctctg caccgggggtc attcatagga ctatgtatt tcatgatata 1260  
ctgtgcacat ccaggcctgt ggcacacgtc cctgtctaaa gttgtctagg tgttctagtc 1320  
ctgacttcac ctttttgatt tgggtgtgtc cctagggtat gtacccttcc ccatctgagc 1380  
ctcgtgtgtc ccatgtgtct ggcgggggat ggggtgactg tatgatttcc aaggactcta 1440  
ccagtcaagt gttctgatgt catcgggtgg aggtgtgtct ctatacctaa aggatgacct 1500  
gctccagaaa caggaccagg acagcatgta tttctctctc ttctgaaagt tctgcttctg 1560  
agaccctccc cctcctttgc aagggtatgg atagaggggg tcagatgcag ctctctactg 1620  
taaatggggc tccctggtat ctctgtctct cctactgct caaaccccta aatttttggt 1680



gtacatttta ttggaaggga aaataaattt tttttttggg ccaaaaaaaa aaaaaaaa 1740  
 aaaaaaaa aa 1752

<210> 66  
 <211> 254  
 <212> PRT  
 <213> Homo sapiens

<400> 66  
 Met Tyr Gln Val Pro Leu Pro Leu Asp Arg Asp Gly Thr Leu Val Arg  
 1 5 10 15  
 Leu Arg Phe Thr Met Val Ala Leu Val Thr Val Cys Cys Pro Leu Val  
 20 25 30  
 Ala Phe Leu Phe Cys Ile Leu Trp Ser Leu Leu Phe His Phe Lys Glu  
 35 40 45  
 Thr Thr Ala Thr His Cys Gly Val Pro Asn Tyr Leu Pro Ser Val Ser  
 50 55 60  
 Ser Ala Ile Gly Gly Glu Val Pro Gln Arg Tyr Val Trp Arg Phe Cys  
 65 70 75 80  
 Ile Gly Leu His Ser Ala Pro Arg Phe Leu Val Ala Phe Ala Tyr Trp  
 85 90 95  
 Asn His Tyr Leu Ser Cys Thr Ser Pro Cys Ser Cys Tyr Arg Pro Leu  
 100 105 110  
 Cys Arg Leu Asn Phe Gly Leu Asn Val Val Glu Asn Leu Ala Leu Leu  
 115 120 125  
 Val Leu Thr Tyr Val Ser Ser Ser Glu Asp Phe Thr Ile His Glu Asn  
 130 135 140  
 Ala Phe Ile Val Phe Ile Ala Ser Ser Leu Gly His Met Leu Leu Thr  
 145 150 155 160  
 Cys Ile Leu Trp Arg Leu Thr Lys Lys His Thr Val Ser Gln Glu Asp  
 165 170 175  
 Arg Lys Ser Tyr Ser Trp Lys Gln Arg Leu Phe Ile Ile Asn Phe Ile  
 180 185 190  
 Ser Phe Phe Ser Ala Leu Ala Val Tyr Phe Arg His Asn Met Tyr Cys  
 195 200 205  
 Glu Ala Gly Val Tyr Thr Ile Phe Ala Ile Leu Glu Tyr Thr Val Val  
 210 215 220  
 Leu Thr Asn Met Ala Phe His Met Thr Ala Trp Trp Asp Phe Gly Asn  
 225 230 235 240  
 Lys Glu Leu Leu Ile Thr Ser Gln Pro Glu Glu Lys Arg Phe  
 245 250

<210> 67  
 <211> 782

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 67

```

cactctctgca gacaaggacac tgattgcccc agaccatgta gttccagctc cagaagagtg 60
ctatgtgtat agtccattgg gctctgctta taaccttcaa agttaccctg aaggataccg 120
taaaacacac agtttagtaa ccttttttat gatttggaa accatgatgg aaccatctat 180
actaagcatt ccttggggca taaaacagge tggctttact actgggaatgt gtgtcatcat 240
actgatgggc cttttaacac ttctattgctg ctacagagta gtgaaatcac ggactatgat 300
gttttcattg gataccacta cctgggaata tcccgatgtc tgcagacatt atttcggctc 360
ctttgggcag tggtcgagtc tctctctctc cttgggtctc ctcatggag caatgatagt 420
ttattgggtg cttatgtcaa atttctttt taatactgga aagtttattt ttagtaagta 480
tctatatcat atgcttttaa ccagtaact tcaatacta ttaccactgt aatgttagtt 540
ctagccitaa attctaggac ttgggataaa taataaaga agtaacatat ataattttgg 600
aaatatatc ttattcagtc ggctttctgt gtttgtgctc tcaatatag tgtatgctta 660
tttccaaaca ttaattcttg aaggaaatc atctctccaa atcttttagt taatatanaa 720
tatgtctata atccaaaaa aaaaaaana aaaaaaana aaaaaaana aaaaaaanaa 780
8

```

781

&lt;210&gt; 68

&lt;211&gt; 127

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

```

Met Ile Trp Asn Thr Met Met Gly Thr Ser Ile Leu Ser Ile Pro Trp
  1             5             10             15
Gly Ile Lys Gln Ala Gly Phe Thr Thr Gly Met Cys Val Ile Ile Leu
          20             25             30
Met Gly Leu Leu Thr Leu Tyr Cys Cys Tyr Arg Val Val Lys Ser Arg
          35             40             45
Thr Met Met Phe Ser Leu Asp Thr Thr Thr Trp Glu Tyr Pro Asp Val
          50             55             60
Cys Arg His Tyr Phe Gly Ser Phe Gly Gln Trp Ser Ser Leu Leu Phe
          65             70             75             80
Ser Leu Val Ser Leu Ile Gly Ala Met Ile Val Tyr Trp Val Leu Met
          85             90             95
Ser Asn Phe Leu Phe Asn Thr Gly Lys Phe Ile Phe Ser Lys Tyr Leu
          100            105            110
Tyr His Met Leu Leu Thr Gln Tyr Phe Gln Ile Leu Leu Pro Leu
          115            120            125

```

&lt;210&gt; 69

&lt;211&gt; 649

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 69

```

gagcaactcc ctcccccata tctgtcacc atgtggagcc tgaatctgc cctggctctg 60
cttctgtgac taccctgcag ctatgccttc atgtctcttt ctctggagaa gaaaactagc 120
gaaccacagg ggaaggtgca atcaggagag cacttkagga ttccgcagaa tctccagag 180
cacacccaag gctggcttgg gagcaaatgg ctctggcttc ttttggttgt tgtgccttt 240

```

```

gtgatactgc agtggtcaag agacagtgag aagaataagg agcagaagtc tectggccct 300
cgggggggac aacttcactc tccattaaa gaaaaangaa atgcttcccc caacaagac 360
tgtgcattca atacttaac ggaactegag gtggagctta tgaatttgt gtccaaagt 420
cggaaatetta aacgtgacat ggaacaggtt agtggcagta accdcaggt tcgaagta 480
gagatgcttg tagatccata ccatgtcaag atctgtgaaa tatggggaga agaaagctct 540
agctgaatgg atttgttgtt caggagagaa aaaaagttag tgttgacaaa ctgtatgaa 600
actaataaa ctattctgaa gaaaagaaa aaaaaaaa aaaaaaaa 649

```

&lt;210&gt; 70

&lt;211&gt; 171

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 70

```

Met Trp Thr Leu Lys Ser Ser Leu Val Leu Leu Leu Cys Leu Thr Cys
  1              5              10              15

Ser Tyr Ala Phe Met Phe Ser Ser Leu Arg Gln Lys Thr Ser Glu Pro
          20              25              30

Gln Gly Lys Val Gln Tyr Gly Glu His Phe Arg Ile Arg Gln Asn Leu
          35              40              45

Pro Glu His Thr Gln Gly Trp Leu Gly Ser Lys Trp Leu Trp Leu Leu
  50              55              60

Phe Val Val Val Pro Phe Val Ile Leu Gln Cys Gln Arg Asp Ser Glu
  65              70              75              80

Lys Asn Lys Glu Gln Ser Pro Pro Gly Leu Arg Gly Gly Gln Leu His
          85              90              95

Ser Pro Leu Lys Lys Lys Arg Asn Ala Ser Pro Asn Lys Asp Cys Ala
          100              105              110

Phe Asn Thr Leu Met Glu Leu Glu Val Glu Leu Met Lys Phe Val Ser
          115              120              125

Lys Val Arg Asn Leu Lys Arg Ala Met Ala Thr Gly Ser Gly Ser Asn
          130              135              140

Leu Arg Leu Arg Lys Ser Glu Met Pro Ala Asp Pro Tyr His Val Thr
          145              150              155              160

Ile Cys Glu Ile Trp Gly Glu Glu Ser Ser Ser
          165              170

```

&lt;210&gt; 71

&lt;211&gt; 1456

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

```

cacggctgtc ttatttgaa gtgcagagag gctctgctt cagctgggac accaatcttg 60
tcgggcaactc gtctcaccgt ggaatgtcat cactataaaa atgagggagag ctggcacgat 120
gggtgcccggg aatgtactctg tctcaatgga cgggaaatgt gtgacctgat caactgccc 180
gtgcttgccct gtggcaaccc ccaattcac cctggacagt gctgcccata atgtgcagat 240
gactttgttg tgcagaagcc agagctcagt actccctcaa ttgcccacgt cctggagga 300
gaataactttg tggagaggaga aacgtggaac attgaactct gtactcagtg cacttgccac 360

```

```

agcggagcggg tgcctgtgtga gacagaggtg tgcacacgcg tgcctctgcca gaacccctca 420
cgcaaccagg attcctgctg cccacagtgt acagatcaac cttctcgccg ttccctgtcc 480
cgcaataaca ggcctaccta ttattgcaaa atgatatgaq gggatatatt cctggcagct 540
gagtcctgga agcctgacgt ttgtaccagg tgcctctgca ttgatagcgt aattagctgt 600
ttctctgagt cctgccttcc tgtatcctgt gaagacctg tcttgagaaa aggcagtggt 660
tgtccctact gcatagaaga cacaattcca aagaagggtg tgtcccaact cagtgggaag 720
gcctatgccc acgaggagcg gtgggacctt gacagctgca cctactgata ctgcttgcag 780
ggcagacccc tctgctcgac cgtcagctgc cccctctgca cctgtgttga gcccatcaac 840
gtgggaaggaa gttcctgccc aatgtgtcca gaattgtatg tcccagaacc aaccaatata 900
ccatttgaga agacaaacca tggaggagag gttagcctgg aggttccctt gtggcccccg 960
cctagtgaaa atgatatcgt ccatctccct agagatatgg gtacactcca ggtagattac 1020
agagataaca ggcctgaccc aagtgaagat tcttccctgg actccattgc ctcaattgtg 1080
gttcccataa tttatgtcct ctctattata atagcattac tattcatcaa tcagaagaaa 1140
cctgggatac cactgctttg ctggatctga acaccaacta agccttcttc cttaataaat 1200
cagctagtat ctgtggactg caagaagga accagagtc aggtggacag tcccagaga 1260
atgctaaaga ttgcagaacc agatgcaga ttcagtgct tctacagcat gcaaaacag 1320
aaccatctac aggcagacaa ttctaccaa acagctgtga gaaaggcanc taggatgag 1380
tttcaaaaga cgggaagcga ctcaatctgc tctaaaagt aaactagaa ttgtgcactt 1440
aaaaaaanaa aaaaaa

```

&lt;210&gt; 72

&lt;211&gt; 400

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 72

```

Met Cys Ala Leu Ile Thr Cys Pro Val Pro Ala Cys Gly Asn Pro Thr
  1             5             10             15

```

```

Ile His Pro Gly Gln Cys Cys Pro Ser Cys Ala Asp Asp Phe Val Val
      20             25             30

```

```

Gln Lys Pro Glu Leu Ser Thr Pro Ser Ile Cys His Ala Pro Gly Gly
      35             40             45

```

```

Glu Tyr Phe Val Glu Gly Glu Thr Trp Asn Ile Asp Ser Cys Thr Gln
      50             55             60

```

```

Cys Thr Cys His Ser Gly Arg Val Leu Cys Glu Thr Glu Val Cys Pro
      65             70             75             80

```

```

Pro Leu Leu Cys Gln Asn Pro Ser Arg Thr Gln Asp Ser Cys Cys Pro
      85             90             95

```

```

Gln Cys Thr Asp Gln Pro Phe Arg Pro Ser Leu Ser Arg Asn Asn Ser
     100             105             110

```

```

Val Pro Asn Tyr Cys Lys Asn Asp Glu Gly Asp Ile Phe Leu Ala Ala
     115             120             125

```

```

Glu Ser Trp Lys Pro Asp Val Cys Thr Ser Cys Ile Cys Ile Asp Ser
     130             135             140

```

```

Val Ile Ser Cys Phe Ser Glu Ser Cys Pro Ser Val Ser Cys Glu Arg
     145             150             155             160

```

```

Pro Val Leu Arg Lys Gly Gln Cys Cys Pro Tyr Cys Ile Glu Asp Thr
     165             170             175

```

```

Ile Pro Lys Lys Val Val Cys His Phe Ser Gly Lys Ala Tyr Ala Asp

```

180										185					190						
Glu	Glu	Arg	Trp	Asp	Leu	Asp	Ser	Cys	Thr	His	Cys	Tyr	Cys	Leu	Gln						
		295					300					205									
Gly	Gln	Thr	Leu	Cys	Ser	Thr	Val	Ser	Cys	Pro	Pro	Leu	Pro	Cys	Val						
	210					215					220										
Glu	Pro	Ile	Asn	Val	Glu	Gly	Ser	Cys	Cys	Pro	Met	Cys	Pro	Glu	Met						
225				230						235				240							
Tyr	Val	Pro	Glu	Pro	Thr	Asn	Ile	Pro	Ile	Glu	Lys	Thr	Asn	His	Arg						
				245					250					255							
Gly	Glu	Val	Asp	Leu	Glu	Val	Pro	Leu	Trp	Pro	Thr	Pro	Ser	Glu	Asn						
		260					265						270								
Asp	Ile	Val	His	Leu	Pro	Arg	Asp	Met	Gly	His	Leu	Gln	Val	Asp	Tyr						
	275						280					285									
Arg	Asp	Asn	Arg	Leu	His	Pro	Ser	Glu	Asp	Ser	Ser	Leu	Asp	Ser	Ile						
	290					295					300										
Ala	Ser	Val	Val	Val	Pro	Ile	Ile	Ile	Cys	Leu	Ser	Ile	Ile	Ile	Ala						
305				310					315						320						
Phe	Leu	Phe	Ile	Asn	Gln	Lys	Lys	Gln	Trp	Ile	Pro	Leu	Leu	Cys	Trp						
			325					330						335							
Tyr	Arg	Thr	Pro	Thr	Lys	Pro	Ser	Ser	Leu	Asn	Asn	Gln	Leu	Val	Ser						
			340					345					350								
Val	Asp	Cys	Lys	Lys	Gly	Thr	Arg	Val	Gln	Val	Asp	Ser	Ser	Gln	Arg						
	355						360					365									
Met	Leu	Arg	Ile	Ala	Glu	Pro	Asp	Ala	Arg	Phe	Ser	Gly	Phe	Tyr	Ser						
	370					375					380										
Met	Gln	Lys	Gln	Asn	His	Leu	Gln	Ala	Asp	Asn	Phe	Tyr	Gln	Thr	Val						
385				390					395					400							

&lt;210&gt; 73

&lt;211&gt; 4723

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 73

```

ggccttcacg ggcctatcttt tttttttttt aaatgataca acttaatttt attaggacaa 60
ggctgggtgg cactggagtg gcacttcacg ggcacaggaga ggcactgggg aggggtcaca 120
ggatgctact cgggcaccta gaagccacag ctgcccctcca cagagcggca ctgcactatg 180
cgacgggaatg tctcgacctt gtccatgtcc ttcttgagtc agtagagcag ccgtagtttc 240
ctgagcagtg cgtcatagtt gtgagagttt gtgtcacaact tgccttaggt ctgcttgagg 300
atctgtccag tccggcggtt gcctgtcttc agcctcccca tcaagcgttt gatgccttcc 360
tctaggtcct ctgagcggtg atagtcacag ttgtcagagg tgtcatacac caggttggtg 420
gcgaacatac tctgaggaa ccgtacgggc tccagccacg actcgatgag cagcagggag 480
atgaggagca gctctagatt ggatttctgt tgcgtttcct ccattgttga gggtgtcgga 540
atagagtcctg agaagcagaa ggaggtcttg gactcatgca ggaatgaata cttctggtcc 600
tttgagatat aggtttcttc aaactccttg taggtgtcaa tggccagctg gtgcgcgcga 660
tggtgttggg gcataagctg gtcaaaaagc ctggataacg gaacgggttg gatggaccca 720

```

gectotttga	gccagggcag	gcagagcagg	gcacacagcca	ggagcagcga	cgtccgggag	780
cctggagcca	ttgcacactag	gtgagctgtc	cacaggaccc	tgaatgggtc	ggggaggttc	840
gccttcactg	cctagggagc	gcgcaggagt	gagcagagtc	gggcggcgcg	agcggagctc	900
ggggatcttt	tgctgcgcga	cgcgcgcac	tcggcagctc	gggaggtcgg	gacgggcccc	960
gaggtctg	cgctgcgggg	cgggcggact	cggagggagga	gagggagggag	gcgcgcctgg	1020
cccggtctgg	agccgagcgc	agcagccacc	gcgcggcgcc	cgcagggagt	ttgggttgaa	1080
ccggagctgc	cgggagggaaa	cttttttttt	ttttccctct	ccctccgggg	aggagggagg	1140
ggagggaggg	gggaagctgc	cgcgcggcgc	agggctcgtg	ggctcggggg	cggcgggggc	1200
cgcagaaggg	gggggggctt	cgcgcggcga	ggggaggggc	gcgcgggggg	cccgaggagg	1260
ggcggttgag	cccgggggct	gctggttgcc	cggcgggggc	ggcggtgtgt	cccgcgcgag	1320
gggagggcgc	cgcgcgcgtt	cccgggcggg	ctgcagggag	gagggcgggg	cggcgcgagg	1380
ggatgtactt	gggggggggg	gacagggggg	tcggcggttg	cgggcgcctc	ctgggtctcg	1440
tgctgggggt	gctgctctct	ctggcgcgct	cgcgcacccg	ggcgctgggt	tgctctgctt	1500
gtgacggagc	caagtggcag	gagcccgagg	actgcccggg	gagcatcgtg	caggcgctct	1560
ggcgctgctg	ctacacgtgc	gcagggcaga	ggaaacgagag	ctgcggcggg	accttcggga	1620
ttlatcgga	ctgggacggg	gggtctgggt	gtgtcatccg	cccccgcctc	aatggcgact	1680
ccctcaccga	gtacgagcgc	gggttttggg	agatggagaa	ctggactggt	gacaaactgc	1740
ttggttttaa	accatgcaat	gaaaaactta	ttgttgggtg	caatataatc	aatgggaaat	1800
gtgaatgtaa	cacatctaga	acctgcagca	atccctttga	gtttccaggt	caggatatgt	1860
gectttcagc	tttaagagga	attgaagag	agaaagccaga	tgctccag	gcccgtctgt	1920
aggtccagtt	ctctccaggt	tgctctgaag	attctgttct	gatcgagggt	tatgctcttc	1980
ctggggagtg	ctgtccctta	cccgagcgct	gctgtgtcaa	cccgcgaggg	tgctctgcga	2040
aggtctgcca	gcccgggaac	ctgaacatac	tagtgtcaaa	agcctcaggg	agccggggag	2100
actgctgtgc	cctctatgag	tgcaaacctg	tttccggcgt	ggactgcagg	actgtggaa	2160
gectcctgt	tcagtagacc	gctgttcccc	cggcagctta	tgaaactcaa	gtcaggaata	2220
ctgcagctgg	ttgctgtact	ttgcacacaa	gatcgaggtg	ttctcttggt	ttatgtgttt	2280
cccccggtg	tgaggtggga	tcacatcccc	gcctagcttc	tcgtggcgt	gggacacctg	2340
gaaagtgtct	tgatgtcttt	gaatgtgtta	atgatacaaa	gcccgcctgc	gtatttaaca	2400
atgtggaaata	ttatgatgga	gacatgtttc	gaatggacaa	ctgtcggttc	tgctgatgac	2460
aaggggcctt	tgcaatctgc	ttcaatgcct	agtgtggtga	gataaactgc	gagaggtact	2520
acgtgcgcga	aggggagctc	tgcccaggtg	gtgaagatcc	agtgtatcct	tttaataatc	2580
cgcctggctg	ctatggcaat	ggcctgacct	ttgcacacgg	agaccggatg	cggggaagac	2640
actgcacatt	ctgcagctgc	gtcaacgggtg	aacgccactg	cgttgcgacc	gtctgcggac	2700
agacctgcac	aaacccctgt	aaagtgcctg	gggaggttat	ccctgtgtgc	gaagcaacaa	2760
ccatcatcac	agttgatcca	cctgcactgt	acaaatggtg	gggacctgt	ctgacaggga	2820
aggactgcct	taatgggttc	aaacggcgatc	aaggctgcac	cttgaaactgt	cagtgcataa	2880
aaacggagga	actatgttca	gaacgttaac	gtgagtgccg	cccaaggccc	cccttcgggtt	2940
tccttactga	tgcccaaaac	tgtagatctt	ttggattgct	gaagaaatag	agagagtcca	3000
gacccataat	ctgtgacag	tttgtctcac	ttccatgcag	taagatctgc	cacggctgtg	3060
acatctgtcg	ctgtacgaaa	tgctcttctc	gcaagtgcag	agagggcctt	cccttggggt	3120
tcagcgaggga	cagtcacggc	tgctcttctc	ccgtggatgg	taagatctgc	gcttcagctg	3180
ggccacccat	cctgtcgggc	actgtgtctc	actgtgtctc	agagggcctt	aaatgggg	3240
agagctggca	cgatgggtgc	cgggaatgct	actgtgtctc	tggaaggga	atgtgtgccc	3300
tgatcacctg	cccggtgctt	gctgtgtgca	accccaacat	tcacccctgg	cagtgtctgc	3360
cactatgtgc	agatgacttt	gtgtgtcaga	agccagggct	cagtactccc	tcctttggcc	3420
acgcccctgg	aggagatatc	tttgtggag	gagaaacgtg	gaacattgac	tcctgtactc	3480
agtgcacctg	ccacagcgga	cgggtgctgt	gtgagacaga	gggtgtccca	cgcctgtctc	3540
gccagaaacc	ctcagcgacc	caggattctt	gctgcccaca	gtgtacagat	caaccttttc	3600
ggccttccct	gtcccgcaat	taagagcttg	ctaattactg	caaaatgat	gaaggggata	3660
ttctcctggc	agctgagctc	ggtagagctg	acgtttgtac	cagctgcata	tgcatggata	3720
gcgttaattag	ctgtttctct	gggtcctgct	cttctgtatc	ctgtgaaaga	cctgtcttga	3780
gaagaggcca	gtgttgtccc	tactgcatag	agacacaaat	tcacagagag	gtggtgtgac	3840
acttcagctg	gaaggcctat	gcccagcagg	agcggtggga	ccttgacagc	tgaccccaat	3900
gctactgcct	gcaggggcag	acccctctgt	cgaacgttcg	ctgccccctt	ctgcctctgt	3960
ttgagcccat	caacgtggaa	ggaagtgtct	gcccattgtg	tcagaaatg	tatgtccag	4020
aaacaaacaa	tatcccaatt	gagaagacaa	accatcgagg	agaggttgac	ctggaggttc	4080
ccctgtggcc	cacgcctagt	gaatgtata	tcgtccatct	ccctagagat	atgggtcacc	4140
tcaggttaga	ctacagagct	aacaggctgc	acccagtgga	agattcttca	ctggactcca	4200
ttgcctcaggt	tggtgttccc	ataatttat	gctctcttat	tataatagca	ttcctattca	4260
tcactcagaa	gaacacgtgg	ataccactgc	ttgtctggtg	tcgaacccca	actaagcctt	4320

```

cttccctttaa taatcagttt gtatctgttg actccacaga aggaaccaga gtccaggttg 4380
acagttccca gagaatgcta agaattgcag aaccagatgc aagattcagt ggtttctaca 4440
gcattcaca acagaccat ctccaggcag acattttcta cccacacagt tgaagaaagg 4500
caactaggat gagttttcaa agacaggag agcactaat ctgtctctaa aagttaacta 4560
gaatttgtgc acttgcctag tggattgtat tggattgtga cttgatgtac agcgttaaga 4620
ccttactggg atggctcttg tctacagcaa tgtgcagAAC aagcattccc cctcaacct 4680
aaaaaaaaaa aaaaaaadaa aaaaaaadaa aaaaaaadaa aaa 4723

```

&lt;210&gt; 74

&lt;211&gt; 1036

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 74

```

Met Tyr Leu Val Ala Gly Asp Arg Gly Leu Ala Gly Cys Gly His Leu
  1              5              10              15

Leu Val Ser Leu Leu Gly Leu Leu Leu Leu Leu Ala Arg Ser Gly Thr
      20              25              30

Arg Ala Leu Val Cys Leu Pro Cys Asp Glu Ser Lys Cys Glu Glu Pro
      35              40              45

Arg Asn Cys Pro Gly Ser Ile Val Gln Gly Val Cys Gly Cys Cys Tyr
      50              55              60

Thr Cys Ala Ser Gln Arg Asn Glu Ser Cys Gly Gly Thr Phe Gly Ile
      65              70              75              80

Tyr Gly Thr Cys Asp Arg Gly Leu Arg Cys Val Ile Arg Pro Pro Leu
      85              90              95

Asn Gly Asp Ser Leu Thr Glu Tyr Glu Ala Gly Val Cys Glu Asp Glu
     100              105              110

Asn Trp Thr Asp Asp Gln Leu Leu Gly Phe Lys Pro Cys Asn Glu Asn
     115              120              125

Leu Ile Ala Gly Cys Asn Ile Ile Asn Gly Lys Cys Glu Cys Asn Thr
     130              135              140

Ile Arg Thr Cys Ser Asn Pro Phe Glu Phe Pro Ser Gln Asp Met Cys
     145              150              155              160

Leu Ser Ala Leu Lys Arg Ile Glu Glu Glu Lys Pro Asp Cys Ser Lys
     165              170              175

Ala Arg Cys Glu Val Gln Phe Ser Pro Arg Cys Pro Glu Asp Ser Val
     180              185              190

Leu Ile Glu Gly Tyr Ala Pro Pro Gly Glu Cys Cys Pro Leu Pro Ser
     195              200              205

Arg Cys Val Cys Asn Pro Ala Gly Cys Leu Arg Lys Val Cys Gln Pro
     210              215              220

Gly Asn Leu Asn Ile Leu Val Ser Lys Ala Ser Gly Lys Pro Gly Glu
     225              230              235              240

Cys Cys Asp Leu Tyr Glu Cys Lys Pro Val Phe Gly Val Asp Cys Arg

```

245	250	255
Thr Val Glu Cys Pro Pro Val Gln Gln Thr Ala Cys Pro Pro Asp Ser 260 265 270		
Tyr Glu Thr Gln Val Arg Leu Thr Ala Asp Gly Cys Cys Thr Leu Pro 275 280 285		
Thr Arg Cys Glu Cys Leu Ser Gly Leu Cys Gly Phe Pro Val Cys Glu 290 295 300		
Val Gly Ser Thr Pro Arg Ile Val Ser Arg Gly Asp Gly Thr Pro Gly 305 310 315 320		
Lys Cys Cys Asp Val Phe Glu Cys Val Asn Asp Thr Lys Pro Ala Cys 325 330 335		
Val Phe Asn Asn Val Glu Tyr Tyr Asp Gly Asp Met Phe Arg Met Asp 340 345 350		
Asn Cys Arg Phe Cys Arg Cys Gln Gly Gly Val Ala Ile Cys Phe Thr 355 360 365		
Ala Gln Cys Gly Glu Ile Asn Cys Glu Arg Tyr Tyr Val Pro Glu Gly 370 375 380		
Glu Cys Cys Pro Val Cys Glu Asp Pro Val Tyr Pro Phe Asn Asn Pro 385 390 395 400		
Ala Gly Cys Tyr Ala Asn Gly Leu Ile Leu Ala His Gly Asp Arg Trp 405 410 415		
Arg Glu Asp Asp Cys Thr Phe Cys Gln Cys Val Asn Gly Glu Arg His 420 425 430		
Cys Val Ala Thr Val Cys Gly Gln Thr Cys Thr Asn Pro Val Lys Val 435 440 445		
Pro Gly Glu Cys Cys Pro Val Cys Glu Glu Pro Thr Ile Ile Thr Val 450 455 460		
Asp Pro Pro Ala Cys Gly Glu Leu Ser Asn Cys Thr Leu Thr Gly Lys 465 470 475 480		
Asp Cys Ile Asn Gly Phe Lys Arg Asp His Asn Gly Cys Arg Thr Cys 485 490 495		
Gln Cys Ile Asn Thr Glu Glu Leu Cys Ser Glu Arg Lys Gln Gly Cys 500 505 510		
Thr Leu Asn Cys Pro Phe Gly Phe Leu Thr Asp Ala Gln Asn Cys Glu 515 520 525		
Ile Cys Glu Cys Arg Pro Arg Pro Lys Lys Cys Arg Pro Ile Ile Cys 530 535 540		
Asp Lys Tyr Cys Pro Leu Gly Leu Leu Lys Asn Lys His Gly Cys Asp 545 550 555 560		
Ile Cys Arg Cys Lys Lys Cys Pro Glu Leu Ser Cys Ser Lys Ile Cys		



565										570					575				
Pro	Leu	Gly	Phe	Gln	Gln	Asp	Ser	His	Gly	Cys	Leu	Ile	Cys	Lys	Cys				
			580					585					590						
Arg	Glu	Ala	Ser	Ala	Ser	Ala	Gly	Pro	Pro	Ile	Leu	Ser	Gly	Thr	Cys				
		595					600					605							
Leu	Thr	Val	Asp	Gly	His	His	His	Lys	Asn	Glu	Glu	Ser	Trp	His	Asp				
		610				615					620								
Gly	Cys	Arg	Glu	Cys	Tyr	Cys	Leu	Asn	Gly	Arg	Glu	Met	Cys	Ala	Leu				
		625			630						635				640				
Ile	Thr	Cys	Pro	Val	Pro	Ala	Cys	Gly	Asn	Pro	Thr	Ile	His	Pro	Gly				
			645						650					655					
Gln	Cys	Cys	Pro	Ser	Cys	Ala	Asp	Asp	Phe	Val	Val	Gln	Lys	Pro	Glu				
			660					665						670					
Leu	Ser	Thr	Pro	Ser	Ile	Cys	His	Ala	Pro	Gly	Gly	Glu	Tyr	Phe	Val				
		675					680						685						
Glu	Gly	Glu	Thr	Trp	Asn	Ile	Asp	Ser	Cys	Thr	Gln	Cys	Thr	Cys	His				
		690				695					700								
Ser	Gly	Arg	Val	Leu	Cys	Glu	Thr	Glu	Val	Cys	Pro	Pro	Leu	Leu	Cys				
		705			710				715					720					
Gln	Asn	Pro	Ser	Arg	Thr	Gln	Asp	Ser	Cys	Cys	Pro	Gln	Cys	Thr	Asp				
			725						730					735					
Gln	Pro	Phe	Arg	Pro	Ser	Leu	Ser	Arg	Asn	Asn	Ser	Val	Pro	Asn	Tyr				
		740						745					750						
Cys	Lys	Asn	Asp	Glu	Gly	Asp	Ile	Phe	Leu	Ala	Ala	Glu	Ser	Trp	Lys				
		755					760					765							
Pro	Asp	Val	Cys	Thr	Ser	Cys	Ile	Cys	Ile	Asp	Ser	Val	Ile	Ser	Cys				
		770				775					780								
Phe	Ser	Glu	Ser	Cys	Pro	Ser	Val	Ser	Cys	Glu	Arg	Pro	Val	Leu	Arg				
		785			790				795					800					
Lys	Gly	Gln	Cys	Cys	Pro	Tyr	Cys	Ile	Glu	Asp	Thr	Ile	Pro	Lys	Lys				
			805						810				815						
Val	Val	Cys	His	Phe	Ser	Gly	Lys	Ala	Tyr	Ala	Asp	Glu	Glu	Arg	Trp				
		820						825				830							
Asp	Leu	Asp	Ser	Cys	Thr	His	Cys	Tyr	Cys	Leu	Gln	Gly	Gln	Thr	Leu				
		835					840					845							
Cys	Ser	Thr	Val	Ser	Cys	Pro	Pro	Leu	Pro	Cys	Val	Glu	Pro	Ile	Asn				
		850				855					860								
Val	Glu	Gly	Ser	Cys	Cys	Pro	Met	Cys	Pro	Glu	Met	Tyr	Val	Pro	Glu				
		865			870						875			880					
Pro	Thr	Asn	Ile	Pro	Ile	Glu	Lys	Thr	Asn	His	Arg	Gly	Glu	Val	Asp				

70

```

ctaggccsaga cagacagtat tggatagggt aaagcaagaa aactcaagot gcagctggac 1500
tgcaggpette ttttgcftaa gtcaacagtg cctcaaacct ccaaacctca atgcagtcas 1560
ttattccagcg catgcacagc ataatttgcet cetttgtgtg gagtgggtgtg tcagcccttg 1620
aacatctctct ccaagagagac tggagagtc ttaaatata tgtgggagga ggggggatag 1680
aacatcacaa cactgcteta gtttcttggg gaatacatt tccttacagg ttaangacaa 1740
accagccccc egggttttta tctagaaagt cattcaagtg aaagaaagag aagggaattg 1800
cttagtagga gtctctcagt atagaacat tacttgtatg aaattatacc tctgaatttt 1860
agaatgtcat gtgtctcttt aaaaaatta gctccccatc ctccctctct actctctccc 1920
tcctctcttc tctctctctc tctctctccc tctctcaag acacatacac acacacacac 1980
acacacacag cgtccacac tcaatttaa cttaagcttt attggaagca aagttagcca 2040
aatctctacg ttaattttcc cttagctgga tcccaagtag ctgggaagtt tttgtgcca 2100
ggagagtcas taactgtgaa caagaggtc tgcctctagg tcttctgtgg tgtttaagtc 2160
accaacaata gactcagggt aaagaatasa aacacttcca tagctctctt cattcaccta 2220
gaagtggtaa taatttttcc cttaatgata cacttttctt ctccctctgt acctatggga 2280
cttcagaaa gaagtcaaat tgagtcaaat catcagaac tgaatccctg taagaaaaaa 2340
taattgttga agaagagagt tgatagant caaaaaggcc atctttttgc ttacagaca 2400
ataaaactta ccaagtaata gatcagtcct cttaaatatt tttagacca tagttgtctg 2460
gtcagaaaaa ttatatttaa ttagtaatt cttagagctc tttagcaagt ggtcgggtcc tcatgagcat 2520
tctctctcaa ttataggagt tgatttttac tttagcaagt ggtcgggtcc tcatgagcat 2580
ctgcatgttg actctctcgt taagaacttc gttgttctat tagggaggtg gatattctga 2640
tgaagatctt tctctcaaac ctctctctca tcttctgtct attcatcag cagatacttt 2700
agtcagaaat tccagagaaq gctgctctca aaatgtctac ttgcagcccc ataccagagc 2760
ataaacctac cattctgggg tctgctctta gaaatcatct ttgtgggaag acctactct 2820
tcacagcaag gatctcagga atgctctcta gatttcttcc ctctgagggg caggaatgaa 2880
ctgtagaaat gttttaagga cccagaaacc ccatatgtct cattccatga ctataggtga 2940
gagaattctt tcttaagagg gtttctatcc aataggggaa atgtataaak gttcagttct 3000
tatgacaccc tggcataaag gactcaattc ttatgaaga gacaaaaggc ccttatggcc 3060
agggtttctt gggacaagac tctcaacagc acatcacaca cgttctctct gtgtgtgtgt 3120
agcagtcacat cctgggttag aggtcacaaa gcttagtgtt gtgtgtgtgt ggtaaaggga 3180
gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt ggtaaaggga ggaaggtgtt 3240
atgagggtgc tctctcgtc cagaggtgtg cagtgtctcc ataatgtgga gactagttaac 3300
tagctcttaa ggcataagag tgttctctct tttggtgat tcaatccaaa gcttccccc 3360
ccaggtgttc tctgaagagt tagctttaag agaaacagca gaggatctcc ctgatatata 3420
tctgctctcc aggtgtgtgg acacaccttt gcaaatgct gtgggaagca ggaagctggg 3480
agctgtgtta agtcaagga gaacccctcc agtgtttgt gtgtgtgtga gaataggaca 3540
taggttaag aggcraagct gctgttagtt agtagagag aatggatgtg gttctctctg 3600
tgtatttatt tglatctaa acatttgaa caacaaagac cataagcatc atttagcagt 3660
tgtagtcatt ttctagttaa ctctgttaa caagtaagag taacataaca gtattactct 3720
tcaactgttc tcaaggaac tgaactaat catgctctt atttatgtag tcaactgtatt 3780
tctggaattt caaattaata aaaaagttta ttttgaaan tcaaaaanaa aaaaaaaa 3840
aaaaaaaana aaaaaaaa a 3861

```

&lt;210&gt; 76

&lt;211&gt; 457

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 76

```

Met Arg Lys Lys Trp Lys Met Gly Gly Met Lys Tyr Ile Phe Ser Leu
  1             5             10             15

```

```

Leu Phe Phe Leu Leu Leu Glu Gly Gly Lys Thr Glu Gln Val Lys His
          20             25             30

```

```

Ser Glu Thr Tyr Cys Met Phe Gln Asp Lys Lys Tyr Arg Val Gly Glu
          35             40             45

```

```

Arg Trp His Pro Tyr Leu Glu Pro Tyr Gly Leu Val Tyr Cys Val Asn
          50             55             60

```

Cys Ile Cys Ser Glu Asn Gly Asn Val Leu Cys Ser Arg Val Arg Cys  
 65 70 75 80  
 Pro Asn Val His Cys Leu Ser Pro Val His Ile Pro His Leu Cys Cys  
 85 90 95  
 Pro Arg Cys Pro Asp Ser Leu Pro Pro Val Asn Asn Lys Val Thr Ser  
 100 105 110  
 Lys Ser Cys Glu Tyr Asn Gly Thr Thr Tyr Gln His Gly Glu Leu Phe  
 115 120 125  
 Val Ala Glu Gly Leu Phe Gln Asn Arg Gln Pro Asn Glu Cys Thr Gln  
 130 135 140  
 Cys Ser Cys Ser Glu Gly Asn Val Tyr Cys Gly Leu Lys Thr Cys Pro  
 145 150 155 160  
 Lys Leu Thr Cys Ala Phe Pro Val Ser Val Pro Asp Ser Cys Cys Arg  
 165 170 175  
 Val Cys Arg Gly Asp Gly Glu Leu Ser Trp Glu His Ser Asp Gly Asp  
 180 185 190  
 Ile Phe Arg Gln Pro Ala Asn Arg Glu Ala Arg His Ser Tyr His Arg  
 195 200 205  
 Ser His Tyr Asp Pro Pro Pro Ser Arg Gln Ala Gly Gly Leu Ser Arg  
 210 215 220  
 Phe Pro Gly Ala Arg Ser His Arg Gly Ala Leu Met Asp Ser Gln Gln  
 225 230 235 240  
 Ala Ser Gly Thr Ile Val Gln Ile Val Ile Asn Asn Lys His Lys His  
 245 250 255  
 Gly Gln Val Cys Val Ser Asn Gly Lys Thr Tyr Ser His Gly Glu Ser  
 260 265 270  
 Trp His Pro Asn Leu Arg Ala Phe Gly Ile Val Glu Cys Val Leu Cys  
 275 280 285  
 Thr Cys Asn Val Thr Lys Gln Glu Cys Lys Lys Ile His Cys Pro Asn  
 290 295 300  
 Arg Tyr Pro Cys Lys Tyr Pro Gln Lys Ile Asp Gly Lys Cys Cys Lys  
 305 310 315 320  
 Val Cys Pro Gly Lys Lys Ala Lys Glu Glu Leu Pro Gly Gln Ser Phe  
 325 330 335  
 Asp Asn Lys Gly Tyr Phe Cys Gly Glu Glu Thr Met Pro Val Tyr Glu  
 340 345 350  
 Ser Val Phe Met Glu Asp Gly Glu Thr Thr Arg Lys Ile Ala Leu Glu  
 355 360 365  
 Thr Glu Arg Pro Pro Gln Val Glu Val His Val Trp Thr Ile Arg Lys  
 370 375 380

Gly Ile Leu Gln His Phe His Ile Glu Lys Val Ser Lys Arg Met Phe  
 385 390 395 400

Glu Glu Leu Pro His Phe Lys Leu Val Thr Arg Thr Thr Leu Ser Gln  
 405 410 415

Trp Lys Ile Phe Thr Glu Gly Glu Ala Gln Ile Ser Gln Met Cys Ser  
 420 425 430

Ser Arg Val Cys Arg Thr Glu Leu Glu Asp Leu Val Lys Val Leu Tyr  
 435 440 445

Leu Glu Arg Ser Glu Lys Gly His Cys  
 450 455

&lt;210&gt; 77

&lt;211&gt; 2050

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 77

```

gtgtctctgag aagccggact acgcccgcgc ggtctcttcaa agccgagccg ggagttttttg 60
ctacagttttt cgcacccatg agtcgcagct ataatgatga gctgcagttc ttggagaaga 120
tcactaaaaa ctgctggagg atcaagaagg gcttctgtgc caacatgcag gttgaaggta 180
ttttctatgt gaattatgt ctggagaaat tgatgtttga ggaattaaag aatgcctgtc 240
gaggttgttg tgttggtggc ttcctgcag ccatgaacca gattggcaat gtggcagccc 300
tgcttgaat tcttcataga tctattgggc tctctgatgt ccattcagga tatgggttcg 360
ctatttggga catggcagcc ttctatata atgacctga agcagtagta tccccaggtg 420
gtgtgggggt tgacatcaac tgtggtgttc gcttgctaag aacccattta gatgaangtg 480
atgtccagcc tgtgaaggag caacttgcgc agctatgtt tgaccacatt cctgttgggg 540
tggttcaaaa aggtgtcctt ccaatgaatg ccaagagctt ggaaggagcc ttggagatcg 600
gggttggact gttcttaca ggaaggtatg cctgggttga agacaaggag cactgcagag 660
agtaagggaag gatgtctcag gctgacccca ataaagtttc tgcaaggggc aagaaaaag 720
gccttctctc gttggggacc ctgggagcag gcaaccatta tgcagaaatc caggttgttg 780
atgagatttc caatgagtat gctgctaaaa aaatgggcct cgaaccataa ggacaggtgt 840
gtgtgatgat ccacagtggg agcagaggtt tggggccacca agtagccaca gatgcgttg 900
tagtatatga gaagcccatg aagagagaca agattatagt caatgatcgg cagttggctt 960
gtgtctgaat cgttcccca gagggtcag actatctgaa ggaatggca gctgctggga 1020
actatgcctg ggtcaaccgc tcttccatga ccttctaac cgtcaggtt ttggccaagg 1080
tcttcaacac aacctctgat gacttgcacc tccatgtgat ctatgatgtt tctcaacaca 1140
ttgccaaggt ggagcagcat gtggtggacg gaaaggaaag gacactgtta gtacacagga 1200
agggatccac cgcgctttc cctctctacc atccctcat tgcgttgtgt caccacacta 1260
ctggacagcc agtctctatt ggtggcacca tgggaacctg tagttatgtt ctacttggca 1320
ctgaacaggg catgactgag acctttggaa caacctgcca tggagcgggc cgtgcattgt 1380
ccggggcaaa atctcagagt aatttagatt gcttcaacca aactgttat ggaagaggct cctgagtcct 1500
tggaatttgc gatcgtgtt gcttcaacca cctatgatgc tggatcagc aagaaggcca 1560
ataaagatgt gacagatgtg gtanatacct gccatgatgc tggacagcag ggtgcctga 1620
ttaaactgag accaattgct gtgatcaag gatagaaact tggacagcag ggtgcctga 1680
caccaccaac cctctctgaa gtggaagtgg actgacatgc tcttctgaca tcagactcaa 1740
ggcgggacaa gttgcaaat gtgcagctgt aactgtctac gccaaaatgg ctgatgggga 1800
ggctgctgtt ttcgggggac cgtgcttcta aaataacctt ccagggaagag gcacattgac 1860
caactttgga aagggaaggaa tatgccttct ccttgggtgt cccacagagt tttaggaaaa 1920
tctgtttagg atgggttagt gtcacactgc cttacagcag catactgate tttagccatc 1980
agattgatct tcttcaacc aagctctgtt tccattccga gagggtgcat gaagaaagtt 2040
ctgttcacaa agggaaaaaa aaaaaaana aaaaaaana aaaaaaana aaaaaaana 2050

```

&lt;210&gt; 78

&lt;211&gt; 505

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 78

```

Met Ser Arg Ser Tyr Asn Asp Glu Leu Gln Phe Leu Glu Lys Ile Asn
 1           5           10           15

Lys Asn Cys Trp Arg Ile Lys Lys Gly Phe Val Pro Asn Met Gln Val
          20           25           30

Glu Gly Val Phe Tyr Val Asn Asp Ala Leu Glu Lys Leu Met Phe Glu
          35           40           45

Glu Leu Arg Asn Ala Cys Arg Gly Gly Gly Val Gly Gly Phe Leu Pro
 50           55           60

Ala Met Lys Gln Ile Gly Asn Val Ala Ala Leu Pro Gly Ile Val His
 65           70           75           80

Arg Ser Ile Gly Leu Pro Asp Val His Ser Gly Tyr Gly Phe Ala Ile
          85           90           95

Gly Asn Met Ala Ala Phe Asp Met Asn Asp Pro Glu Ala Val Val Ser
          100          105          110

Pro Gly Gly Val Gly Phe Asp Ile Asn Cys Gly Val Arg Leu Leu Arg
          115          120          125

Thr Asn Leu Asp Glu Ser Asp Val Gln Pro Val Lys Glu Gln Leu Ala
          130          135          140

Gln Ala Met Phe Asp His Ile Pro Val Gly Val Gly Ser Lys Gly Val
          145          150          155          160

Ile Pro Met Asn Ala Lys Asp Leu Glu Glu Ala Leu Glu Met Gly Val
          165          170          175

Asp Trp Ser Leu Arg Glu Gly Tyr Ala Trp Ala Glu Asp Lys Glu His
          180          185          190

Cys Glu Glu Tyr Gly Arg Met Leu Gln Ala Asp Pro Asn Lys Val Ser
          195          200          205

Ala Arg Ala Lys Lys Arg Gly Leu Pro Gln Leu Gly Thr Leu Gly Ala
          210          215          220

Gly Asn His Tyr Ala Glu Ile Gln Val Val Asp Glu Ile Phe Asn Glu
          225          230          235          240

Tyr Ala Ala Lys Lys Met Gly Ile Asp His Lys Gly Gln Val Cys Val
          245          250          255

Met Ile His Ser Gly Ser Arg Gly Leu Gly His Gln Val Ala Thr Asp
          260          265          270

Ala Leu Val Ala Met Glu Lys Ala Met Lys Arg Asp Lys Ile Ile Val
          275          280          285

Asn Asp Arg Gln Leu Ala Cys Ala Arg Ile Ala Ser Pro Glu Gly Gln
          290          295          300

```

Asp Tyr Leu Lys Gly Met Ala Ala Ala Gly Asn Tyr Ala Trp Val Asn  
 305 310 315 320  
 Arg Ser Ser Met Thr Phe Leu Thr Arg Gln Ala Phe Ala Lys Val Phe  
 325 330 335  
 Asn Thr Thr Pro Asp Asp Leu Asp Leu His Val Ile Tyr Asp Val Ser  
 340 345 350  
 His Asn Ile Ala Lys Val Glu Gln His Val Val Asp Gly Lys Glu Arg  
 355 360 365  
 Thr Leu Leu Val His Arg Lys Gly Ser Thr Arg Ala Phe Pro Pro His  
 370 375 380  
 His Pro Leu Ile Ala Val Asp Tyr Gln Leu Thr Gly Gln Pro Val Leu  
 385 390 395 400  
 Ile Gly Gly Thr Met Gly Thr Cys Ser Tyr Val Leu Thr Gly Thr Glu  
 405 410 415  
 Gln Gly Met Thr Glu Thr Phe Gly Thr Thr Cys His Gly Ala Gly Arg  
 420 425 430  
 Ala Leu Ser Arg Ala Lys Ser Arg Arg Asn Leu Asp Phe Gln Asp Val  
 435 440 445  
 Leu Asp Lys Leu Ala Asp Met Gly Ile Ala Ile Arg Val Ala Ser Pro  
 450 455 460  
 Lys Leu Val Met Glu Glu Ala Pro Glu Ser Tyr Lys Asn Val Thr Asp  
 465 470 475 480  
 Val Val Asn Thr Cys His Asp Ala Gly Ile Ser Lys Lys Ala Ile Lys  
 485 490 495  
 Leu Arg Pro Ile Ala Val Ile Lys Gly  
 500 505

&lt;210&gt; 79

&lt;211&gt; 1178

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 79

gccaaatgtc cggtaaatgat gtaacacagc tccagtggcc cagccagttc gagtcagggt 60  
 tctccagggg aagaaacaga tcaactgaa accgtgtcag ttcagttctc ggtattgggg 120  
 aagggtgtaa aacatcgacc cccaccaatc aaacttccct caagctcagg aaatagttcc 180  
 tcaggtaact attttacacc acaacagaca agcagcttct tcaaatctcc aactctctcc 240  
 ccttctctca agccatcgag tatctctcgg aaactctctg tggatctcaa tcaagtttagc 300  
 atgtttcttc cagctgcctt atcaactgac agtctctaac aaagaaccac ggcaccccag 360  
 gtcattgcaa actctgctgg acttaacttc atcaatgtag tgggtctctg ttgtggggcc 420  
 caggttttga tgaagtggtc aaaccccatt ctgggtgtga acactgggtc cataactctc 480  
 gcaggataaa aactgagcgg cttctacccc tcaagggttc tgcatacaaa tgcactgccc 540  
 agtgcaatgc aggcagcttc tcaagcaggt gttccatttc gtttaaaaaa tacttcaagt 600  
 ctacaggccct caactctact caagcttcca ggtggtttac ttatttttaa cactctgccc 660  
 cagcagcaac agcagctctc ccagtttacc caacaacnac ctccagcagc cacaacttgt 720  
 agtctctaac agccagggga gcagggttct gagcaaggtt caaccagtca agacagggcc 780

```

ttatctgtctc agcagctgc tgttattaac cttactegag taggaagtct tatgcagtcn 840
caggcagctc cagttgcgat tottgtagta tcaaatggct atggcagcag cagcagcaca 900
aacagctcag ctacatcatt atgggcattc aggcagccag tcaaaaagta aaatgagag 960
aggcatgcca accactcaca aattttgagt ctgcaattac tttttgttc ttttttaaaa 1020
atcacagagc actgaatcaa aagaattgag ttctacttt ttgtttttt caatgltgca 1080
gtattttaca ttgtatgatg tacaacttt atcagaagc acaccttat cattttttaa 1140
tcaaacaggg gaatatgttt acaaaaaaa aaaaaaaa 1178

```

&lt;210&gt; 80

&lt;211&gt; 310

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

```

Met Ser His Ser Ser Ser Gly Ser Ala Ser Leu Ser Gln Val Ser Pro
  1                      5                      10                      15

Gly Lys Glu Thr Asp Gln Thr Glu Thr Val Ser Val Gln Ser Ser Val
    20                      25                      30

Leu Gly Lys Gly Val Lys His Arg Pro Pro Pro Ile Lys Leu Pro Ser
    35                      40                      45

Ser Ser Gly Asn Ser Ser Ser Gly Asn Tyr Phe Thr Pro Gln Gln Thr
    50                      55                      60

Ser Ser Phe Leu Lys Ser Pro Thr Pro Pro Pro Ser Ser Lys Pro Ser
    65                      70                      75                      80

Ser Ile Pro Arg Lys Ser Ser Val Asp Leu Asn Gln Val Ser Met Leu
    85                      90                      95

Ser Pro Ala Ala Leu Ser Pro Ala Ser Ser Ser Gln Arg Thr Thr Ala
   100                      105                      110

Thr Gln Val Met Ala Asn Ser Ala Gly Leu Asn Phe Ile Asn Val Val
   115                      120                      125

Gly Ser Val Cys Gly Ala Gln Ala Leu Met Ser Gly Ser Asn Pro Met
   130                      135                      140

Leu Gly Cys Asn Thr Gly Ala Ile Thr Pro Ala Gly Ile Asn Leu Ser
   145                      150                      155                      160

Gly Leu Leu Pro Ser Gly Gly Leu Leu Pro Asn Ala Leu Pro Ser Ala
   165                      170                      175

Met Gln Ala Ala Ser Gln Ala Gly Val Pro Phe Gly Leu Lys Asn Thr
   180                      185                      190

Ser Ser Leu Arg Pro Leu Asn Leu Leu Gln Leu Pro Gly Gly Ser Leu
   195                      200                      205

Ile Phe Asn Thr Leu Gln Gln Gln Gln Gln Leu Ser Gln Phe Thr
   210                      215                      220

Pro Gln Gln Pro Gln Gln Pro Thr Thr Cys Ser Pro Gln Gln Pro Gly
   225                      230                      235                      240

Glu Gln Gly Ser Glu Gln Gly Ser Thr Ser Gln Glu Gln Ala Leu Ser

```



	245		250		255
Ala Gln Gln Ala Ala Val Ile Asn Leu Thr Gly Val Gly Ser Phe Met					
260		265		270	
Gln Ser Gln Ala Ala Ala Val Ala Ile Leu Ala Ala Ser Asn Gly Tyr					
275	280		285		
Gly Ser Ser Ser Ser Thr Asn Ser Ser Ala Thr Ser Ser Ser Ala Tyr					
290	295		300		
Arg Gln Pro Val Lys Lys					
305	310				

<210> B1  
 <211> 641  
 <212> DNA  
 <213> Homo sapiens

<400> B1  
 gcgagtgta cccgtgctc cctccctgct tcttgctctg ctaactcaac ctgctcttcc 60  
 tcttttttcat tcttctactc tgcctctatc ggaggacaaa tggacactag ggggtgctaac 120  
 cttattgggtg cctgccccag cctaccccag gtgcccagca actctcgtgc acaggaggt 180  
 cccacagtta tggagccgg aaagaatttc tctgcactgg atggactgta tattgagatt 240  
 aaaaattata ttccttatat tcttgcttat atcaatgctc tctctgtaaa acctcttcc 300  
 agcctcaatt ctctcaactg atcttggtta ggcgttgat tctcttatt tactctttgc 360  
 ttgactggtt cctcctaac ctctccccc tagcactcta ctctcttaag ctggttggtc 420  
 attaacctctg ttggatccac tctctgggaa agacttctgt taatgtaagt gaccttactc 480  
 cctggatggt gtcactagtc tagtggtctt cgttaaataa acctttctta tttctaaaa 540  
 aaaaaaa aaaaaaa aaaaaaa aaaaaaa aaaaaaa aaaaaaa 600  
 aaaaaaa aaaaaaa aaaaaaa aaaaaaa a 641

<210> B2  
 <211> 94  
 <212> PRT  
 <213> Homo sapiens

<400> B2  
 Met Ser Pro Cys Pro Pro Ser Leu Leu Leu Ala Leu Leu Thr Gln Leu  
 1 5 10 15  
 Cys Leu Pro Leu Phe His Ser Ser Thr Leu Pro Tyr Met Glu Asp Lys  
 20 25 30  
 Trp Thr Pro Gly Val Leu Thr Leu Leu Val Pro Ala Pro Ala Tyr Pro  
 35 40 45  
 Arg Cys Gln Gln Thr Leu Val His Arg Arg Leu Pro Gln Leu Trp Ser  
 50 55 60  
 Gln Glu Arg Ile Ser Leu His Trp Met Asp Cys His Leu Arg Leu Lys  
 65 70 75 80  
 Ile Ile Phe Leu Ile Phe Leu Leu Ile Ser Met Leu Ser Leu  
 85 90

<210> B3  
 <211> B32

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 83

```

ccttgcatccta ccttctctctg ccttatctctg tcttaggtgt aaacctgttt tttctcacc 60
tgaacttggtg aaccaatcct ggcattataa caaaagcaaa tgaattatta tttcttcctg 120
tttatgaattc tgaatgaagt atgtttccaa agaactgtag gtcctctact tgtgatttaa 180
ggaaaccagc tcatcccaag caatgcagtg tctgtaactg gtgtgtgcac cgtttcgacc 240
atcaactgtgt ttgggtgca cactgcacat gggcctggaa catcaggtac ttcctcatct 300
acgtcttgac cttgacggcc tgggtgcca ccttcctcat tctgagcacc acttttcttg 360
tccactctgt ggtgatgtca gatttatacc aggaactta catcgatgac cttggacacc 420
ttccatgtta tggacacggt cttctctatt cactacctgt tctgacttt tccacggatt 480
gtcttcacatc tgggtcttgt cgtgcttctg agcttctctc tgggtggcta cctgttcttt 540
gtctctgata tggcgggcac caaccagact actaacgagt ggtacagagg tgaactggcc 600
tgggtccacc gttgtccct tctgctctgg cctcctgca cagagcccca agtccacgg 660
aacatccact cccatgggtc tggagacaa cttcaagaga tcttctacc tgcctttcca 720
tgtcatgaga ggaagaacaa agaatgaca gtgtatgact gcctttgagc tatagttccc 780
gtttattttac acatgtggat cctggttttc ctcccgattg aattctagac ct 832

```

&lt;210&gt; 84

&lt;211&gt; 140

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 84

```

Met Phe Pro Lys Asn Val Arg Cys Ser Thr Cys Asp Leu Arg Lys Pro
  1             5             10             15

Ala Arg Ser Lys His Cys Ser Val Cys Asn Trp Cys Val His Arg Phe
          20             25             30

Asp His His Cys Val Trp Val Asn Asn Cys Ile Gly Ala Trp Asn Ile
          35             40             45

Arg Tyr Phe Leu Ile Tyr Val Leu Thr Leu Thr Ala Ser Ala Ala Thr
          50             55             60

Val Ala Ile Val Ser Thr Thr Phe Leu Val His Leu Val Val Met Ser
          65             70             75             80

Asp Leu Tyr Gln Glu Thr Tyr Ile Asp Asp Leu Gly His Leu Pro Cys
          85             90             95

Tyr Gly His Gly Leu Ser Tyr Ser Val Pro Val Pro Asp Phe Ser Thr
          100            105            110

Asp Cys Leu His Ala Gly Leu Cys Arg Gly Ser Glu Leu Pro Pro Gly
          115            120            125

Trp Leu Pro Val Val Cys Pro Val Ser Gly Gly His Gln Pro Asp Tyr
          130            135            140

```

&lt;210&gt; 85

&lt;211&gt; 3790

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 85

```

aacggcagtc tcaatctggc cccacacatt tcttgggctt gtaggaaggt ggacatgggc 60

```

tcacggagac	aagacaagtg	atatgttgaa	ctgttcgggt	gttggaatca	actgctcctg	120
gagtgcctta	agggcagtg	ttatcagaa	ttapccagg	ccagccagc	agggcagat	180
gctctgcat	gaatggccac	gtaggcagag	actgacaagc	ggtaggcaat	gagcttccc	240
ctcggactgc	tccttccctg	tgtgttcagg	ggagggggtc	actttctggc	aactctgctg	300
ctgctgctgc	tgtgtgctgc	acttcagctt	ctctctcact	caagggtcagc	aggtcagggg	360
agggcaggt	gctagggaaa	gctttgtacc	atgaaccagg	tcggaaagtt	ttccggagg	420
agtggggag	tcttggcatt	tatcttctga	gcttctgtca	tctggctcct	ctttgacatg	480
gcagctctcc	gctctctcatt	cagtgcagtc	accctctggg	tcctcaggga	agcatttgtg	540
aggaggagg	ggataggatt	cagcgttcag	ccagaccag	gaaaaatttt	ttcagcagc	600
atgaaggag	tgaacctcc	ctaaaggga	catgggaag	gggcattggg	caaggagat	660
gttagaaaa	ctgaggagag	tgtactcagg	gttaggttg	acttggacca	aacccaggag	720
gaagaaaaa	tgcagaaagc	cctgggaagg	ggcagggtg	tgcgcttgtg	gcctcctgca	780
catctggaga	ccctccctgt	gaactcctaa	aagcagagag	cagacgggag	aggcacaaaa	840
cctgagggct	cctctcacc	ggggacacca	aagcaaacga	cagctcaggg	ggtcccaag	900
acctcattca	tagcagcaca	aggaactcag	gtagtcaaaa	tatcagtaac	catgggacgt	960
gtcagtttaa	aacaggagcc	ccggaagag	catagtccca	gcagtgcac	atcaaaacta	1020
gcagctgaaa	gggacttgaa	tgtgacctac	agtcttagta	ctgatagacc	aaagcagcga	1080
tcacagggag	tagcaaacga	gagggcacac	cctgaccaga	cagcagtgcc	gaagtctggg	1140
gaagccatgg	ccttaaacca	aactaagact	cagagcaaa	aaagcaatgc	aaataaacac	1200
aaagtcata	cggctcttcc	tttctcctaa	ttcactgtca	attcacaatc	cttaagggaag	1260
caatctacta	atgagacacc	ttgggaaggt	ttgtcaaaag	atgatggagc	tagaggggct	1320
cctggagaga	aactcatttt	ctctggaagc	catcttgtga	ttataaccac	agagggaagag	1380
caaaaggcag	accccaaaag	ggtctctaat	totaaaacta	aaacaaatct	tcctaaagta	1440
ttgggttaaa	gcaaaagtaa	acacatttcc	aggaatagaa	gtgagatgtc	ttctctctca	1500
cctgtccac	atagagtgc	actgtcccaa	actaacacag	ctttaaactg	agggctagag	1560
ccagcaaaa	tcacataac	tgcacaagcc	cctctacacg	aatacaacca	gagtcataa	1620
aaagcccttt	tacctgaaga	cagtggaaag	caccaggtgt	taagcaattg	tgtgacactt	1680
tctccaaagg	accccaaaag	tccagggcag	tttggggctc	ctgtagtgtt	cccccatgga	1740
aaggagagag	aggcagaaag	aagatggaaa	gaaggaaact	ccaatgtcta	ccttagcgat	1800
ttgatccca	tggatagagc	cattgaagac	accagacctg	ctggatgtgc	agagcageta	1860
gttcacaata	acctcccaac	caccagtgcc	atcatgttgt	ctgtgatgta	agtgtgggtc	1920
actctcctga	gactgttcca	cagtgtcacc	aactgtcttc	ctccacacct	catcaaggag	1980
attctgtctg	tagatgactt	cagccacca	gactatctaa	aagataaatt	ggataaatac	2040
atgtccaggt	ttccaaaggt	tgggattctt	cgctccaaag	agagacatgg	cttaataaag	2100
gcccaggctg	caggagcaca	gaatgcacaa	ggtgatgtgt	tgacattttt	agattctcat	2160
gtggaaatgt	acgttgggtt	gttggaaact	cttctggaaa	gagtttattt	aagtagaaag	2220
aaagtggcct	gtccagtaac	cgaaagtcac	aatgataagg	atatgagtta	catgacagtg	2280
gataactctc	aaaggggcat	ctttgtgttg	cccatgacct	ttggttggag	aaacattcct	2340
ccagatgtcc	tggcaaaa	cagaattaaa	gaactgtata	caataagggt	ccttgtctat	2400
gctgggtggat	tttttcttat	tgaacaaagt	tacttlttly	aacttgggac	atacgacct	2460
ggccttgatg	tttgggggtg	gaaaaatakt	gagctctcat	tcaagggtgt	gatgtgtggt	2520
ggtgaaattg	agatcatttc	ctgtctccga	gtgggocata	tattcagaaa	tgacaatcca	2580
tattctctcc	ccaaagaccg	gatgaagacc	gtggggggga	acttgggtgc	ggttgcagg	2640
gtctgggtgg	atgagtata	ggagctgttc	tatggccatg	gagacacacct	catcgaccaa	2700
gggttagatg	ttggcaacct	cacccagcaa	agggagctgc	gaaggaacct	gaagtgcana	2760
agtttccaat	ggtaccttga	gaatgtcttt	cctgacttaa	gggtcccat	tgtgagagct	2820
agtgtgtgtc	ttactaatgt	ggttttgggt	aatgacattt	ccattgaana	cactacagtc	2880
attctggaa	actgcgttgg	gagcaaaag	cttcaacaa	ttaatcacac	ctggttaaga	2940
cttatttaaa	gtggaggaatg	gtgtatagcc	cccatccctg	ataaaggagc	cgtaaggctg	3000
caccccttgt	ataacagana	caaagggtca	aatggtgtgc	ataaatcaac	ataagctctt	3060
catccagaac	tggtyaatca	cattgttttt	gaacaacatc	agcaattatt	atgcttggga	3120
ggaaattttt	ctcaaaagat	cctgaabgta	gctgcctgtg	accagtgaa	gcatatcaa	3180
agtggaatat	ttgaanaata	ttatgaagcc	tgaagtgtaa	ctgatgtttt	tatatagtaa	3240
acccattaaa	tactgtgaaa	ataacactga	acttggaaac	takatctctc	agcggtagtt	3300
taaatcttca	atttctatca	catttgaatg	gaagattttt	tataaatcac	aatatttggg	3360
ataccaaaag	atgactcagg	aaaacagtc	aacttggac	tgaagtccct	cttgggaact	3420
gggtggcctt	tgaattgctt	gctttccacc	ctatgtctaga	cctcatcatg	caaatctccc	3480
tgtgaaagct	aacaggtaac	tggaaatgaa	gacagaagga	cttgagaaag	catgaggaata	3540
ttcccaatga	ctatgttttg	taataatcag	ctctctctgg	ccacacgtag	gaatgatcaa	3600
tgagaaactta	acttagtctt	ttatttgggg	atttttctac	caaacaaaaa	tttcttgggt	3660

ctcttatgggt agagagacctc agatgcccac agctgtccacc ttctgtgmaat cccctccagac 3720  
 tacatgcctg cctacctaac agtttgaat agtattgata tactgctggt aaaaaaaaaa 3780  
 aaaaaaaaaa 3790

<210> 86  
 <211> 940  
 <212> PRT  
 <213> Homo sapiens

<400> 86  
 Met Asn Arg Ile Arg Lys Phe Phe Arg Gly Ser Gly Arg Val Leu Ala  
 1 5 10 15  
 Phe Ile Phe Val Ala Ser Val Ile Trp Leu Leu Phe Asp Met Ala Ala  
 20 25 30  
 Leu Arg Leu Ser Phe Ser Glu Ile Asn Thr Arg Val Ile Lys Glu Asp  
 35 40 45  
 Ile Val Arg Arg Glu Arg Ile Gly Phe Arg Val Gln Pro Asp Gln Gly  
 50 55 60  
 Lys Ile Phe Tyr Ser Ser Ile Lys Glu Met Lys Pro Pro Leu Arg Gly  
 65 70 75 80  
 His Gly Lys Gly Ala Trp Gly Lys Glu Asn Val Arg Lys Thr Glu Glu  
 85 90 95  
 Ser Val Leu Lys Val Glu Val Asp Leu Asp Gln Thr Gln Arg Glu Arg  
 100 105 110  
 Lys Met Gln Asn Ala Leu Gly Arg Gly Lys Val Val Pro Leu Trp His  
 115 120 125  
 Pro Ala His Leu Gln Thr Leu Pro Val Thr Pro Asn Lys Gln Lys Thr  
 130 135 140  
 Asp Gly Arg Gly Thr Lys Pro Glu Ala Ser Ser His Gln Gly Thr Pro  
 145 150 155 160  
 Lys Gln Thr Thr Ala Gln Gly Ala Pro Lys Thr Ser Phe Ile Ala Ala  
 165 170 175  
 Lys Gly Thr Gln Val Val Lys Ile Ser Val His Met Gly Arg Val Ser  
 180 185 190  
 Leu Lys Gln Glu Pro Arg Lys Ser His Ser Pro Ser Ser Asp Thr Ser  
 195 200 205  
 Lys Leu Ala Ala Glu Arg Asp Leu Asn Val Thr Ile Ser Leu Ser Thr  
 210 215 220  
 Asp Arg Pro Lys Gln Arg Ser Gln Ala Val Ala Asn Glu Arg Ala His  
 225 230 235 240  
 Pro Ala Ser Thr Ala Val Pro Lys Ser Gly Glu Ala Met Ala Leu Asn  
 245 250 255  
 Lys Thr Lys Thr Gln Ser Lys Glu Val Asn Ala Asn Lys His Lys Ala  
 260 265 270

Asn Thr Ser Leu Pro Phe Pro Lys Phe Thr Val Asn Ser Asn Arg Leu  
 275 285  
 Arg Lys Gln Ser Ile Asn Glu Thr Pro Leu Gly Ser Leu Ser Lys Asp  
 290 295 300  
 Asp Gly Ala Arg Gly Ala His Gly Lys Lys Leu Asn Phe Ser Glu Ser  
 305 310 315 320  
 His Leu Val Ile Ile Thr Lys Glu Glu Glu Gln Lys Ala Asp Pro Lys  
 325 330 335  
 Glu Val Ser Asn Ser Lys Thr Lys Thr Ile Phe Pro Lys Val Leu Gly  
 340 345 350  
 Lys Ser Gln Ser Lys His Ile Ser Arg Asn Arg Ser Glu Met Ser Ser  
 355 360 365  
 Ser Ser Leu Ala Pro His Arg Val Pro Leu Ser Gln Thr Asn His Ala  
 370 375 380  
 Leu Thr Gly Gly Leu Glu Pro Ala Lys Ile Asn Ile Thr Ala Lys Ala  
 385 390 395 400  
 Pro Ser Thr Glu Tyr Asn Gln Ser His Ile Lys Ala Leu Leu Pro Glu  
 405 410 415  
 Asp Ser Gly Thr His Gln Val Leu Arg Ile Asp Val Thr Leu Ser Pro  
 420 425 430  
 Arg Asp Pro Lys Ala Pro Gly Gln Phe Gly Arg Pro Val Val Val Pro  
 435 440 445  
 His Gly Lys Glu Lys Glu Ala Glu Arg Arg Trp Lys Glu Gly Asn Phe  
 450 455 460  
 Asn Val Tyr Leu Ser Asp Leu Ile Pro Val Asp Arg Ala Ile Glu Asp  
 465 470 475 480  
 Thr Arg Pro Ala Gly Cys Ala Glu Gln Leu Val His Asn Asn Leu Pro  
 485 490 495  
 Thr Thr Ser Val Ile Met Cys Phe Val Asp Glu Val Trp Ser Thr Leu  
 500 505 510  
 Leu Arg Ser Val His Ser Val Ile Asn Arg Ser Pro Pro His Leu Ile  
 515 520 525  
 Lys Glu Ile Leu Leu Val Asp Asp Phe Ser Thr Lys Asp Tyr Leu Lys  
 530 535 540  
 Asp Asn Leu Asp Lys Tyr Met Ser Gln Phe Pro Lys Val Arg Ile Leu  
 545 550 555 560  
 Arg Leu Lys Glu Arg His Gly Leu Ile Arg Ala Arg Leu Ala Gly Ala  
 565 570 575  
 Gln Asn Ala Thr Gly Asp Val Leu Thr Phe Leu Asp Ser His Val Glu  
 580 585 590

Cys Asn Val Gly Trp Leu Glu Pro Leu Leu Glu Arg Val Tyr Leu Ser  
 595 600 605  
 Arg Lys Lys Val Ala Cys Pro Val Ile Glu Val Ile Asn Asp Lys Asp  
 610 615 620  
 Met Ser Tyr Met Thr Val Asp Asn Phe Gln Arg Gly Ile Phe Val Trp  
 625 630 635 640  
 Pro Met Asn Phe Gly Trp Arg Thr Ile Pro Pro Asp Val Ile Ala Lys  
 645 650 655  
 Asn Arg Ile Lys Glu Thr Asp Thr Ile Arg Cys Pro Val Met Ala Gly  
 660 665 670  
 Gly Leu Phe Ser Ile Asp Lys Ser Tyr Phe Phe Glu Leu Gly Thr Tyr  
 675 680 685  
 Asp Pro Gly Leu Asp Val Trp Gly Gly Glu Asn Met Glu Leu Ser Phe  
 690 695 700  
 Lys Val Trp Met Cys Gly Gly Glu Ile Glu Ile Ile Pro Cys Ser Arg  
 705 710 715 720  
 Val Gly His Ile Phe Arg Asn Asp Asn Pro Tyr Ser Phe Pro Lys Asp  
 725 730 735  
 Arg Met Lys Thr Val Glu Arg Asn Leu Val Arg Val Ala Glu Val Trp  
 740 745 750  
 Leu Asp Glu Tyr Lys Glu Leu Phe Tyr Gly His Gly Asp His Leu Ile  
 755 760 765  
 Asp Gln Gly Leu Asp Val Gly Asn Leu Thr Gln Gln Arg Glu Leu Arg  
 770 775 780  
 Lys Lys Leu Lys Cys Lys Ser Phe Lys Trp Tyr Leu Glu Asn Val Phe  
 785 790 795 800  
 Pro Asp Leu Arg Ala Pro Ile Val Arg Ala Ser Gly Val Leu Ile Asn  
 805 810 815  
 Val Ala Leu Gly Lys Cys Ile Ser Ile Glu Asn Thr Thr Val Ile Leu  
 820 825 830  
 Glu Asp Cys Asp Gly Ser Lys Gln Leu Gln Gln Phe Asn Tyr Thr Trp  
 835 840 845  
 Leu Arg Leu Ile Lys Cys Gly Glu Trp Cys Ile Ala Pro Ile Pro Asp  
 850 855 860  
 Lys Gly Ala Val Arg Leu His Pro Cys Asp Asn Arg Asn Lys Gly Leu  
 865 870 875 880  
 Lys Trp Leu His Lys Ser Thr Ser Val Phe His Pro Glu Leu Val Asn  
 885 890 895  
 His Ile Val Phe Glu Asn Asn Gln Gln Leu Leu Cys Leu Glu Gly Asp  
 900 905 910

Phe Ser Gln Lys Ile Leu Lys Val Ala Ala Cys Asp Pro Val Lys Pro  
 915 920 925

Tyr Gln Lys Trp Lys Phe Glu Lys Tyr Tyr Glu Ala  
 930 935 940

<210> 87  
 <211> 1200  
 <212> DNA  
 <213> Homo sapiens

<400> 87  
 ggctttctcgg agcggcgctg ggcgcaccgga gcagggtcga gatgtccctac atcccggggcc 60  
 agccgggtcac cgcggtggcg caaagagttg aaatttcaca gctgcgtcaa ggtgagaaact 120  
 taatccctggg ttccagcatt ggaaggtgga tgcaccagga tccctcccg aatccctct 180  
 atgaagacaa gacggacaa ggtattttatgc caaccgggt gctcgaagga ggcctctgtg 240  
 aaatcgctgg gctgcagatt ggaacacaga tcatgcaggt gaaagggtgg gacatgacca 300  
 tggtcacaca cgaaccaggcc cgaagcggc tcaaccaggc ctccggaggag gtgtgtcgtc 360  
 tctgtgtgac ggggcagtcg ctgcagaagg cgtgcagcag tcatgtctgtc ctgcagacca 420  
 ccaccatctg cgaactctgc ctgcgcctc tctgtacagt aatgcacact ccacactctg 480  
 tcccatctg gcttctgtgt accagctttc tctctggac aacgaggact ggaataaagg 540  
 acctggagct gctcagtcgc cagtcctgtc tgaccacagg ctccaggtccg acctgtgtgc 600  
 ttggccacag cagtggctgg gcaagtcgga accactctct ctggggagcc cccaaagct 660  
 ggaatatgct ggaaggaaca ggcctttccc gcttttgcct ggtgcaggc ttggcctccg 720  
 cccctgcccc ccaagctctg tctgtccaca cgcagtgct tctgcccctc gggggactgg 780  
 acacacatcc tgcacagagc gctacgaagg ttgtccaga tgaagccagg tgggctccgc 840  
 gttcactccc actctccaga ggggtgtcgg cctcccagc gtttgccttc ttaaggattt 900  
 agacagaggt cgaaggtcac ctatcagggc agctctcagg attgkcttt tcccttttgc 960  
 ctgtgggttt aacttttcta ttttttaat ccaagtttg atacaaaatg tttttatcgt 1020  
 actctttgga gatgccatt ctacttttga atttgcctt tactaattcg catctggaag 1080  
 ctccgcaagt gcaacagct tacttttgtt accgtggaaa ccactgcgc cctcccgga 1140  
 tctgtgtgtc tgaataaaaa tgcctggcatt caaaaaaaa aaaaaaaa aaaaaaaa 1200

<210> 88  
 <212> 286  
 <212> PRT  
 <213> Homo sapiens

<400> 88  
 Met Ser Tyr Ile Pro Gly Gln Pro Val Thr Ala Val Val Gln Arg Val  
 1 5 10 15  
 Glu Ile His Lys Leu Arg Gln Gly Glu Asn Leu Ile Leu Gly Phe Ser  
 20 25 30  
 Ile Gly Gly Gly Ile Asp Gln Asp Pro Ser Gln Asn Pro Phe Tyr Glu  
 35 40 45  
 Asp Lys Thr Asp Lys Gly Ile Tyr Val Thr Arg Val Ser Glu Gly Gly  
 50 55 60  
 Pro Ala Glu Ile Ala Gly Leu Gln Ile Gly Asp Lys Ile Met Gln Val  
 65 70 75 80  
 Asn Gly Trp Asp Met Thr Met Val Thr His Asp Gln Ala Arg Lys Arg  
 85 90 95  
 Leu Thr Lys Arg Ser Glu Glu Val Val Arg Leu Leu Val Thr Arg Gln

100	105	110
Ser Leu Gln Lys Ala Cys Ser Ser His Ala Val Leu Ala Ala Thr Thr		
115	120	125
Ile Cys Asp Ser Cys Leu Pro Pro Leu Cys Thr Val Thr Pro Leu Pro		
130	135	140
His Ser Val Pro Ile Trp Leu Leu Leu Thr Ser Phe Leu Ser Trp Thr		
145	150	155
Pro Arg Ile Gly Asn Lys Gly Leu Glu Leu Ser Ser Ser Gln Ser Ala		
165	170	175
Val Thr Thr Gly Ser Gly Pro Thr Leu Leu Leu Gly His Ser Ser Gly		
180	185	190
Trp Ala Ser Gly Asn His Tyr Leu Leu Gly Ala Pro Lys Ser Trp Glu		
195	200	205
Met Leu Glu Glu Pro Gly Leu Ser Arg Phe Cys Leu Ala Ala Gly Leu		
210	215	220
Gly Ser Ala Pro Ala Pro Gln Pro Trp Cys Val His Thr Ala Val Leu		
225	230	235
Leu Pro Leu Gly Gly Leu Asp Thr His Pro Ala Arg Gly Ala Thr Lys		
245	250	255
Leu Cys Pro Asp Glu Ala Arg Trp Ala Pro Arg Ser Leu Pro Leu Ser		
260	265	270
Arg Gly Val Leu Ala Ser Pro Gly Phe Ala Phe Leu Arg Ile		
275	280	285

&lt;210&gt; B9

&lt;211&gt; 1023

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; B9

```

ccacacatgga gactttgtac cgtgtcccggt tcttagtgcg cgaatgtccc aacctgaagc 60
tgaagaagcc gccctgggttg cccatgccgt cggccatgac tgtgtatgct ctggtggtgg 120
tgtcttaact cctcctcacc ggaaggaataa tttatgatgt tattgtttaa cctccaagtg 180
tcggttctat gactgatgaa catgggcac cagagccagt agctttcttg gcctacagag 240
taastggaca atatattatg gaaggacttg catccagctt cctatttaca atggggaggtt 300
taggtttcat aatcctgggc ccatcgaatg ccccaaatat cccaaaactc aatagattcc 360
ttcttctggt catcggatgc gtctgtgtgc tattgagctt ttccatggct agagtattca 420
tgagaatgaa actgcggggc tatctgatgg gtttaggtgc ctttgagagc aaateagttg 480
atcctggatt tgcctcctgc aatgaagttt taaggctgtt aaccaatcctc caatatgaa 540
tgtggaaag aatgaagagc agcagtaaaa gaatatctta ctgaasaaac agggagcgta 600
ttgaagcttg gactagaatt ccttcttggt attaaagaga caagttttat acagaatttt 660
ttttctgctt ggcctattgc tatacceaag atgtcagatg ccattttctt tttagttttt 720
cattaaaba tttcccatat ctacaactat aatatcaaat aaagtgaata ttctttaca 780
ccctcttaac attttttgga gatgaatttt ctgattttca gaatttaaca taaaatccag 840
aagcaagatt ccgttaagctg agaacctctg acagttgac agctttacat atgggtgctt 900
gcctttaact agagtgtgt atggttagatt atttcagata tgtatgtaa actgtttcct 960
gaacataag atgtatgaac ggaagcagaa taataacttt ttctaatca aaaaaaada 1020
aaa

```

1023



<210> 90  
 <211> 149  
 <212> PRT  
 <213> Homo sapiens

<400> 90  
 Met Glu Thr Leu Tyr Arg Val Pro Phe Leu Val Leu Glu Cys Pro Asn  
   1                  5                  10                  15  
 Leu Lys Leu Lys Lys Pro Pro Trp Leu His Met Pro Ser Ala Met Thr  
                   20                  25                  30  
 Val Tyr Ala Leu Val Val Val Ser Tyr Phe Leu Ile Thr Gly Gly Ile  
           35                  40                  45  
 Ile Tyr Asp Val Ile Val Glu Pro Pro Ser Val Gly Ser Met Thr Asp  
   50                  55                  60  
 Glu His Gly His Gln Arg Pro Val Ala Phe Leu Ala Tyr Arg Val Asn  
   65                  70                  75                  80  
 Gly Gln Tyr Ile Met Glu Gly Leu Ala Ser Ser Phe Leu Phe Thr Met  
                   85                  90                  95  
 Gly Gly Leu Gly Phe Ile Ile Leu Asp Arg Ser Asn Ala Pro Asn Ile  
           100                  105                  110  
 Pro Lys Leu Asn Arg Phe Leu Leu Leu Phe Ile Gly Phe Val Cys Val  
   115                  120                  125  
 Leu Leu Ser Phe Phe Met Ala Arg Val Phe Met Arg Met Lys Leu Pro  
   130                  135                  140  
 Gly Tyr Leu Met Gly  
 145

<210> 91  
 <211> 3901  
 <212> DNA  
 <213> Homo sapiens

<400> 91  
 gccatggagg gactgagcgc gctgctggcc cgtctccccc cggccggcct gccctgtggc 60  
 ctggggatca cggcgtgcgc cggggccggc gtgttgctct accggtatgc gccgaggatg 120  
 aagccaacgc acacgatggt caactgctgg ttctgcacac aggatacgt gtgacctat 180  
 ggggaaccgca actgctggga ctgtccccc tggagacagt acaacggctt ccaggagaac 240  
 ggcgactaca acaagccgat ccccgcccag tacttggagc acctgaacca cgtggtgagc 300  
 agcgcgcccc gcctgcgcga cccttcgcag ccgcagcagt ggtgagcag ccaagtccctg 360  
 ctgtgcaaga ggtgcaccca ccaccagacc accaagatca agcagctggc cgccttcgct 420  
 cctcgcgagg agggcaggtc tgacgaggag gtccagctgt accggcatca cctggagcag 480  
 atgtacaagc tgtgcgggcc gtgccaagcy gctgtggagt actacatcaa gccaccagac 540  
 cgcagctgc gccacctgtt gtcagccac cagttcaagc gccgggaggc cgcaccagac 600  
 cagcgaacga ccttctcttc cgcctggaag tcccgggtcc aggtcctct gctccgtgct 660  
 ctgccttcc tggcctgcgc ctctctactg attaccgcgc tctatggggt cagcggacac 720  
 ttgcccccag gcaccactgt gctcttggcc ctgccacctg gtggcaatgg ctacgccata 780  
 cctgacaactg gcaccacccc tggggccgag ggttggcggc agttgctggg cctactcccc 840  
 gagccatggt cggagagact gtgtgaggcc tgggctcttg ggcagagcca ccagacgggc 900  
 gtctggcac tgggctact cactgcctg ctggcaatgc tcttggctgg ccgcacagg 960

```

ctccggaggga  tccatgcctt  ctgcactgco  ctgtggggcc  tctgtctggg  gctgcacctg  1020
gctgaacagc  acctgcaggc  cgcctcgcc  agctggctaa  acaggtctca  gttcagcacc  1080
acatctttgt  gctgctgggt  tggcttcaag  gggctctgtg  ccacaaggaa  ggcaacgggc  1140
ccacggaggt  tccggcccg  aaggtcagag  aagcagccat  gactgcgggg  ggaggacaca  1200
cggatgctca  ggccaggt  ttgccaagtc  cgaagcgggc  cctctctgt  cctgctctt  1260
ttcactgct  cagctctac  cacccccacc  ctacagcccc  aggtcttgg  cttagtccct  1320
cactgcctcg  aagagtcagt  ctgccttgc  ctctctctcg  gggccaccac  agccatcccc  1380
gagtgcctg  tagccactca  ccactgctgc  cactctctcg  gccaatggcc  ctttcaactg  1440
cttggtagc  ggaatgtgg  cagcgtccac  acaggtctct  gccaatggcc  ctttcaactg  1500
agctccaggc  ccccccct  caccacggcc  ttgctgggt  ctgacactgt  tgggtgagg  1560
tctgtgtct  ggtgttcc  ctctgtggt  ctgcacagg  gtgtgacag  tggctacagg  1620
ctggggcccc  ggcgtgcct  gacgtgcag  cagagtggg  ctggggcagc  agagagccc  1680
agccctcccc  ctgaggagca  cctgtgtct  gtccctctg  tctgtctat  ggtgtgctg  1740
gacctgcagg  aggtgtggg  gctgtgaag  aggcaggct  gcagctctg  ctgtgtctg  1800
gctctctgt  ccaagacct  cctggtccc  cggaaatgg  gactgcagt  ctggggccc  1860
gctgtgctt  cgcctaggt  ttggggagcg  agaccccc  tgagacagg  agtaggagcc  1920
tgtgtgac  ttgggggact  tgagctttt  caaggttaag  gggccaggcg  tatgcaggcc  1980
ctcagtaca  ccaggtgtt  gtcactctt  cctctctga  cctgtacga  gctctgccc  2040
gtgctgtct  acctggctt  agcgcactc  ctgcttcca  ctgaggtgag  gggcagccc  2100
acagacctgc  tctcagtag  caggccctg  ccaggccct  gctgtctca  gctcagttt  2160
gacctctatg  aatgggggt  gacctctct  catagctct  ggggtgcagc  tcaagtgggt  2220
tggggatcac  atgaggtgg  tcatgaggac  ccactctgga  agtgcaggcg  ctgcccagtg  2280
cagagggaag  tccgggctg  ggggtttat  ccagggtct  cagtcagag  tggctcagg  2340
ccttccctca  ccgggcatt  tccctctgg  tgcacctc  ctgagggc  acctgtctg  2400
ccactgcga  ctctgagca  aggttacct  tccagatgt  gccaaggtgt  gccaactct  2460
ctctctgtg  cgttggacc  ccgggggag  caagagcag  tccagggtgt  atgaggtta  2520
cccccgtcac  ctgggtctg  cgggatggg  tggggggca  ggtgcaggc  ggtgcaggc  2580
tgaatctgcc  acgctgggg  gtctagagg  ctgcccacc  ccagtgttg  ggtgcaggc  2640
ccatccccc  ccagcttcc  aagtgaggaa  cccaggtgca  tggggagac  ctggggggt  2700
tctgtggct  ctgtgcctga  tgacctgct  ggtttcagg  aaggccccc  cgttaactgg  2760
ctcactgtg  tgttctgtg  ttccactgg  ttctgagtg  aggtgtggg  ggggtgagc  2820
ggggccagc  tctacctgag  ggcagggaa  ccgctgccc  gtgaggtgag  gtgaggtgag  2880
ccccatggtg  gggccctct  gagagaagg  cgtgtgtgg  gaactctgg  gcaggacatc  2940
agccacaaa  gagccctct  gctgaggtg  ctgtgtgtg  ttgtgtgtg  cccctgctg  3000
gtcagctgg  atgtgtgtg  ctgtgtgtg  ttgtgtgtg  ttgtgtgtg  cccctgctg  3060
tctgtgtgt  atgtgtgtg  ttgtgtgtg  ttgtgtgtg  ttgtgtgtg  cccctgctg  3120
ctctgtgtg  atgtgtgtg  ttgtgtgtg  ttgtgtgtg  ttgtgtgtg  cccctgctg  3180
ggtgtgtgt  agcaggtga  ggtgtgtgt  ttgtgtgtg  ttgtgtgtg  cccctgctg  3240
ttctctatg  gtcctttaa  agtttcaat  aggtgtgtg  ttgtgtgtg  cccctgctg  3300
ggcaacagct  gagaggtgc  gagaggtgc  aggtgtgtg  ttgtgtgtg  cccctgctg  3360
acccctgcca  gggactctt  cctctgctg  aggtgtgtg  ttgtgtgtg  cccctgctg  3420
ggtctgtgt  tccctctag  acttgaggt  cccctgctg  aggtgtgtg  ttgtgtgtg  3480
tccctctag  gagcaaacg  ccaggagca  gcttcaact  ttgtgtgtg  cccctgctg  3540
ccttgcctg  aggaacacat  gacctctct  tctgtgtgt  actgtgtgt  cccctgctg  3600
ctgctcagg  aggaacacat  gacctctct  tctgtgtgt  actgtgtgt  cccctgctg  3660
gtctgtgag  ttgtgtgtg  ttgtgtgtg  ttgtgtgtg  ttgtgtgtg  cccctgctg  3720
cctagatgt  ggtttttgg  ttgtgtgtg  ttgtgtgtg  ttgtgtgtg  cccctgctg  3780
ccacacacac  aaaaaaaaa  aaaaaaaaa  aaaaaaaaa  aaaaaaaaa  aaaaaaaaa  3840
a

```

3901

&lt;210&gt; 92

&lt;211&gt; 392

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 92

```

Met  Glu  Gly  Val  Ser  Ala  Leu  Leu  Ala  Arg  Cys  Pro  Thr  Ala  Gly  Leu
  1              5              10              15

```

Ala Gly Gly Leu Gly Val Thr Ala Cys Ala Ala Ala Gly Val Leu Leu  
                     20                    25                    30  
 Tyr Arg Ile Ala Arg Arg Met Lys Pro Thr His Thr Met Val Asn Cys  
                     35                    40                    45  
 Trp Phe Cys Asn Gln Asp Thr Leu Val Pro Tyr Gly Asn Arg Asn Cys  
                     50                    55                    60  
 Trp Asp Cys Pro His Cys Glu Gln Tyr Asn Gly Phe Gln Glu Asn Gly  
                     65                    70                    75                    80  
 Asp Tyr Asn Lys Pro Ile Pro Ala Gln Tyr Leu Glu His Leu Asn His  
                     85                    90                    95  
 Val Val Ser Ser Ala Pro Ser Leu Arg Asp Pro Ser Gln Pro Gln Gln  
                     100                    105                    110  
 Trp Val Ser Ser Gln Val Leu Leu Cys Lys Arg Cys Asn His His Gln  
                     115                    120                    125  
 Thr Thr Lys Ile Lys Gln Leu Ala Ala Phe Ala Pro Arg Glu Glu Gly  
                     130                    135                    140  
 Arg Tyr Asp Glu Glu Val Glu Val Tyr Arg His His Leu Glu Gln Met  
                     145                    150                    155                    160  
 Tyr Lys Leu Cys Arg Pro Cys Gln Ala Ala Val Glu Tyr Tyr Ile Lys  
                     165                    170                    175  
 His Gln Asn Arg Gln Leu Arg Ala Leu Leu Leu Ser His Gln Phe Lys  
                     180                    185                    190  
 Arg Arg Glu Ala Asp Gln Thr His Ala Gln Asn Phe Ser Ser Ala Val  
                     195                    200                    205  
 Lys Ser Pro Val Gln Val Ile Leu Leu Arg Ala Leu Ala Phe Leu Ala  
                     210                    215                    220  
 Cys Ala Phe Leu Leu Thr Thr Ala Leu Tyr Gly Ala Ser Gly His Phe  
                     225                    230                    235                    240  
 Ala Pro Gly Thr Thr Val Pro Leu Ala Leu Pro Pro Gly Gly Asn Gly  
                     245                    250                    255  
 Ser Ala Thr Pro Asp Asn Gly Thr Thr Pro Gly Ala Glu Gly Trp Arg  
                     260                    265                    270  
 Gln Leu Leu Gly Leu Leu Pro Glu His Met Ala Glu Lys Leu Cys Glu  
                     275                    280                    285  
 Ala Trp Ala Phe Gly Gln Ser His Gln Thr Gly Val Val Ala Leu Gly  
                     290                    295                    300  
 Leu Leu Thr Cys Leu Leu Ala Met Leu Leu Ala Gly Arg Ile Arg Leu  
                     305                    310                    315                    320  
 Arg Arg Ile Asp Ala Phe Cys Thr Cys Leu Trp Ala Leu Leu Leu Gly  
                     325                    330                    335

Leu His Leu Ala Glu Gln His Leu Gln Ala Ala Ser Pro Ser Trp Leu  
340 345 350

Asn Thr Leu Lys Phe Ser Thr Thr Ser Leu Cys Cys Leu Val Gly Phe  
355 360 365

Thr Ala Ala Val Ala Thr Arg Lys Ala Thr Gly Pro Arg Arg Phe Arg  
370 375 380

Pro Arg Arg Ser Glu Lys Gln Pro  
385 390

<210> 93

<211> 2203

<212> DNA

<213> Homo sapiens

<400> 93

```

cagcgggtggg aggtgggggac cagccgggttg agggcccccagg ctgggcccctca ccacaaatgtg 60
gcacgagggtt cgggaagcatg agcgggaagct tcgggggcctg atggtcgact acaagaagag 120
ggcggagcggg agacggggagt attatgaana gatcaagaag gacccagccc agttcctgca 180
ggtaacatggc cggagcttgca aggtgcacct ggaattctgca gtcccctggg ccgctcgagag 240
ccctgttaat atgatgcctt ggcaggggga caaccaaac atgattgacc gattcgatgt 300
ccgtgccccac ctggaccaca tcccggacta cacccccct ctgctacca ccactctccc 360
agacacaggag tcggacgcac ggaagtgtac ctacgagcgc tacgagggcc tgggtgcagaa 420
cgactttgcc ggcattctag agcagcagtg cctgtaccag atctacattg atgagctgtg 480
cggaggccctc cagagaccac gcaagatga gaaagagaag ctggcagaga agaaagctto 540
catcggttat acctacgagg acagcacggt ggcggaggta gagaaggcgg cagaaaagcc 600
agaggaggag gagtccagcg ccgaggagga gagaacctcg gacgaagatg aggtcatccc 660
cgacatcgac gtggaggttg acgtggatga attgaaccag gaggaggttg negatctcaa 720
caaacaggcc acgacttatg gcctggccga caggctctt gaggaggana aggccatgta 840
caaggaggag cgtctctgac gccagtcgag agagtttcgg gagaaaggcc tgaagggtcg 900
ctcgggagcg ccacccagct atgcttcgag agacagccc acctatgacc cctataagcg 960
gbcacccctg gagtccagct cagagtcctg ctcccgtctc cgttcccga ccccgggcgg 1020
cgaggagag ataacgttc taccagttt tgggggcaga gatgaggagg cagcccgagc 1080
cgtctctgca gcagcagcat caggagtcac caccaggag ccccccgcac ctccccagcc 1140
tggcggcccc gccccgggac gtaatgctag cgcgcgcgc cgtctctctt ctctcctctc 1200
ctcctcttat gctcagagg catccagctc cctccacggt agccacggtt cgtctccggt 1260
tcgcctggtt gggggtatct accgttccgg ccggcacggt cgtctccggt cccgtctctg 1320
gtcccgtctc cgtctccgct cccggcgcta ttctcgttc cgtacgctg gccggcgga 1380
ctcagctggg ggtctccgag acgggacccg gtactccctc tgcgcgcgc cggctggttg 1440
ttacggggcc cggcgagaa gccggagccg ctcccactca ggggaccggt acagccgggg 1500
cgcccggggc ctacggcacc acagccagtg ccgcagctgc agcagctggt cctcagccc 1560
gtcccgcagt cgcagctga cccagagccc ctgcacatca cccgcaagag agaagctgac 1620
cagccgcagc cgcagccga gccagagccc ctgcacatca cccgcaagag agaagctgac 1680
taggcggccc gctctctctg ctgtggcgga gaaagtgaaa agacccgac ctgcctctgg 1740
taaaagagaa ggaagtgcac aacccaagct gacgctctag gagaagctga aactgagat 1800
gcagaaggcg ctgacacagg agttcaaggc ggaatagaag ggggacacag aaagatgat 1860
cacgcaggag catgagcggc agggagcgga agcagagctt cagcccatgg cccgcaagat 1920
ccgcattgag gaggcggaac gccagagaga ggagagaga gagtgggaat gccagtacag 1980
ccggcagagc cgtccacctt ccccgcgata cagtgcagaa taccgctctt ctgcaggcg 2040
ctcaaggctc cgtcccgaa gccctcatta ccgacattag gcagaagagt ggggggtggg 2100
ggggcagag ggtgggttaa ggggtcagag ctgtgatgct gctggtttta tctctagtga 2160
aatbaagtc aaggtattt aattccgctc aaacaaana aas
2203

```

<210> 94

<211> 674

<212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 94

```

Met Trp His Glu Ala Arg Lys His Glu Arg Lys Leu Arg Gly Met Met
 1              5              10              15

Val Asp Tyr Lys Lys Arg Ala Glu Arg Arg Arg Glu Tyr Tyr Glu Lys
      20              25              30

Ile Lys Lys Asp Pro Ala Gln Phe Leu Gln Val His Gly Arg Ala Cys
      35              40              45

Lys Val His Leu Asp Ser Ala Val Ala Leu Ala Ala Glu Ser Pro Val
      50              55              60

Asn Met Met Pro Trp Gln Gly Asp Thr Asn Asn Met Ile Asp Arg Phe
      65              70              75              80

Asp Val Arg Ala His Leu Asp His Ile Pro Asp Tyr Thr Pro Pro Leu
      85              90              95

Leu Thr Thr Ile Ser Pro Glu Gln Glu Ser Asp Glu Arg Lys Cys Asn
      100              105              110              115

Tyr Glu Arg Tyr Arg Gly Leu Val Gln Asn Asp Phe Ala Gly Ile Ser
      115              120              125

Glu Glu Gln Cys Leu Tyr Gln Ile Tyr Ile Asp Glu Leu Tyr Gly Gly
      130              135              140

Leu Gln Arg Pro Ser Glu Asp Glu Lys Lys Lys Leu Ala Glu Lys Lys
      145              150              155              160

Ala Ser Ile Gly Tyr Thr Tyr Glu Asp Ser Thr Val Ala Glu Val Glu
      165              170              175

Lys Ala Ala Glu Lys Pro Glu Glu Glu Glu Ser Ala Ala Glu Glu Glu
      180              185              190

Ser Asn Ser Asp Glu Asp Glu Val Ile Pro Asp Ile Asp Val Glu Val
      195              200              205

Asp Val Asp Glu Leu Asn Gln Glu Gln Val Ala Asp Leu Asn Lys Gln
      210              215              220

Ala Thr Thr Tyr Gly Met Ala Asp Gly Asp Phe Val Arg Met Leu Arg
      225              230              235              240

Lys Asp Lys Glu Glu Ala Glu Ala Ile Lys His Ala Lys Ala Leu Glu
      245              250              255

Glu Glu Lys Ala Met Tyr Ser Gly Arg Arg Ser Arg Arg Gln Arg Arg
      260              265              270

Glu Phe Arg Glu Lys Arg Leu Arg Gly Arg Lys Ile Ser Pro Pro Ser
      275              280              285

Tyr Ala Arg Arg Asp Ser Pro Thr Tyr Asp Pro Tyr Lys Arg Ser Pro
      290              295              300

```

Ser Glu Ser Ser Ser Glu Ser Arg Ser Arg Ser Arg Ser Pro Thr Pro  
 305 310 315 320  
 Gly Arg Glu Glu Lys Ile Thr Phe Ile Thr Ser Phe Gly Gly Ser Asp  
 325 330 335  
 Glu Glu Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ser Gly Val Thr  
 340 345 350  
 Thr Gly Lys Pro Pro Ala Pro Pro Gln Pro Gly Gly Pro Ala Pro Gly  
 355 360 365  
 Arg Asn Ala Ser Ala Arg Arg Arg Ser Ser Ser Ser Ser Ser Ser Ser  
 370 375 380  
 Ser Ala Ser Arg Thr Ser Ser Ser Arg Ser Ser Ser Arg Ser Ser Ser  
 385 390 395 400  
 Arg Ser Arg Arg Gly Gly Gly Tyr Tyr Arg Ser Gly Arg His Ala Arg  
 405 410 415  
 Ser Arg Ser Arg Ser Trp Ser Arg Ser Arg Ser Arg Ser Arg Arg Tyr  
 420 425 430  
 Ser Arg Ser Arg Ser Arg Gly Arg Arg His Ser Gly Gly Gly Ser Arg  
 435 440 445  
 Asp Gly His Arg Tyr Ser Arg Ser Pro Ala Arg Arg Gly Gly Tyr Gly  
 450 455 460  
 Pro Arg Arg Arg Ser Arg Ser Arg Ser His Ser Gly Asp Arg Tyr Arg  
 465 470 475 480  
 Arg Gly Gly Arg Gly Leu Arg His His Ser Ser Ser Arg Ser Arg Ser  
 485 490 495  
 Ser Trp Ser Leu Ser Pro Ser Arg Ser Arg Ser Leu Thr Arg Ser Arg  
 500 505 510  
 Ser His Ser Pro Ser Pro Ser Gln Ser Arg Ser Arg Ser Arg Ser Arg  
 515 520 525  
 Ser Gln Ser Pro Ser Pro Ser Pro Ala Arg Glu Lys Leu Thr Arg Pro  
 530 535 540  
 Ala Ala Ser Pro Ala Val Gly Glu Lys Leu Lys Lys Thr Glu Pro Ala  
 545 550 555 560  
 Ala Gly Lys Glu Thr Gly Ala Ala Lys Pro Lys Leu Thr Pro Gln Glu  
 565 570 575  
 Lys Leu Lys Leu Arg Met Gln Lys Ala Leu Asn Arg Gln Phe Lys Ala  
 580 585 590  
 Asp Lys Lys Ala Ala Gln Glu Lys Met Ile Gln Gln Glu His Glu Arg  
 595 600 605  
 Gln Glu Arg Glu Asp Glu Leu Arg Ala Met Ala Arg Lys Ile Arg Met  
 610 615 620

Lys Glu Arg Glu Arg Arg Glu Lys Glu Arg Glu Glu Trp Glu Arg Gln  
625 630 635 640

Tyr Ser Arg Gln Ser Arg Ser Pro Ser Pro Arg Tyr Ser Arg Glu Tyr  
645 650 655

Ser Ser Ser Arg Arg Arg Ser Arg Ser Arg Ser Arg Ser Pro His Tyr  
660 665 670

Arg His

<210> 95

<211> 1014

<212> DNA

<213> Homo sapiens

<400> 95

```

gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 60
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 120
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 180
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 240
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 300
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 360
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 420
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 480
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 540
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 600
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 660
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 720
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 780
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 840
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 900
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 960
gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg gggggggggg 1014

```

<210> 96

<211> 204

<212> PRF

<213> Homo sapiens

<400> 96

```

Met Gly Ala Val Thr Asp Asp Glu Val Ile Arg Lys Arg Leu Leu Ile
  1          5          10          15

Asp Gly Asp Gly Ala Gly Asp Asp Arg Arg Ile Asn Leu Leu Val Lys
 20          25          30

Ser Phe Ile Lys Trp Cys Asn Ser Gly Ser Gln Glu Glu Gly Tyr Ser
 35          40          45

Gln Tyr Gln Arg Met Leu Ser Thr Leu Ser Gln Cys Glu Phe Ser Met
 50          55          60

Gly Lys Thr Leu Leu Val Tyr Asp Met Asn Leu Arg Glu Met Glu Asn
 65          70          75          80

Tyr Glu Lys Ile Tyr Lys Glu Ile Glu Cys Ser Ile Ala Gly Ala His
 85          90          95

```

Glu Lys Ile Ala Glu Cys Lys Lys Gln Ile Leu Gln Ala Lys Arg Ile  
 100 105 110  
 Arg Lys Asn Arg Gln Glu Tyr Asp Ala Leu Ala Lys Val Ile Gln His  
 115 120 125  
 His Pro Asp Arg His Glu Thr Leu Lys Glu Leu Glu Ala Leu Gly Lys  
 130 135 140  
 Glu Leu Glu His Leu Ser His Ile Lys Glu Ser Val Glu Asp Lys Leu  
 145 150 155 160  
 Glu Leu Arg Arg Lys Gln Phe His Val Leu Leu Ser Thr Ile His Glu  
 165 170 175  
 Leu Gln Gln Thr Leu Glu Asn Asp Glu Lys Leu Ser Glu Val Glu Glu  
 180 185 190  
 Ala Gln Glu Ala Ser Met Glu Thr Asp Pro Lys Pro  
 195 200

<210> 97  
 <211> 955  
 <212> DNA  
 <213> Homo sapiens

<400> 97  
 aatcctcaac aaaatagtag caaaccaaat ccaatagtaa atcaccgaaga taatacagta 60  
 tgcataaatg ggaatttaatt caaggatgca tagatgattc aacattcaag atcacaataa 120  
 ttattctctg taatttttca taaggtatgt ttttaaatca gaatgggtatt gtatattaga 180  
 cataaatga cgtttttagt tagcattctt agagctagct tcataatcca attaatatac 240  
 ttgcacactg agtgcaggty ttttaatttt tataactgta tctgttatgc tattcaaatg 300  
 agctaattgt agttaktctt ataccattg geattgggtt ccatagkata cataegtttt 360  
 atttttgttt tctctgttag accttcaaat atttaacttct catagtttct ctggcataaa 420  
 agctcccaag ttccatcttc aacagtccag gtcttgggat atctatcgtt ttatttgttt 480  
 ttataacttt tatttagaag agtcaatcc ttttttagct atttaagat aaaaagtcca 540  
 ctttttccac ttttgtcttc gattgaattg ctgcacctaa catggakcta tcttgggttt 600  
 acagaagaa gagaagagga agatattgaa gagaagaat cpattaagaa aaaaattaaa 660  
 gaacttaagt ttttagatcc caaacttgc cagaaacttt gtaaglatca tattccaata 720  
 ccattcaagg acagtggaa tatltcttca aatgatttca tttttctaa gaccgattat 780  
 tcattacttg ctatlttcat ttgtttatta tatgcctgat aattccacag ataactctcc 840  
 tttaggtaaa ttatgggatt aactgcttca aaagataaag gcataattaga aaatacaaat 900  
 aagaagaggt ttttaaatga aattctacct ttcataactg aaaaaaaaaa aaaaa 955

<210> 98  
 <211> 97  
 <212> PRT  
 <213> Homo sapiens

<400> 98  
 Met Ile Lys Ser Ser Leu Phe Pro Leu Leu Tyr Leu Asn Glu Leu Leu  
 1 5 10 15  
 Pro Leu Thr Trp Ile Tyr Leu Gly Phe Thr Glu Arg Arg Glu Glu Glu  
 20 25 30  
 Asp Ile Glu Glu Lys Lys Ser Ile Lys Lys Lys Ile Lys Glu Leu Lys  
 35 40 45



Phe Leu Asp Ser Lys Ile Ala Gln Asn Leu Cys Lys Tyr His Ile Pro  
50 55 60

Ile Pro Phe Lys Asp Ser Gly Asn Ile Ser Leu Asn Asp Phe Ile Phe  
65 70 75 80

Phe Lys Thr Asp Tyr Ser Leu Phe Ala Ile Phe Ile Leu Leu Leu Tyr  
85 90 95

Ala

<210> 99

<211> 1375

<212> DNA

<213> Homo sapiens

<400> 99

```

gtctctcttt agggagcagg agtgcctctg gtaattgagg gtggatgttg tgtgtgctgg 60
ggaggggtcc ttctgtttgg tctacacctt gtctactctg cccctggatg gtgggggttg 120
ctttctctac ccccaactc cctgctcagg tctctgtgtt gccctgcatg cccaggcttg 180
tgagccnagg tctcttttgg ggcagggagt agcagcaggt ggggggggtt ccccatcagg 240
ccttgcaggt ccccaactca ggcctctgga aggtccaggg atgggtctctg atgagagggg 300
aaaagtgtgt cagggaacaa caggcctcag ctgcttagag gacctcccc ctgccttgca 360
gtgggctcgg gtagagcagt atcaggagct aggttgtctt gctgcccaca ctctgtctt 420
ttgggtatcc taactgtctc ggaggaggtt gacatcccc ttctgggtca tgtgtctgac 480
accaacaaca tgggtctctgt cctctctctt ttgactctcc ctttgtcttc cccatagagg 540
tggggtgggg tggatcccta cacttggggc aggcagcccc aaagtggggg agggggatgg 600
cagagactgt aaaggcgcca ctggactctg gcaaggcctt tattaccttc actccccctc 660
ctctccctc accagctca aggcctgagg gctgcagggg ctcttgccag ctactgggtg 720
aggtctctct gcaacagact ccccttcttc ctggcaccac ctcttccctt ttgagagaga 780
cagcaacagg cgtagcaaaa gcagctgctg ctctgtctat gagggtgtat atatttttta 840
cccgaagctc tggacttgta catlakttt ttcaaacctc aagagggaaa gaggcttgta 900
tcatatgtga acattgtctc ataggtaatg ttgtacagac cctttctatac agtgatctgt 960
cttgttctct cagcaaacat cctctatgga cataggaggt gctgtgtccc atgcctctt 1020
gccctgacag tctcccatgg gccctctctt gctccctgcc cctccctgct tactgtgtgt 1080
gcctctctct ctccctgacg cccctggctt cccagccttc ctctgaccc ctcccaacag 1140
ccttgggaat cagctgcca cccctctg ctggcaccac tgggaaccaa tggcccagtc 1200
ttggctgtgt cttaccccta gccttgatgc ctgcccaggg acccccagcc cctcccggtt 1260
gccctgcagg tttaacagag tgaacctgtt gtattgtaca ggcgcggttg tcattgcaga 1320
aacgcgtggg tggagaagaa gcgataaag tctatgactc aaaaaa 1375

```

<210> 100

<211> 132

<212> PRT

<213> Homo sapiens

<400> 100

Met Ala Glu Thr Val Lys Ala Pro Leu Asp Ser Gly Lys Ala Phe Ile  
1 5 10 15

Thr Phe Thr Pro Leu Pro Leu Pro Ser Pro Ala Ser Arg Pro Glu Gly  
20 25 30

Cys Arg Gly Ser Trp Gln Leu Leu Gly Glu Val Ser Trp His Arg Leu  
35 40 45

Thr Leu Leu Ser Gly Thr Thr Ser Phe Pro Phe Glu Glu Thr Ala Thr

50                      55                      60  
 Ala Val Ala Lys Ala Ala Ala Pro Ala Met Arg Val Tyr Ile Phe  
 65                      70                      75                      80  
 Phe Thr Gln Ser Ser Gly Ile Val His Leu Phe Phe Lys Thr Gln Arg  
 85                      90                      95  
 Gly Lys Glu Pro Cys Ile Ile Cys Glu His Cys Ile Ile Gly Asn Val  
 100                      105                      110  
 Val Gln Thr Leu Leu Tyr Ser Asp Leu Ser Cys Ser Cys Ser Lys Asn  
 115                      120                      125  
 Pro Leu Trp Thr  
 130

<210> 101  
 <211> 1213  
 <212> DNA  
 <213> Homo sapiens

<400> 101  
 ggcttcaggc tgaagtcctc gcttcctcca gttcctcaag ggttaggtag ggctcctgc 60  
 atcaccctca gaatccagtt ccaaccccca ctctccttag gccttgctct ctgctctgcc 120  
 ctgcacaggt gcccttgctc atgtcagtag catgggcggg tgggtgggac ggcagtggtg 180  
 atgaaggggg tgcaccacag gccctcatga gcagcttcca catgggcgtg tggctggggc 240  
 gtggccacca cagagcaccat ggctgtgtct aggcgcagc accttagcag tatctgttta 300  
 catgcpcaag gatcaagccg actacctgtg ctgtctactg ggacagcagc ctccgagcta 360  
 ctccgtacct cctctctcca ggtcgtggag ttaggctcca gtccttactt gtccctggtt 420  
 ccactgtgc tctaacctgt ccagccacctg ggagctcttg cctggggctc gaggccctgg 480  
 taggagctgc agttggaggg cgttctgtgc ccagcagcgg tggagcggct ccctgggccc 540  
 tgtctctgca gggagccagg gctgcggcac atgtgctgtg aaactggcac ccactggcg 600  
 tgctgtctgc gccacttgt tctgcagca cctctaccc tgcctcgtgt cctccctctc 660  
 ccgcgccttg gctcaggagt gctgganaag ctacagcctc ggcttgggag cctggcctct 720  
 tgatatacct cagagcttcc ctgtgctccc cagcccccagg accactggcc ccttggcctg 780  
 aggggctcgg ggcaccacga cctgtcagct caggtccggg agagagcccg gacggcgctg 840  
 ccatctcggc tgggcttgc tgagagcctc cgtcttgggt ttctcctgt ctgagattca 900  
 tggctcactg tgggtctaca cagctagaat agatatactt agagagagag atattttta 960  
 gacaaagccc acaattagct gtcctttaac accctccag aagaagagcg 1020  
 atccctcgga cggtcggggc gggcaccctc agccgggctc ttgacagag cagcaccgt 1080  
 gactgtgggc ccggccctca gatgtgtaca tatacggcta ttctctatt tactgttctt 1140  
 cagattttagt actgtaat aaacacacac attagggaga gattanacat ttttgctaaa 1200  
 aaaaaaaaaa aa 1213

<210> 102  
 <211> 108  
 <212> PRT  
 <213> Homo sapiens

<400> 102  
 Met Lys Gly Val His His Arg Pro His Glu Ala Val Pro Thr Trp Ala  
 1                      5                      10                      15  
 Cys Gly Trp Gly Val Ala Thr Thr Glu His Met Ala Val Ser Arg Arg  
 20                      25                      30  
 Lys His Phe Ser Ser Ile Cys Leu His Ala Gln Gly Ser Ser Arg Leu  
 35                      40                      45

Pro Val Leu Ser Thr Gly Thr Ala Val Ser Glu Leu Leu Arg Thr Ser  
 50 55 60

Leu Cys Gln Val Val Glu Leu Gly Pro Ser Pro Tyr Leu Ser Leu Val  
 65 70 75 80

Pro Thr Val Leu Leu Thr Val Gln His Leu Gly Ala Leu Ala Trp Gly  
 85 90 95

Trp Arg Pro Trp  
 100

<210> 103  
 <211> 1036  
 <212> DNA  
 <213> Homo sapiens

<400> 103  
 cctcaaatgc tttctttctt cagatgcttt tctgtgtaca tgatactagt agacactttt 60  
 ctcttttatat ttactgatag tgaasatcat agcaataaa atattgatgt ttgaaggcag 120  
 tggtaaccaa ttggtcaaaa aactatgaaa tglasactga attgttatat ctctatcctt 180  
 tttgttttct tctgtgtttt taatgtatgg aataatctc ataatagaa agaaaaataa 240  
 cttagaatt ttccaaagct agtactcttt ctcttataa atgtacataa ttctaatctt 300  
 ttacaaaatt takttaactg taactactgt acttattgtt gattcaatga cgcagttacg 360  
 tcaatcaca aaagtattatg aattcgagat tactgaactg tttcttctat attgcattca 420  
 catcaatatt tglgaatttg ttgttcagct ttctattcaa acaaaaataa ttccctcaag 480  
 aaagctccat ttttatcata aacatttcaa cataactaac attagaacaa gtctgccatg 540  
 ttcaaaataa tttaagact tatctctgaa aacggtatcc agaaagcag gtgttcccag 600  
 taatgtagct tcaaaaataa aatgtgctat ttatatgaca tgaattcat aacttttggg 660  
 aggttatatt tatgacagca caaaaaataa attctgtgtt ataaagaaga tccaaacaa 720  
 taacatata agcaagaaa alagagaac acagttattg aatctactct tgtcaattac 780  
 attttcaaaa acaaaaatgc atattgtaat atttggtaca tgaacttgc atgttgat 840  
 gctatatac ttcaaaagta ttcaatgtgt acttagagac gottaaataa tgtcatgtac 900  
 aactcttata aacatttita cagggttccc atttgcactt catctttcag taaagtcttg 960  
 tcagaaabaa attgtctgat aatatcgaa aataaaatt tgaattttag ttaaaaaaa 1020  
 aaaaaaaa aaaaaa 1036

<210> 104  
 <211> 87  
 <212> PRT  
 <213> Homo sapiens

<400> 104  
 Met Tyr Thr Ile Leu Ile Phe Leu Gln Ile Tyr Leu Thr Val Pro Thr  
 1 5 10 15

Val Leu Ile Val Asp Ser Met Thr Gln Leu Ser His His Pro Arg Ile  
 20 25 30

Tyr Glu Phe Glu Ile Thr Asp Leu Phe Ser Ser Tyr Cys Ile His Ile  
 35 40 45

Asn Ile Cys Glu Phe Val Val Gln Leu Phe Ile Gln Thr Lys Asn Ile  
 50 55 60

Pro Ser Arg Lys Leu His Phe Tyr His Lys His Phe Asn Ile Thr Asn  
 65 70 75 80

Ile Arg Thr Ser Leu Pro Cys

85

&lt;210&gt; 105

&lt;211&gt; 2349

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 105

```

tttttttttt tttttgggtc ctttggtaaaa ttttatccaa aaaaacaggat acatataatat 60
tttagagagg aaatatgaas tcaagagttt tggcagcccc tgcttttttt ttttttttag 120
ctccctaagg actgtagcag gataaaagga tcaatggctc cggctctctt tgagataaca 180
agtgatgaan taaaaaagaa agccctatcc ctcaaatuag gtcaggtaac cccattggcc 240
acccctcccta caaggtaaaa aatgagtact tttagtaaca gttcagaatt catcttttct 300
tcctacctgc ctcatcgggt gaagttttaa gtcatgattt ttttttagaa ttgtactctg 360
tgtctataga caaatgaact catatttagc agccaaagag gcttaaccct gctgcaccag 420
cagtatacct caagcactgc ctcaacctcg cccctgcgcc cagatgcttc tgttganaag 480
tcaaccgagg agcagctac ccaggctccc agtctggaga gtctgacttt aaagctagag 540
caagaggttg tggccaggag ccgaaccaac atgagatgct agtatcccc 600
tctgatacta cccctctggt ttcccgagct gtccaccag tcnaactgga ggtatgagat 660
gattcggaact ctgagctgga cttagcgcaag ctgtaccact cttctttctt tctctatcc 720
tcctccagct ccagctccag cactgatgag agtgaggatg gaaaggagga gaagctaat 780
gaacagtcct gctcaagact ctatgatgan gagagctctc tctccctcac tatgtcccaa 840
gatggatttc caaatgaaga tggagaacaa atgacccctg agtctctgct actgcaggaa 900
agacaaagag cctctgagtg gcccaaggat cgtgtctctg taacccgtat tgaactctgt 960
tgcagagctg tactctcagg gaagtctgct tctagcccta gggcccgga aatggttaaa 1020
ggaggaattt tggggccagg caaccacttg cttagcagtc cctcattgac tcttgagaa 1080
tatggctgact ctccagtcct caccaccaga agtagtagtg capcttccat ggcagagga 1140
gaagcctctg cagtccagac agtggcagcc cagttcacca aacttcgcc agcatggat 1200
gaaaaggagt ttacagttaa aatcaaaagt gagggaagat tgaagttaac attccagaag 1260
cacaaagtga tggcgaatgg agtaatggga gatggacatc cactgtttca taagaagaag 1320
gggaacagaa agaacctagt agagctggag gtggagtgca tggaaagacc taatcaccct 1380
gatgtggccc tggagacccc gatccctgtc atcaataagg tggatggtac tttgctggtg 1440
ggtgaggatg cccctctgcc ggtgnaactg gagatgtggt tacagagctc tccagagttt 1500
gctgttgatc cccgatttct agcgtatctg gaggatcgca gaaacagaa gtggcaaga 1560
tgtaaaaaaa ataatgaagg agaatgaac tgtttgggaa tggaaaccag acagacagct 1620
aaactctgaa atgggaaaaa gggtcctcac actgaaaacc tpttcaacc ggttttgcac 1680
gggcttattg caccagagag cagcaagaa ggggtgaaag accgggtctc tatctctgca taacacgttc 1740
aagatgatgg cctcctgca ggttgaaagc accgggtctc tatctctgca taacacgttc 1800
caacacagca gtagtgacct acagtctgtg tcatcttttg gtcacagtag tgcacttct 1860
gcacttttgc ctttttatgc atttgtgttg ggtggtgca cactccccc catcctccc tctgtagac 1920
tccagcacca tgcctcatca ccacacacc cactccccc cccaccatc ccacatcac 1980
catccaggct tgagagcccc tgcctacccc tcttcaccag tgaataccg cctcgttact 2040
accttgcggt tcccacccct gcaacctgag aggatgagc atgaggatga agaagatgat 2100
gatgacttat ctcaaggcta tgatagctcc gaaagggact tctcactcat tgatgatct 2160
atgatgccag ctcaactaga ctccagtgaa gtgtgtgat actgaagccc cagcatgggc 2220
ccattgctt gggcggctgc tctatttctc tttactcttg ccttggact atggaacgt 2280
gggaggggca ggggagctgt ggggaagctc aggaactccag gaggtgaan ggaasaaaa 2340
aaaaaanaa

```

2349

&lt;210&gt; 106

&lt;211&gt; 539

&lt;212&gt; gnt

&lt;213&gt; Homo sapiens

&lt;400&gt; 106

```

Met Arg Val Ser Pro Ser Asp Thr Thr Pro Leu Val Ser Arg Ser Val
1 5 10 15

```

Pro Pro Val Lys Leu Glu Asp Glu Asp Asp Ser Asp Ser Glu Leu Asp  
 20 25 30  
 Leu Ser Lys Leu Ser Pro Ser Ser Ser Ser Ser Ser Ser Ser Ser  
 35 40 45  
 Ser Ser Ser Ser Thr Asp Glu Ser Glu Asp Glu Lys Glu Glu Lys Leu  
 50 55 60  
 Thr Asp Glu Ser Arg Ser Lys Leu Tyr Asp Glu Glu Ser Leu Leu Ser  
 65 70 75 80  
 Leu Thr Met Ser Gln Asp Gly Phe Pro Asn Glu Asp Gly Glu Gln Met  
 85 90 95  
 Thr Pro Glu Leu Leu Leu Leu Gln Glu Arg Gln Arg Ala Ser Glu Trp  
 100 105 110  
 Pro Lys Asp Arg Val Leu Ile Asn Arg Ile Asp Leu Val Cys Gln Ala  
 115 120 125  
 Val Leu Ser Gly Lys Trp Pro Ser Ser Arg Arg Ser Gln Glu Met Val  
 130 135 140  
 Thr Gly Gly Ile Leu Gly Pro Gly Asn His Leu Leu Asp Ser Pro Ser  
 145 150 155 160  
 Leu Thr Pro Gly Glu Tyr Gly Asp Ser Pro Val Pro Thr Pro Arg Ser  
 165 170 175  
 Ser Ser Ala Ala Ser Met Ala Glu Glu Glu Ala Ser Ala Val Ser Thr  
 180 185 190  
 Ala Ala Ala Gln Phe Thr Lys Leu Arg Arg Gly Met Asp Glu Lys Glu  
 195 200 205  
 Phe Thr Val Gln Ile Lys Asp Glu Glu Gly Leu Lys Leu Thr Phe Gln  
 210 215 220  
 Lys His Lys Leu Met Ala Asn Gly Val Met Gly Asp Gly His Pro Leu  
 225 230 235 240  
 Phe His Lys Lys Lys Gly Asn Arg Lys Lys Leu Val Glu Leu Glu Val  
 245 250 255  
 Glu Cys Met Glu Glu Pro Asn His Leu Asp Val Asp Leu Glu Thr Arg  
 260 265 270  
 Ile Pro Val Ile Asn Lys Val Asp Gly Thr Leu Leu Val Gly Glu Asp  
 275 280 285  
 Ala Pro Arg Arg Ala Glu Leu Glu Met Trp Leu Gln Gly His Pro Glu  
 290 295 300  
 Phe Ala Val Asp Pro Arg Phe Leu Ala Tyr Met Glu Asp Arg Arg Lys  
 305 310 315 320  
 Gln Lys Trp Gln Arg Cys Lys Lys Asn Asn Lys Ala Glu Leu Asn Cys  
 325 330 335

Leu Gly Met Glu Pro Val Gln Thr Ala Asn Ser Arg Asn Gly Lys Lys  
 340 345 350  
 Gly His His Thr Glu Thr Val Phe Asn Arg Val Leu Pro Gly Pro Ile  
 355 360 365  
 Ala Pro Glu Ser Ser Lys Lys Arg Ala Arg Arg Met Arg Pro Asp Leu  
 370 375 380  
 Ser Lys Met Met Ala Leu Met Gln Gly Gly Ser Thr Gly Ser Leu Ser  
 385 390 395 400  
 Leu His Asn Thr Phe Gln His Ser Ser Ser Gly Leu Gln Ser Val Ser  
 405 410 415  
 Ser Leu Gly His Ser Ser Ala Thr Ser Ala Ser Leu Pro Phe Met Pro  
 420 425 430  
 Phe Val Met Gly Gly Ala Pro Ser Ser Pro His Val Asp Ser Ser Thr  
 435 440 445  
 Met Leu His His His His His His Pro His Pro His His His His His  
 450 455 460  
 His His Pro Gly Leu Arg Ala Pro Gly Tyr Pro Ser Ser Pro Val Thr  
 465 470 475 480  
 Thr Ala Ser Gly Thr Thr Leu Arg Leu Pro Pro Leu Gln Pro Glu Glu  
 485 490 495  
 Asp Asp Asp Glu Asp Glu Glu Asp Asp Asp Asp Leu Ser Gln Gly Tyr  
 500 505 510  
 Asp Ser Ser Glu Arg Asp Phe Ser Leu Ile Asp Asp Pro Met Met Pro  
 515 520 525  
 Ala Asn Ser Asp Ser Ser Glu Asp Ala Asp Asp  
 530 535

&lt;210&gt; 107

&lt;211&gt; 3004

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 107

```

ggggcatgag catctcaggg ctgccagaat ggccttttgc cagtgcatag caccagcgtg 60
tgtcatgtct tggctgcgtt tctggggccc atggcccttc cttaactggc aactattgtc 120
tttactegtc aaggaaggtc agcctctggt gtgggtccag gacccgctcc agctgaactc 180
caaccccttg gggccacctg agccttggtc ttctcgatcc tcccaatcc catgggaatc 240
tccccatgca cctgctcccc cagcagcccc gggggacttt gattacctgg agcctctgca 300
ttcttcgcaag atgtcagccc tgcctcagga actcaactga aatttggttc catctctgaa 360
ggaattggat tcagctggag agctgcctct ggggccagag cgttctctgg ctgcacatca 420
ggaattaaat gacagagcga ctccagaaga aaggctccca gagggtggtc cgtctctcaa 480
ccggatccag aaccaggccc tagttcagct tcttcgcttc aagtgggttc aaactacaga 540
tcagatcggg gctgcaggtc atcaggcaga tgaataactt gtccactag acagtaaggt 600
ttcaagacca accaaatttg ttgtttcgcc caagacctg aagaagatc tagctgaacg 660
ttggagcctt cctgagattg ttgggatctc acaccaatha tccaaactc agcgtcagaa 720
acagactttg ccagatgakt acttgagtat ggaacactg tatccgggca gctactctc 780
agactccggg gtgaacgcag atgaactcc agggcctctc gacaggttg gactttctca 840

```

```

actccatctc  gagcccaaaa  gtcaasatcc  agagaccctt  gaagacatcc  aetccctctc  900
actccaggaa  gaagcccccag  cgcagctttt  acagctccct  caggaggtag  aactttccac  960
ccagcaggag  gccccagctc  tgcctccaga  gkctctctatg  gagagtcctag  ctcaaaatcc  1020
actgaatcat  gaagtgaag  tccascctcc  aggtgaggat  caagctcatt  ataatttgcc  1080
caagltttca  gtcaaacctg  cagatgtgga  ggttaccatg  acttcagagc  ctasaaatga  1140
gacagaatct  attcaagccc  agccggaggc  cccattccag  cctcccgagg  agggcggaac  1200
ttcttctaca  gccctgaggc  ctacagatcc  tcttccagaa  caccctgagg  tgacacttcc  1260
accttcagac  aagggtcagg  ctacagcttc  acactgact  gaagccacac  tccacctct  1320
ggacctggag  cttagcataa  ctacagagcc  tcttccagag  gttaaacctg  ctccaaacct  1380
ggaggaaacc  tccagtcagg  ctccagacc  ggggtttccc  ataactccag  aacccactac  1440
agagatttga  cattccacac  cctggagaa  gactagagct  cctcatccag  accaggttca  1500
gectctgcat  cgaagctga  ctgaagtcac  aggtccacct  acaaagttag  aatcttcgca  1560
ggatctcattg  gtgcagctc  aacctgcacc  agaggaaacag  aaggctccca  caagcaccac  1620
catatgttag  ctctgcacct  gccggagatga  gactctgtcc  tctgttggtc  tcagcccaaa  1680
gcagaggctc  cgcacagtc  cctgtccaga  gcccgacacc  tccaatggca  tcttcacacc  1740
cttaaatctc  caaggaaact  atatttccat  ccttgatgga  aatgtatgga  aagcatacag  1800
ctggaccgag  aaactaatc  tcagtgaacc  ttatttgact  gaattaccta  aggattcatt  1860
tgaaggcctg  ctataacctc  agtatttga  ttatctctgc  aataaaatac  gatataatga  1920
aagcccaaca  ttgcaatcac  taccatttct  gcagtatata  aatctgggct  gcaatttcat  1980
tacaasactg  agccttgga  catttcaggc  ctggcacgga  atgcagtttt  tacacaactt  2040
aacttctaat  cgcacactcc  tgactactgt  cgaagatcca  tatctctttg  aactgcgggc  2100
attaaaatac  ctagacatgg  gacaaacaca  catcacactt  acacacetta  agaactttct  2160
ccgctgact  gttgaactgg  aaaaactgat  ctacactagc  catatggcct  actgcctctg  2220
ccactttaaa  aatagcattg  agctgtctg  caagacagtc  aagctccatt  gcaacacctc  2280
atgtctgact  aacagcatac  attgtccctg  aagacatct  gtagggatcc  cagaaggagc  2340
gttcatgaag  atgttacaa  cccggaagca  gcacatgagc  actcagctga  ctatttgaat  2400
ggagggccac  tcagacagcc  gtggcatcaa  ctctcaggc  tttgggggtg  attagcttga  2460
aattcagcta  accgagcagc  tacggtccct  cttccccaac  gaggatgtga  gaaagtccat  2520
gtctcatgtt  atccggacct  cgaacatgga  atgttcagaa  acacatgtgc  aaggagctg  2580
tgccaaagctc  atgtcgcgaa  caggctccct  gatgaagctt  ctacagcagc  agcaggaagc  2640
aaaggtcattg  aatgtagaat  gggatagga  ccaacaaaaa  acacattata  ttaatgagaa  2700
catggacacg  aatgaacaga  aagagcagaa  gtcnagtyag  ctcatgaag  aagttccagg  2760
agatgactat  aagaacaaac  tcatcttcgc  aatatctgtg  actctaatac  taataatttt  2820
gattataatt  ttttgtctta  tagaggtgaa  ctacataaaa  agggcatcag  aaaaatacaa  2880
agacaaacca  tcaatatcag  ggcctgagc  atgagttaaa  gcattgtggat  ggcttggagc  2940
tatgttttta  aaaktgttat  taatatctgg  ttttttactt  aaaaaaadaa  aaaaaaadaa  3000
aaaa 3004

```

&lt;210&gt; 108

&lt;211&gt; 959

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 108

```

Met Ala Phe Ala Glu Cys Ile Ala Pro Ala Cys Val Met Ser Trp Leu
  1             5             10             15
Arg Phe Trp Gly Pro Trp Pro Leu Leu Thr Trp Gln Leu Leu Ser Leu
          20             25             30
Leu Val Lys Glu Ala Gln Pro Leu Val Trp Val Lys Asp Pro Leu Gln
      35             40             45
Leu Thr Ser Asn Pro Leu Gly Pro Pro Glu Pro Trp Ser Ser Arg Ser
  50             55             60
Ser His Leu Pro Trp Glu Ser Pro His Ala Pro Ala Pro Pro Ala Ala
  65             70             75             80
Pro Gly Asp Phe Asp Tyr Leu Gly Pro Ser Ala Ser Ser Gln Met Ser

```

85										90					95				
Ala	Leu	Pro	Gln	Glu	Pro	Thr	Glu	Asn	Leu	Ala	Pro	Phe	Leu	Lys	Glu				
			100						105						110				
Leu	Asp	Ser	Ala	Gly	Glu	Leu	Pro	Leu	Gly	Pro	Glu	Pro	Phe	Leu	Ala				
		115						120					125						
Ala	His	Gln	Asp	Leu	Asn	Asp	Lys	Arg	Thr	Pro	Glu	Glu	Arg	Leu	Pro				
		130					135						140						
Glu	Val	Val	Pro	Leu	Leu	Asn	Arg	Asp	Gln	Asn	Gln	Ala	Leu	Val	Gln				
	145				150				155						160				
Leu	Pro	Arg	Leu	Lys	Trp	Val	Gln	Thr	Thr	Asp	Leu	Asp	Arg	Ala	Ala				
			165					170						175					
Gly	His	Gln	Ala	Asp	Glu	Ile	Leu	Val	Pro	Leu	Asp	Ser	Lys	Val	Ser				
		180						185						190					
Arg	Pro	Thr	Lys	Phe	Val	Val	Ser	Pro	Lys	Asn	Leu	Lys	Lys	Asp	Leu				
		195					200						205						
Ala	Glu	Arg	Trp	Ser	Leu	Pro	Glu	Ile	Val	Gly	Ile	Pro	His	Gln	Leu				
	210					215					220								
Ser	Lys	Pro	Gln	Arg	Gln	Lys	Gln	Thr	Leu	Pro	Asp	Asp	Tyr	Leu	Ser				
	225				230						235				240				
Met	Asp	Thr	Leu	Tyr	Pro	Gly	Ser	Leu	Pro	Pro	Glu	Leu	Arg	Val	Asn				
			245					250						255					
Ala	Asp	Glu	Pro	Pro	Gly	Pro	Pro	Glu	Gln	Val	Gly	Leu	Ser	Gln	Phe				
		260					265							270					
His	Leu	Glu	Pro	Lys	Ser	Gln	Asn	Pro	Glu	Thr	Leu	Glu	Asp	Ile	Gln				
		275					280						285						
Ser	Ser	Ser	Leu	Gln	Glu	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Gln	Leu	Pro				
	290					295							300						
Gln	Glu	Val	Glu	Pro	Ser	Thr	Gln	Gln	Glu	Ala	Pro	Ala	Leu	Pro	Pro				
	305				310				315						320				
Glu	Ser	Ser	Met	Glu	Ser	Leu	Ala	Gln	Thr	Pro	Leu	Asn	His	Glu	Val				
			325					330						335					
Thr	Val	Gln	Pro	Pro	Gly	Glu	Asp	Gln	Ala	His	Tyr	Asn	Leu	Pro	Lys				
		340					345							350					
Phe	Thr	Val	Lys	Pro	Ala	Asp	Val	Glu	Val	Thr	Met	Thr	Ser	Glu	Pro				
		355				360							365						
Lys	Asn	Glu	Thr	Glu	Ser	Thr	Gln	Ala	Gln	Gln	Glu	Ala	Pro	Ile	Gln				
	370					375							380						
Pro	Pro	Gln	Glu	Ala	Glu	Pro	Ser	Ser	Thr	Ala	Leu	Arg	Thr	Thr	Asp				
	385				390				395						400				
Pro	Pro	Pro	Glu	His	Pro	Glu	Val	Thr	Leu	Pro	Pro	Ser	Asp	Lys	Gly				



	405		410		415
Gln Ala Gln His Ser His Leu Thr Glu Ala Thr Val Gln Pro Leu Asp	420		425		430
Leu Glu Leu Ser Ile Thr Thr Glu Pro Thr Thr Glu Val Lys Pro Ser	435		440		445
Pro Thr Thr Glu Glu Thr Ser Ala Gln Pro Pro Asp Pro Gly Leu Ala	450		455		460
Ile Thr Pro Glu Pro Thr Thr Glu Ile Gly His Ser Thr Ala Leu Glu	465		470		475
Lys Thr Arg Ala Pro His Pro Asp Gln Val Gln Thr Leu His Arg Ser	485		490		495
Leu Thr Glu Val Thr Gly Pro Pro Thr Lys Leu Glu Ser Ser Gln Asp	500		505		510
Ser Leu Val Gln Ser Glu Thr Ala Pro Glu Glu Gln Lys Ala Ser Thr	515		520		525
Ser Thr Asn Ile Cys Glu Leu Cys Thr Cys Gly Asp Glu Thr Leu Ser	530		535		540
Cys Val Gly Leu Ser Pro Lys Gln Arg Leu Arg Gln Val Pro Val Pro	545		550		555
Glu Pro Asp Thr Tyr Asn Gly Ile Phe Thr Thr Leu Asn Phe Gln Gly	565		570		575
Asn Tyr Ile Ser Tyr Leu Asp Gly Asn Val Trp Lys Ala Tyr Ser Trp	580		585		590
Thr Glu Lys Leu Ile Leu Ser Glu Asn Tyr Leu Thr Glu Leu Pro Lys	595		600		605
Asp Ser Phe Glu Gly Leu Leu Tyr Leu Gln Tyr Leu Asp Leu Ser Cys	610		615		620
Asn Lys Ile Arg Tyr Ile Glu Arg Gln Thr Phe Glu Ser Leu Pro Phe	625		630		635
Leu Gln Tyr Ile Asn Leu Gly Cys Asn Leu Ile Thr Lys Leu Ser Leu	645		650		655
Gly Thr Phe Gln Ala Trp His Gly Met Gln Phe Leu His Asn Leu Ile	660		665		670
Leu Asn Arg Asn Pro Leu Thr Thr Val Glu Asp Pro Tyr Leu Phe Glu	675		680		685
Leu Pro Ala Leu Lys Tyr Leu Asp Met Gly Thr Thr His Ile Thr Leu	690		695		700
Thr Thr Leu Lys Asn Ile Leu Thr Met Thr Val Glu Leu Glu Lys Leu	705		710		715
Ile Leu Pro Ser His Met Ala Cys Cys Leu Cys Gln Phe Lys Asn Ser					720

725										730					735				
Ile	Glu	Ala	Val	Cys	Lys	Thr	Val	Lys	Leu	His	Cys	Asn	Thr	Ala	Cys				
740										745					750				
Leu	Thr	Asn	Ser	Ile	His	Cys	Pro	Glu	Glu	Ala	Ser	Val	Gly	Asn	Pro				
755										760					765				
Glu	Gly	Ala	Phe	Met	Lys	Met	Leu	Gln	Ala	Arg	Lys	Gln	His	Met	Ser				
770										775					780				
Thr	Gln	Leu	Thr	Ile	Glu	Ser	Glu	Ala	Pro	Ser	Asp	Ser	Ser	Gly	Ile				
785										790					795				
Asn	Leu	Ser	Gly	Phe	Gly	Gly	Asp	Gln	Leu	Glu	Ile	Gln	Leu	Thr	Glu				
805										810					815				
Gln	Leu	Arg	Ser	Leu	Ile	Pro	Asn	Glu	Asp	Val	Arg	Lys	Phe	Met	Ser				
820										825					830				
His	Val	Ile	Arg	Thr	Leu	Lys	Met	Glu	Cys	Ser	Glu	Thr	His	Val	Gln				
835										840					845				
Gly	Ser	Cys	Ala	Lys	Leu	Met	Ser	Arg	Thr	Gly	Leu	Leu	Met	Lys	Leu				
850										855					860				
Leu	Ser	Glu	Gln	Gln	Glu	Ala	Lys	Ala	Leu	Asn	Val	Glu	Trp	Asp	Thr				
865										870					875				
Asp	Gln	Gln	Lys	Thr	Asn	Tyr	Ile	Asn	Glu	Asn	Met	Glu	Gln	Asn	Glu				
885										890					895				
Gln	Lys	Glu	Gln	Lys	Ser	Ser	Glu	Leu	Met	Lys	Glu	Val	Pro	Gly	Asp				
900										905					910				
Asp	Tyr	Lys	Asn	Lys	Leu	Ile	Phe	Ala	Ile	Ser	Val	Thr	Val	Ile	Leu				
915										920					925				
Ile	Ile	Leu	Ile	Ile	Ile	Phe	Cys	Leu	Ile	Glu	Val	Asn	Ser	His	Lys				
930										935					940				
Arg	Ala	Ser	Glu	Lys	Tyr	Lys	Asp	Asn	Pro	Ser	Ile	Ser	Gly	Ala					
945										950					955				

&lt;210&gt; 109

&lt;211&gt; 1331

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 109

```

gttctttttct tttccatgat atcatttatat ggacagttta ggggtggtctc atggattata 60
accatttggg tatttggttc actaacaatt ttcttaactg ccagagttct tgggtggagaa 120
gttgcataatg gccaaagtcct tggagttata ggatattcat taattccctct cattgtcata 180
gccccgtgtac ttttctggc tggatcattt gaagcgggtgt ctacacttat aaactgttt 240
gggtgtcttct gggtgccta cagtgcctgt ccattgttag tgggtgaaga attcaagacc 300
aaaaagcttc ttctgattta tccaatcttt taattataga tttatatttt gtcgttatat 360
actggtctgt gatccaaagt ctacatgat agaaaaagat ggtgttaaat ttgtgtgtag 420
gatgggaatt ctgctgaag gaattggaga aaactgttg ctgcaaat ttcatgttc 480
cagatggaaa ggggaagtcta agcgtttttt aaacaaatt ttttttgtat ttaattaaag 540

```

```

aattggcagtt atctggggtt ttgggggcag aatttttaatt tctgtttgat tctccatatt 600
cncgtgcaata aactacaaaa gcatttgtgt tttaagattg tctcgacatt caccataaaa 660
cttctgcccc aagcaccctgg attggtaatt atatttccct taaagggtta atttgacaa 720
atcttgataa tcaaaagtgc aatttttctt tcaaaatgt ttctccagc atcacagatc 780
ctgcagatst atatttatat ttatccatat atatttatga aataatctt cctcacaaaa 840
tatattttct ataacatta agatattaaa tctgatgcac aaacttttta atttggccat 900
taattttttt tatttaaaaa cttaaatctt ctthtaaaat tgtatatagt ttttaaaatc 960
tcacacatgc ttcgatacct ccttgcttaag aattcttaat aactactaaa actgattctt 1020
aatagttgct gatatatatt tgggttgttt gggataact ttcaaaacca tttttgaatg 1080
tcaaacatc tgatttaag ttctctgtta tcttctgac caaaggagca agaggatata 1140
tggatatact attcattaaa atctttacta tgtacaaaa cagtaatttt tacagcatcc 1200
gtaatatttt ttaagtggtt ctctaaatc ataaaagtc gggaaagaga cctttaaat 1260
cttctgggtc tgaacaaatg tatatgaag agaaaaata aaatacttc cagttgaaa 1320
aaaaaaaaa a

```

1331

&lt;210&gt; 110

&lt;211&gt; 118

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

```

Met Ile Ser Leu Tyr Gly Gln Phe Arg Val Val Ser Trp Ile Ile Thr
  1             5             10             15

```

```

Ile Trp Ile Phe Gly Ser Leu Thr Ile Phe Leu Leu Ala Arg Val Leu
          20             25             30

```

```

Gly Gly Glu Val Ala Tyr Gly Gln Val Leu Gly Val Ile Gly Tyr Ser
          35             40             45

```

```

Leu Leu Pro Leu Ile Val Ile Ala Pro Val Leu Leu Val Val Gly Ser
          50             55             60

```

```

Phe Glu Val Val Ser Thr Leu Ile Lys Leu Phe Gly Val Phe Trp Ala
          65             70             75             80

```

```

Ala Tyr Ser Ala Ala Ser Leu Leu Val Gly Glu Glu Phe Lys Thr Lys
          85             90             95

```

```

Lys Pro Leu Leu Ile Tyr Pro Ile Phe Leu Leu Tyr Ile Tyr Phe Leu
          100            105            110

```

```

Ser Leu Tyr Thr Gly Val
          115

```

&lt;210&gt; 111

&lt;211&gt; 2610

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 111

```

aattgacct ctctgtgtta ttctgattgt gtaccttgag acatacagag gattctcttt 60
tctctttttt gcttaagaat gtataattag tgtcttttta ttgggggagaa attttatttt 120
tttgtcattt ctctgaagtt gacatttgat gagtgtatc tgaattctac cactcctctg 180
gggaaaaaaa tctcaatgga agattgtaca gttaagtag ttgttttttc ctcttggagt 240
atgtattttt taaggtcatt aatttaaaat agaaataatt ctttaacca ttggggtaaa 300
atttttcaaa atatgtgtgt ttgatataat ttgagaaaa gtgcctgaat cctaaacta 360
actgtctata aggttaagtc cgtataacaa atgattatat atataacaaa gtadaagtaa 420
atagtgtaga atattaaagt tctcttccag ggcattgggt attgtggggg attaaacaga 480

```

```

agccccccttg ctatgatcaa ctgcacagga aagaacctaa atctgggttca gtgataaaca 540
aaaatagcaa agactgatca ggaanagag aagagactat gcaaatccag atgatccaca 600
ggcagagaga ttttaagtgc taatagcttt actccctgat attccctcag tagaatgggc 660
tttttctttt agtctaagtg accttttgaa ggtcttttga ttcattacgc ttttatatgc 720
tgcctttctt tctctttctt tttttttcat ccaaccacgt aagaastgae gaagtttatt 780
gatataattg ccaataaatt accaatctgt tgcacacaca tgcataatgc ttattagcat 840
ctctgttagt totaaaatte acttcaccca tagaacatta tcccagacct tcaaatggaa 900
agctctcttt atgcaagaga aaaaaggtat taaaagtga ccaattttag atgagggaat 960
tggttatgtt gaatttctgt tcaaaaatca ctaggteaaa ttcccatatc taataatgta 1020
tttttaggac aagcaagggt tbgtcattaa gacatagtaa cattaaacta atatttaaaa 1080
atagatttag ctatatattt tcatgagaaa aggtttctgt tggccaggac aaaaaccatc 1140
ttgatcctaa agagacatta gcagtgcaca tgtcagaqcc ttctcacatt ctgggaata 1200
agtaaacacc tccagagaaq gtgagggcac caaatggatc attcttcaaa atgaaagga 1260
gtctcttact ttctcttgag acatttcttc cctatttggg gatctttatg ctgctgttct 1320
ggagcgccca catgtcttgc ttccatgcag gccaaatcct tctcttgggt ttgggcaga 1380
gaagttaatt caccatacag catcttgatt agacagaact atcatctgat ctatccagg 1440
gcaaggaag tggcatgccc aaaggttctc ctatgtcttc ttcttttaa tgatttcta 1500
attttcagaa gggctctgga tgaactagtt tctgtaagg agattttgct gtggtgtctg 1560
ttctccagg tttaaggcac acaaaacctt ctgtggtcca ctgtgcttg tcttgtaat 1620
tcactgaacc atggcagaa atgggttaagg gaaastgaat tttgacttta catagttaa 1680
tgagtaggag ctacagcaaa aaaaaaaaaa agaaccttaa ccttttattt aatagtatto 1740
ttcacttcag gaatttcaga ctttagttg ttccaggtcag aaaaatgtga tatatcctgt 1800
aagactgtct tcaactcaat caagttaaat gttaacagcaa atactttcag ctctacttcc 1860
aaaagcttcc cagaacttcc caccttccat caotltaact tctaccttc tggtaaaaat 1920
taaccaagatt tttagtagtt cactgttccc tggttctct gatttgatcc ttgtacct 1980
ctgttttaqt cctcaagaa aaaaactcaa cattttaa cgtttcact ctactaatg 2040
gttctcactc cagaagaaa acaaacgaaa tatettatgt cctctcttt ttctgaata catcaacaa 2160
gaactacttt cctctcttt aatgaacaaa gatggcagga ctlttgaaat ctlttgaaat ccttctct 2220
ttatctctct cttaggtgac ttctcttata caccatgtg cagagggtca ggaagaaaa caccactgc 2340
tagtctttgt ttatataat attattatca ccacttact tgttttgtt tctctgtag 2400
tgcgccccac actcatccac ctgacctgta caacttact taatgtgtac attctctca 2460
actcatgac tttttgaaat ataattttt taatgtgtac attctctca 2460
ttcatatgta aggtctgaa gacacagact ggttttttg ttaactgttg atgctcagt 2520
ccttaaaaaa cgcatagcac gaattgcac tttttaaatt aataaatctg gaaattgtt 2580
aaaattcaaa aaaaaaaaaa aaaaaaaana

```

<210> 112  
 <211> 116  
 <222> PRT  
 <213> Homo sapiens

```

<400> 112
Met Ala Gly Leu Leu Asn Val Thr Phe Ile Tyr Leu Leu Leu Glu Cys
  1             5             10             15

Leu Ser Leu Tyr Thr His Val Thr Cys Ser Ser Leu Pro Ser Ser Leu
  20             25             30

Cys Leu Tyr Ile Tyr Tyr Tyr His Arg Gly Leu Gly Lys Lys Thr Pro
  35             40             45

Thr Ala Ala Pro His Thr His Pro Pro Ala Leu Tyr His Leu Leu Cys
  50             55             60

Phe Val Phe Leu Cys Arg Ile His Asp Phe Leu Lys Tyr Asn Phe Phe
  65             70             75             80

Asn Val Tyr Ile Leu Tyr Ala Phe Ser His Ser Tyr Val Lys Ser Gly
  85             90             95

```

Arg His Arg Leu Val Phe Leu Phe Thr Val Asp Ala Ser Val Pro Lys  
100 105 110

Ile Cys Ile Ala  
115

<210> 113  
<211> 2759  
<212> DNA  
<213> Homo sapiens

<400> 113  
tttttttttt tttgaaagac acaagtttatt ttattaataa agcaatctct cccactctgc 60  
ccagtgtgtga aggtgttttg cctgtccttgc ggaaggggac caaatgtctt gggggcccta 120  
gcccgtctgc acaggtctagg cctgtccttgc gccaagaaag ctgctcaaac tctagatgac 180  
atttggaagg atgaggacct gagccagag gtggcagtg cctaccacag gaagtcacaa 240  
gctgtgtctc caggtccacg ctctgggttg ggcaggact aaatcctggc tcccccttct 300  
tggtaactaag gggatttagt cttgtgtgtc tgtaggggtt cagagtaggg aggtttccag 360  
gaagggttcc agagtgggtt cacaagggac ctctccctt ggcctcttgg agtccaggtc 420  
gtcgaagggg caaagcttga cgcctcctg gccaagctgg gccctcagcg tgggcccgtt 480  
gaggaagcgc agtctatgca gcccctccca agagcaagg aaaggtctgg agccttcacc 540  
gcagccgcgc gtgggaggca cactgggtga gcccgggttg gccatcagct cggctgtcag 600  
gggtgtggccc gctagggtac ctccagggac ccgcgccagg gccccggaca cgcggtgagc 660  
ggacatgttg cggccgcaag tctccaggcc caccgaaggg tctgtctccc gcaggccctg 720  
pggggagaa ggcctcaggg ggcgggttg tgcgctcca cggccaggg gaaggccagg 780  
pggggggccc ccagccaagt gcagccaccc acactcggct ccagcggcag tgcgtaaaag 840  
cgcaccccat aggccttgcg cgcctcggcg aaacactggc acacgccttg gacacagtg 900  
acgtgtctgt gccggtccgc gtgtgtgggg gccctgcaca gcagctcccg gaagcagctt 960  
agttgggcct cagcttctc ccgacctag cgcgggagcg agggcagcg atgctccag tgtttgaatg 1020  
ggagtgtctg ccagggtat cctctctctg ctccctctgt cctccgggg cgggtctctg 1140  
cggaaagtcac ccacggcccg ctcttaacgg ctccacagta cctccgggg cgggtctctg 1200  
gggtccctgc gacatccctc ctgtagcttg ggcctccgac ctgaggcaaa tccacgtctc 1260  
cggccgcac ccgctcccg aatcccatct tgcacaggaa gaagccttcc gggccagcca 1320  
ggatcaggg gcaacggggg gcgggaccca cggggcggcc ctgggacagg tctctcctg 1380  
gdcctgtggg gatgtgttcc ctgggggcca gctccgcgc gctctcctg gccgcacgt 1440  
tgacagcag ggacacgtg gtacaggtcc cggccagaaa ggctccacag ataccctcat 1500  
cgccttcggg caggtacacg aggtgtccg cgggtccccc caggcggccg cgggagccgg 1560  
cagatccggg tctgtgtggt aagagtgcg tggctactcg aatcaaaaac caggcagtcg 1620  
agggaagcgg ccggcagaaa cggcagtgcc agcaggttcc ggagcagtcg cagccttctg 1680  
gaagctccag ccggtcttct tgcagagct cggctccggc ccccatctc cccgcccct 1740  
cggltgttgt ctgggcggat ttcaacagtc aagtaaaatc aagctgggta atcatgccc 1800  
aaggtggatt tcatccctgt gaatgtgttt gctctcctga acatgcaatg agaagcctga 1860  
tcaatctgtt acggcagktc cagtcctct gtacagacac agagtgtctt caggaaattc 1920  
cgggacccct tgggtgataat ggcacagtg ttacaatgat cttggtagcc tggatggtta 1980  
ttgcatgtat ctgtttctta ctgagacctc cttaactaag aggtaccagc ctacttgga 2040  
agccacccag tctcataaat ggacaagatc caccagttcc tctgttgcc taactttgtg 2100  
atatgggaag tgaactagt taacacctt cagcaccaaa cgaacganga tgaccagagt 2160  
actcttaacc ccattagaac tgttttctct tttgtabctg caatatggga tggatattgt 2220  
tctatgagct tctagaactt cactttgcaa gtttattttt gcttctgtg ttaactgcat 2280  
tccatcttcc agtatatttg agtgaatgat tatattttta aaagtttaca tggggctttt 2340  
ttggtttgtc taacttaca aacattccca tcaattctgt tgttaactgt attatanatt 2400  
ttgtgtatct ttctggctg attgaaggaa atttggagg cctgcattta tatattttta 2460  
atagatttga taggttttta aaktgtcttt ttccatcagg tatttatatt taccttggt 2520  
gttgtctggg attgtgcgaa agaaaattag aaccacgtg tatttacatt ttggatgct 2580  
gttlatattgt ggcgtggcag tctctgtagt ttgggggact gtggtagctc ttggattgt 2640  
ttgcnaatta cagctgaatc ctgtgtcatg gattaaactg gcttatgttg ctgaatagg 2700  
aagapaaaaa aaatgaattg gttgtttact aaktttatcc tccattaaa aactctcaat 2759  
gttaagaaaa ccttaaatna acatgattga tcaatatgaa aaaaaaaa aaaaaaaa

<210> 114  
 <211> 99  
 <212> PRT  
 <213> Homo sapiens

<400> 114  
 Met Ala Glu Gly Gly Phe Asp Pro Cys Glu Cys Val Cys Ser His Glu  
 1 5 10 15  
 His Ala Met Arg Arg Leu Ile Asn Leu Leu Arg Gln Ser Gln Ser Tyr  
 20 25 30  
 Cys Thr Asp Thr Glu Cys Leu Gln Gly Leu Pro Gly Pro Ser Gly Asp  
 35 40 45  
 Asn Gly Ile Ser Val Thr Met Ile Leu Val Ala Trp Met Val Ile Ala  
 50 55 60  
 Leu Ile Leu Phe Leu Leu Arg Pro Pro Asn Leu Arg Gly Ser Ser Leu  
 65 70 75 80  
 Pro Gly Lys Pro Thr Ser Pro His Asn Gly Gln Asp Pro Pro Ala Pro  
 85 90 95  
 Pro Val Asp

<210> 115  
 <211> 1404  
 <212> DNA  
 <213> Homo sapiens

<400> 115  
 aatcgggacg ggaagaatta ttggttgggg gaaccccaag aggggacgcy gccgaggagg 60  
 gtcgtgtgac acccgggggc gtgggagtg ggtaccagat tcagcccaatt tggcccccag 120  
 gctctgtgtt tcggaatecg ggtgctgcgg attgagggtc cggttccctaa cgggtgggac 180  
 ggtgtctctg ggtgagatt tggcgttttc tggggccttt ggtgggacg gtgtctctca 240  
 gatgagattt aggttttctt cggggctttt gggatcttca cctaataacc ggtattattt 300  
 tatgagagga gtgtgtttgg ctgtcagAAC tggatccctg ggtgatatt tgggaattag 360  
 tggagtgtac tctgaagacc tagggctatg ctctggagct gctgtggctg aattttgggg 420  
 cctctgaagt ggcattggaga ttgaggtcca gagagccctg gatcttgagg gctgacattt 480  
 ggagagatgg ggtcgagggt tctcttttgg ccttgactgc ttggggcctt tctcactctc 540  
 attccggga tcttttgcA gaattctctg cggattggcc gtaaccctgt ccccgagcgg 600  
 gctcacaggg tctgagggc acgcatgagg caaaggtaaa gttctgagcc acccggtgac 660  
 tctttccag gactccaaga tggagggaag cgggaacctc ggaggcctga ttaagatgg 720  
 ccactacttg gtcttgctag gtgcctgggg catgcacatg tgggtgacct tctctccagg 780  
 tagggacctt cagcttgggt gtcattgggt cctgggggtg ggtatggaaat aagaggggaa 840  
 ccgggaagt ccttaacacc cctgtgggtc cctaccctg caggcttctt gcttttccga 900  
 agccttcccc gacataacct cggactagt catcaacctc tgcattcttg cttaacagca tgcctgggct 1020  
 ctctccatgg gctgtgctt catcaacctc tgcattcttg cttaacagca tgcctgggct 1080  
 cagctcacat tctggaggc cagccagctt taactgctgt tcttgagcct tccgtgggac 1140  
 actgtcaacg cccgtctgct ggaacccccc accacagctg ccatgtgggc cctgcacacc 1200  
 gtggggaagg agcgaggcct ggttggggag gtaccaggca cctaccaggg tccgatccc 1260  
 taccgcccgc tggagagaa ggaaccccaag tacagtgtct tccgccagaa tctctccg 1320  
 taccatgggc tgtctctctt ttgcaatctg ggtgctgtcc tggagcaatg gctctgtctc 1380  
 gctggccttg cctgggaat aaggagcctc tagcctgggc cctgcctgct aataaatgct 1404  
 tcttcagaaa aaaaaaaaaa aaaa

<210> 116  
 <211> 180  
 <212> PRP  
 <213> Homo sapiens

<400> 116

```

Met Ser Trp Val Pro Gly Val Gly Met Glu Ile Arg Gly Glu Pro Gly
 1             5             10             15

Ser Ala Leu Thr Pro Leu Trp Ser Pro Tyr Pro Ala Gly Phe Leu Leu
      20             25             30

Phe Arg Ser Leu Pro Arg His Thr Phe Gly Leu Val Gln Ser Lys Leu
      35             40             45

Phe Pro Phe Tyr Phe His Ile Ser Met Gly Cys Ala Phe Ile Asn Leu
      50             55             60

Cys Ile Leu Ala Ser Gln His Ala Trp Ala Gln Leu Thr Phe Trp Glu
      65             70             75             80

Ala Ser Gln Leu Tyr Leu Leu Phe Leu Ser Leu Thr Leu Ala Thr Val
      85             90             95

Asn Ala Arg Trp Leu Glu Pro Arg Thr Thr Ala Ala Met Trp Ala Leu
     100             105             110

Gln Thr Val Glu Lys Glu Arg Gly Leu Gly Gly Glu Val Pro Gly Ser
     115             120             125

His Gln Gly Pro Asp Pro Tyr Arg Gln Leu Arg Glu Lys Asp Pro Lys
     130             135             140

Tyr Ser Ala Leu Arg Gln Asn Phe Phe Arg Tyr His Gly Leu Ser Ser
     145             150             155             160

Leu Cys Asn Leu Gly Cys Val Leu Ser Asn Gly Leu Cys Leu Ala Gly
     165             170             175

Leu Ala Leu Glu Ile Arg Ser Leu
     180
  
```

<210> 117  
 <211> 1801  
 <212> DNA  
 <213> Homo sapiens

<400> 117

```

tgaagaaggt gtttactttt tttagaantta ccttgagaca ttcaaaactg tgcagaagat 60
atatgcacaa sagcaaatgt cttgcagttt gctatagcca cttataacat atctggctct 120
tgaatagctt taattcagct gttgaatctc acttgaantt gagcaaaacc ttcattcttta 180
tatgtatctg gacaaattac ttcaattgct tgacagtaat gaccaatcaa tttattttaa 240
atagtatcat ttagtaggac agtgttttct cctgttctga gcaacgaatt caaccagctc 300
cttgggttga tcatcatcat catctccttt tgggtatcag ttcttgagtt atttttacca 360
gggagtttta tactttttag caactatttt gaattatctc aggaatgtca tatactctctg 420
cctcttttag gtcagtcact ggcactctgt ctgkktggtg acatcatgtt tccctgactg 480
ttcttcactc ttgtagttac acattgatat ctgtgcattg aatatgtagg taattataaa 540
cagtcctttg aatctggctt tgtctgtgat tgtccttgca tagtgggtct gtccagaaat 600
tgtaagcata ctgtcttttt tggctcttaa gccctgacac gctacagccc gtgtagtgc 660
  
```

```

aatgggtgccc ctaagcccccag gtcccccctgca gtccactctcgc tgcattctgttt gttgactggt 720
gtgagccccc ccccccattctc ttgtttcttaa ttaccaccta gcaggctaac cctgctggca 780
cctgcagtgc ttccaggggg aagggaaccak agtgtgcccc tctgaagagt ctcagaatgg 840
tgcggaaaggt gaatgcccac ctcccgctct ctcttccac tctagaaact gattccagag 900
aaattctcca agtgcgggtgc tatgtgggct tgcgggagag gtgttatgat caaacagaac 960
cattctctctt accctctgtt catgggtttt ctgggtctcg tggccagtg agttgtcaca 1020
gcttcaactc caatctctgg gatattccagg gcantaatct tgcactcggg tatttgctag 1080
ttgaaattat gtggtaggga gagaagccag taagcttcc tctcccgttt ggttgatgta 1140
actctcgaat cctgtacttt catacggatt gtaataggga ggtctcaca ggaagcttca 1200
gttttggatt tttagagctt ctctcaagta caattgttga atcagaggt aagcatggtc 1260
tggtataact ggaataccgc tgagattaaa aaggagataa attagctcca ttgggacagt 1320
gtatttggga atgtgaattg ataccacttt cctggagctc agtttagtat ttaktttata 1380
tgtaabgcct gaaagatgt gtgtctctt tgactcaata ctataggga ttacacaca 1440
ggttataact aagattata gaactgttat actgcaaaa cagcaatat tctattatg 1500
aattaatgg ggaattggtt aataaactat ggtttgatat gctctaaaa taatgttaca 1560
gaaaaagtg tactgatat gcaaatgta tgacttateg ttaaaaaagg aggttagatg 1620
ttgatagata cagtatgata gaaaaagatc eggaaggtat atctgacat ttaaatctgg 1680
atttttatga gtgtttttt cattccaatc ttctacatg catgtatttt ctageaattg 1740
tattctcttc ttgtaataa actcaattat ttttaaggga ctaaaaaaa aaaaaaaa 1800
a

```

&lt;210&gt; 119

&lt;211&gt; 66

&lt;212&gt; FR7

&lt;213&gt; Homo sapiens

&lt;400&gt; 118

```

Met Val Arg Lys Val Asn Ala His Leu Pro Leu Ser Phe Pro Thr Val
  1             5             10             15

Glu Thr Asp Ser Arg Glu Ile Leu Gln Val Arg Cys Tyr Val Gly Leu
          20             25             30

Arg Glu Arg Cys Tyr Asp Gln Thr Glu Pro Phe Ser Leu Pro Ser Val
          35             40             45

His Gly Phe Ser Trp Leu Cys Gly Pro Val Ser Cys His Ser Phe Thr
          50             55             60

Pro Asn Phe Trp Asp Ile Gln Gly Asn Asn Leu Ala Thr Gly Tyr Leu
          65             70             75             80

Leu Val Glu Ile Met Trp
          85

```

&lt;210&gt; 119

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (2)

&lt;223&gt; biotinylated phosphoramidite residue

&lt;400&gt; 119



anagggagcc tctgaccat ctctcttc

29

<210> 120

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 120

cncacagaaa attcaataag accctcgt

29

<210> 121

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 121

cncagctctt cgtaggaag ttctgactt

29

<210> 122

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 122

antctgcac accagccagt aacgcaccc

29

<210> 123

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 123  
 ggggtgggag agatgtgtgg ggaacaaga 29

<210> 124  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 124  
 aatgggtgtt agccacaca acagggtga 29

<210> 125  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 125  
 aacggcaggg aactacagg gacagagct 29

<210> 126  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 126  
 gggttttcgg tgtcgatggt gtagaggat 29

<210> 127  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 127

cnagsacaca tagggatgag agagcaagc

29

<210> 128

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 128

anactgaaa ctgagtatgt gagagtgtg

29

<210> 129

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 129

gmlatccatt tctcttctct catctgagt

29

<210> 130

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 130

tncttgagga atgggttgaa gtccggcgg

29

<210> 131

<211> 29

<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 131  
 tctctgctg tgcctctc tctcgac 29  
  
 <210> 132  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 132  
 ggcctctca ctggatgtca tctcatca 29  
  
 <210> 133  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 133  
 tngtcctgt gaaggcata gccagttg 29  
  
 <210> 134  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 134

gngcactgta ttgagctgat tgetgaagc

29

<210> 135

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 135

tnccagagca gaagcatgac aggcacac

29

<210> 136

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 136

anacattctg agtagttgca tgatttccc

29

<210> 137

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 137

gncagaaag ttgaggacat gctgggcag

29

<210> 138

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 138  
 atggggaacaa gacaactgga gaagggtca 29  
  
 <210> 139  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 139  
 tngtttccca ggtagacaga ggggttcag 29  
  
 <210> 140  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 140  
 atccatctac atgtgcattg acaagctta 29  
  
 <210> 141  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 141  
 tngtgataga tcctttcgta acacaaagt 29  
  
 <210> 142  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>

<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> {2}  
<223> biotinylated phosphoramidite residue

<400> 142  
angaccagat ctcccccagc acatcacc 29

<210> 143  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> {2}  
<223> biotinylated phosphoramidite residue

<400> 143  
tntctggggc aagatggctg ttaagcagt 29

<210> 144  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> {2}  
<223> biotinylated phosphoramidite residue

<400> 144  
tnggttggttc cgggcagggc attcttgtc 29

<210> 145  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> {2}  
<223> biotinylated phosphoramidite residue

<400> 145  
tnacacgctc tgtgctagga cttttatat 29

<210> 146  
<211> 29

<212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> {2}  
 <223> biotinylated phosphoramidite residue

<400> 146  
 cggcagctgtgt gatattcgag attgctggt

29

<210> 147  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> {2}  
 <223> biotinylated phosphoramidite residue

<400> 147  
 aacatcaagtg gtgcctaat aagtctcca

29

<210> 148  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> {2}  
 <223> biotinylated phosphoramidite residue

<400> 148  
 aatcaacttgc atttgtcttg gaacctgac

29

<210> 149  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> {2}  
 <223> biotinylated phosphoramidite residue

<400> 149



gntgacttca atctctctcag ctctcaccg

29

&lt;210&gt; 150

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; {2}

&lt;223&gt; biotinylated phosphoramidite residue

&lt;400&gt; 150

gnagtgtccac ctatgactac caatttctc

29

&lt;210&gt; 151

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; {2}

&lt;223&gt; biotinylated phosphoramidite residue

&lt;400&gt; 151

gnaggatgag gcaatgcaca caatgaaag

29

&lt;210&gt; 152

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; {2}

&lt;223&gt; biotinylated phosphoramidite residue

&lt;400&gt; 152

cnaaactgggt gttttaccg tatccttca

29

&lt;210&gt; 153

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

<222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 153  
 tngattctgc cgaatacaga agtgccttc

29

<210> 154  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 154  
 angttgatgg gctcaacaca ggcacaggg

29

<210> 155  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 155  
 anggatgcca tctctacacc actctgtac

29

<210> 156  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 156  
 anacacccac ctgacacggc attacttaa

29

<210> 157  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> {2}

<223> biotinylated phosphoramidite residue

<400> 157

cncctgaggg tagaagccg ctcaggttt

29

<210> 158

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> {2}

<223> biotinylated phosphoramidite residue

<400> 158

tngagttagc agagcagcaa gcaaggagg

29

<210> 159

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 159

tttgcattga tagttctctg cagtagcat

29

<210> 160

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> {2}

<223> biotinylated phosphoramidite residue

<400> 160

anttgcatg tcaagagga gccagatga

29

<210> 161

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 161  
 tnaggtttgt gacattgctg agcttcag 29  
  
 <210> 162  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 162  
 cnttcagaca ctacgaacgg gcacaggta 29  
  
 <210> 163  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 163  
 angagaagtt ctgtgcgtgg gtctggtcg 29  
  
 <210> 164  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> oligonucleotide  
  
 <220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue  
  
 <400> 164  
 angctttct gcacaggac ctgttagcg 29  
  
 <210> 165  
 <211> 29  
 <212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 165

gntcatctcc agtaccatct ccataaatg

29

<210> 166

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 166

actgtctttg aatggtattg

20

<210> 167

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 167

angttcacat atgatacaag gctctttcc

29

<210> 168

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<220>

<221> misc\_feature

<222> (2)

<223> biotinylated phosphoramidite residue

<400> 169

cnagtagaca gacacggtag tgggttga

29

<210> 169

<211> 29

<212> DNA

<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 169  
gngaatgcac tatgaagana acaggtcag 29

<210> 170  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 170  
anatacagggt gattaggctc ttccatgca 29

<210> 171  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 171  
tntacattca atgcctttgc ttctgctg 29

<210> 172  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide

<220>  
<221> misc\_feature  
<222> (2)  
<223> biotinylated phosphoramidite residue

<400> 172  
antccatgag accaccctaa actgtccat 29

<210> 173  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 173  
 gngaggaaaa gtgctctgtg ttgatgtat

29

<210> 174  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 174  
 antccacctt ctgcctgat taccagct

29

<210> 175  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

<400> 175  
 anccagatg cagaggtga tgaaggac

29

<210> 176  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide

<220>  
 <221> misc\_feature  
 <222> (2)  
 <223> biotinylated phosphoramidite residue

&lt;400&gt; 176

tntaatgoot gaaaagatgt gtgtctctct

29

&lt;210&gt; 177

&lt;211&gt; 388

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 177

Met	His	Leu	Tyr	Lys	Thr	Asn	Lys	Met	Thr	Ser	Leu	Lys	Glu	Asp	Val
1				5					10					15	
Arg	Arg	Ser	Ala	Met	Leu	Cys	Ile	Leu	Thr	Val	Pro	Ala	Ala	Met	Thr
			20					25					30		
Ser	His	Asp	Leu	Met	Lys	Phe	Val	Ala	Pro	Phe	Asn	Glu	Val	Ile	Glu
		35					40					45			
Gln	Met	Lys	Ile	Ile	Arg	Asp	Ser	Thr	Pro	Asn	Gln	Tyr	Met	Val	Leu
	50					55					60				
Ile	Lys	Phe	Arg	Ala	Gln	Ala	Asp	Ala	Asp	Ser	Phe	Tyr	Met	Thr	Cys
	65				70					75					80
Asn	Gly	Arg	Gln	Phe	Asn	Ser	Ile	Glu	Asp	Asp	Val	Cys	Gln	Leu	Val
				85					90					95	
Tyr	Val	Glu	Arg	Ala	Glu	Val	Leu	Lys	Ser	Glu	Asp	Gly	Ala	Ser	Leu
			100					105					110		
Pro	Val	Met	Asp	Leu	Thr	Glu	Leu	Pro	Lys	Cys	Thr	Val	Cys	Leu	Glu
		115					120					125			
Arg	Met	Asp	Glu	Ser	Val	Asn	Gly	Ile	Leu	Thr	Thr	Leu	Cys	Asn	His
	130					135					140				
Ser	Phe	His	Ser	Gln	Cys	Leu	Gln	Arg	Trp	Asp	Asp	Thr	Thr	Cys	Pro
145					150					155					160
Val	Cys	Arg	Tyr	Cys	Gln	Thr	Pro	Glu	Pro	Val	Glu	Glu	Asn	Lys	Cys
				165				170						175	
Phe	Glu	Cys	Gly	Val	Gln	Glu	Asn	Leu	Trp	Ile	Cys	Leu	Ile	Cys	Gly
		180					185					190			
His	Ile	Gly	Cys	Gly	Arg	Tyr	Val	Ser	Arg	His	Ala	Tyr	Lys	His	Phe
	195						200					205			
Glu	Glu	Thr	Gln	His	Thr	Tyr	Ala	Met	Gln	Leu	Thr	Asn	His	Arg	Val
	210					215					220				
Trp	Asp	Tyr	Ala	Gly	Asp	Asn	Tyr	Val	His	Arg	Leu	Val	Ala	Ser	Lys
225					230					235					240
Thr	Asp	Gly	Lys	Ile	Val	Gln	Tyr	Glu	Cys	Glu	Gly	Asp	Thr	Cys	Gln
			245					250					255		
Glu	Glu	Lys	Ile	Asp	Ala	Leu	Gln	Leu	Glu	Tyr	Ser	Tyr	Leu	Leu	Thr
		260					265						270		



Ser Gln Leu Glu Ser Gln Arg Ile Tyr Trp Glu Asn Lys Ile Val Arg  
 275 280 285  
 Ile Glu Lys Asp Thr Ala Glu Glu Ile Asn Asn Met Lys Thr Lys Phe  
 290 295 300  
 Lys Glu Thr Ile Glu Lys Cys Asp Asn Leu Glu His Lys Leu Asn Asp  
 305 310 315 320  
 Leu Leu Lys Glu Lys Gln Ser Val Glu Arg Lys Cys Thr Gln Leu Asn  
 325 330 335  
 Thr Lys Val Ala Lys Leu Thr Asn Glu Leu Lys Glu Glu Gln Glu Met  
 340 345 350  
 Asn Lys Cys Leu Arg Ala Asn Gln Val Leu Leu Gln Asn Lys Leu Lys  
 355 360 365  
 Glu Glu Glu Arg Val Leu Lys Glu Thr Cys Asp Gln Lys Asp Leu Gln  
 370 375 380  
 Ile Thr Glu Ile  
 385

&lt;210&gt; 175

&lt;211&gt; 171

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 178

Met Met Met Gln Cys Val Ser Arg Met Leu Ala His Pro Leu His Val  
 1 5 10 15  
 Ile Ser Met Arg Cys Met Val Gln Phe Val Gly Arg Glu Ala Lys Tyr  
 20 25 30  
 Ser Gly Val Leu Ser Ser Ile Gly Lys Ile Phe Lys Glu Glu Gly Leu  
 35 40 45  
 Leu Gly Phe Phe Val Gly Leu Ile Pro His Leu Leu Gly Asp Val Val  
 50 55 60  
 Phe Leu Trp Gly Cys Asn Leu Leu Ala His Phe Ile Asn Ala Tyr Leu  
 65 70 75 80  
 Val Asp Asp Ser Phe Ser Gln Ala Leu Ala Ile Arg Ser Tyr Thr Lys  
 85 90 95  
 Phe Val Met Gly Ile Ala Val Ser Met Leu Thr Tyr Pro Phe Leu Leu  
 100 105 110  
 Val Gly Asp Leu Met Ala Val Asn Asn Cys Gly Leu Gln Ala Gly Leu  
 115 120 125  
 Pro Pro Tyr Ser Pro Val Phe Lys Ser Trp Ile His Cys Trp Lys Tyr  
 130 135 140  
 Leu Ser Val Gln Gly Gln Leu Phe Arg Gly Ser Ser Leu Leu Phe Arg

```

<400> 100
Met Gly Pro Val Ser Ala Gly Ser Gln Gly Cys Gly Thr Cys Ala Val
  1          5          10          15
Iys Leu Ala Pro Thr Trp Arg Ala Ala Ala Thr Cys Phe Leu Gln
          20          25          30
His Leu Leu Pro Cys Ser Val Ser Ser Leu Ser Pro Arg Leu Ala Gln
          35          40          45
Glu Cys Trp Lys Ser Ser Arg Leu Gly Leu Gly Ala Trp Pro Leu Asp
  50          55          60
Ile Pro Arg Ala Ser Pro Val Leu Pro Ser Pro Arg Thr Thr Gly Pro
  65          70          75          80

```

WO 99/57132

PCT/US99/09970

Leu Ala

**This Page Blank (uspto)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

***This Page Blank (uspto)***